

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

HW 5, NE 548, Spring 2017

Nasser M. Abbasi

December 30, 2019

Contents

0.1	Part a	2
0.2	Part (b)	3
0.3	Part (c)	5
0.4	Part (d)	6

NE548 Problem: Similarity solution for the 1D homogeneous heat equation

Due Thursday April 13, 2017

1. (a) Non-dimensionalize the 1D homogeneous heat equation:

$$\frac{\partial u(x, t)}{\partial t} = \nu \frac{\partial^2 u(x, t)}{\partial x^2} \quad (1)$$

with $-\infty < x < \infty$, and $u(x, t)$ bounded as $x \rightarrow \pm\infty$.

(b) Show that the non-dimensional equations motivate a similarity variable $\xi = x/t^{1/2}$.

(c) Find a similarity solution $u(x, t) = H(\xi)$ by solving the appropriate ODE for $H(\xi)$.

(d) Show that the similarity solution is related to the solution we found in class on 4/6/17 for initial condition

$$u(x, 0) = 0, \quad x < 0 \quad u(x, 0) = C, \quad x > 0.$$

solution

0.1 Part a

Let \bar{x} be the non-dimensional space coordinate and \bar{t} the non-dimensional time coordinate. Therefore we need

$$\begin{aligned} \bar{x} &= \frac{x}{l_0} \\ \bar{t} &= \frac{t}{t_0} \\ \bar{u} &= \frac{u}{u_0} \end{aligned}$$

Where l_0 is the physical characteristic length scale (even if this infinitely long domain, l_0 is given) whose dimensions is $[L]$ and t_0 of dimensions $[T]$ is the characteristic time scale and $\bar{u}(\bar{x}, \bar{t})$ is the new dependent variable, and u_0 characteristic value of u to scale against (typically this is the initial conditions) but this will cancel out. We now rewrite the PDE $\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial x^2}$ in terms of the new dimensionless variables.

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{\partial u}{\partial \bar{u}} \frac{\partial \bar{u}}{\partial \bar{t}} \frac{\partial \bar{t}}{\partial t} \\ &= u_0 \frac{\partial \bar{u}}{\partial \bar{t}} \frac{1}{t_0} \end{aligned} \quad (1)$$

And

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial u}{\partial \bar{u}} \frac{\partial \bar{u}}{\partial \bar{x}} \frac{\partial \bar{x}}{\partial x} \\ &= u_0 \frac{\partial \bar{u}}{\partial \bar{x}} \frac{1}{l_0} \end{aligned}$$

And

$$\frac{\partial^2 u}{\partial x^2} = u_0 \frac{\partial^2 \bar{u}}{\partial \bar{x}^2} \frac{1}{l_0^2} \quad (2)$$

Substituting (1) and (2) into $\frac{\partial u}{\partial t} = v \frac{\partial^2 u}{\partial x^2}$ gives

$$u_0 \frac{\partial \bar{u}}{\partial \bar{t}} \frac{1}{t_0} = v u_0 \frac{\partial^2 \bar{u}}{\partial \bar{x}^2} \frac{1}{l_0^2}$$

$$\frac{\partial \bar{u}}{\partial \bar{t}} = \left(v \frac{t_0}{l_0^2} \right) \frac{\partial^2 \bar{u}}{\partial \bar{x}^2}$$

The above is now non-dimensional. Since v has units $\left[\frac{L^2}{T} \right]$ and $\frac{t_0}{l_0^2}$ also has units $\left[\frac{T}{L^2} \right]$, therefore the product $v \frac{t_0}{l_0^2}$ is non-dimensional quantity.

If we choose t_0 to have same magnitude (not units) as l_0^2 , i.e. $t_0 = l_0^2$, then $\frac{t_0}{l_0^2} = 1$ (with units $\left[\frac{T}{L^2} \right]$) and now we obtain the same PDE as the original, but it is non-dimensional. Where now $\bar{u} \equiv \bar{u}(\bar{t}, \bar{x})$.

0.2 Part (b)

I Will use the Buckingham π theorem for finding expression for the solution in the form $u(x, t) = f(\xi)$ where ξ is the similarity variable. Starting with $\frac{\partial u}{\partial t} = v \frac{\partial^2 u}{\partial x^2}$, in this PDE, the diffusion substance is heat with units of Joule J . Hence the concentration of heat, which is what u represents, will have units of $[u] = \frac{J}{L^3}$. (heat per unit volume). From physics, we expect the solution $u(x, t)$ to depend on x, t, v and initial conditions u_0 as these are the only relevant quantities involved that can affect the diffusion. Therefore, by Buckingham theorem we say

$$u \equiv f(x, t, v, u_0) \quad (1)$$

We have one dependent quantity u and 4 independent quantities. The units of each of the above quantities is

$$[u] = \frac{J}{L^3}$$

$$[x] = L$$

$$[t] = T$$

$$[v] = \frac{L^2}{T}$$

$$[u_0] = \frac{J}{L^3}$$

Hence using Buckingham theorem, we write

$$[u] = [x^a t^b v^c u_0^d] \quad (2)$$

We now determine a, b, c, d , by dimensional analysis. The above is

$$\frac{J}{L^3} = L^a T^b \left(\frac{L^2}{T} \right)^c \left(\frac{J}{L^3} \right)^d$$

$$(J)(L^{-3}) = (L^{a+2c-3d})(T^{b-c})(J^d)$$

Comparing powers of same units on both sides, we see that

$$\begin{aligned}d &= 1 \\b - c &= 0 \\a + 2c - 3d &= -3\end{aligned}$$

From second equation above, $b = c$, hence third equation becomes

$$\begin{aligned}a + 2c - 3d &= -3 \\a + 2c &= 0\end{aligned}$$

Since $d = 1$. Hence

$$\begin{aligned}c &= -\frac{a}{2} \\b &= -\frac{a}{2}\end{aligned}$$

Therefore, now that we found all the powers, (we have one free power a which we can set to any value), then from equation (1)

$$\begin{aligned}[u] &= [x^a t^b v^c u_0^d] \\ \frac{u}{u_0} &= \bar{u} = x^a t^b v^c\end{aligned}$$

Therefore \bar{u} is function of all the variables in the RHS. Let this function be f (This is the same as H in problem statement). Hence the above becomes

$$\begin{aligned}\bar{u} &= f\left(x^a t^{-\frac{a}{2}} v^{-\frac{a}{2}}\right) \\ &= f\left(\frac{x^a}{v^{\frac{a}{2}} t^{\frac{a}{2}}}\right)\end{aligned}$$

Since a is free variable, we can choose

$$a = 1 \tag{2}$$

And obtain

$$\bar{u} \equiv f\left(\frac{x}{\sqrt{vt}}\right) \tag{3}$$

In the above $\frac{x}{\sqrt{vt}}$ is now non-dimensional quantity, which we call, the similarity variable

$$\xi = \frac{x}{\sqrt{vt}} \tag{4}$$

Notice that another choice of a in (2), for example $a = 2$ would lead to $\xi = \frac{x^2}{vt}$ instead of $\xi = \frac{x}{\sqrt{vt}}$ but we will use the latter for the rest of the problem.

0.3 Part (c)

Using $u \equiv f(\xi)$ where $\xi = \frac{x}{\sqrt{vt}}$ then

$$\begin{aligned}\frac{\partial u}{\partial t} &= \frac{df}{d\xi} \frac{\partial \xi}{\partial t} \\ &= \frac{df}{d\xi} \frac{\partial}{\partial t} \left(\frac{x}{\sqrt{vt}} \right) \\ &= -\frac{1}{2} \frac{df}{d\xi} \left(\frac{x}{\sqrt{vt^2}} \right)\end{aligned}$$

And

$$\begin{aligned}\frac{\partial u}{\partial x} &= \frac{df}{d\xi} \frac{\partial \xi}{\partial x} \\ &= \frac{df}{d\xi} \frac{\partial}{\partial x} \left(\frac{x}{\sqrt{vt}} \right) \\ &= \frac{df}{d\xi} \frac{1}{\sqrt{vt}}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial^2 u}{\partial x^2} &= \frac{\partial}{\partial x} \left(\frac{df}{d\xi} \frac{1}{\sqrt{vt}} \right) \\ &= \frac{1}{\sqrt{vt}} \frac{\partial}{\partial x} \left(\frac{df}{d\xi} \right) \\ &= \frac{1}{\sqrt{vt}} \left(\frac{d^2 f}{d\xi^2} \frac{\partial \xi}{\partial x} \right) \\ &= \frac{1}{\sqrt{vt}} \left(\frac{d^2 f}{d\xi^2} \frac{1}{\sqrt{vt}} \right) \\ &= \frac{1}{vt} \frac{d^2 f}{d\xi^2}\end{aligned}$$

Hence the PDE $\frac{\partial u}{\partial t} = v \frac{\partial^2 u}{\partial x^2}$ becomes

$$\begin{aligned}-\frac{1}{2} \frac{df}{d\xi} \left(\frac{x}{\sqrt{vt^2}} \right) &= v \frac{1}{vt} \frac{d^2 f}{d\xi^2} \\ \frac{1}{t} \frac{d^2 f}{d\xi^2} + \frac{1}{2} \frac{x}{\sqrt{vt^2}} \frac{df}{d\xi} &= 0 \\ \frac{d^2 f}{d\xi^2} + \frac{1}{2} \frac{x}{\sqrt{vt}} \frac{df}{d\xi} &= 0\end{aligned}$$

But $\frac{x}{\sqrt{vt}} = \xi$, hence we obtain the required ODE as

$$\begin{aligned}\frac{d^2 f(\xi)}{d\xi^2} + \frac{1}{2} \xi \frac{df(\xi)}{d\xi} &= 0 \\ f'' + \frac{\xi}{2} f' &= 0\end{aligned}$$

We now solve the above ODE for $f(\xi)$. Let $f' = z$, then the ODE becomes

$$z' + \frac{\xi}{2}z = 0$$

Integrating factor is $\mu = e^{\int \frac{\xi}{2} d\xi} = e^{\frac{\xi^2}{4}}$, hence

$$\begin{aligned} \frac{d}{d\xi}(z\mu) &= 0 \\ z\mu &= c_1 \\ z &= c_1 e^{-\frac{\xi^2}{4}} \end{aligned}$$

Therefore, since $f' = z$, then

$$f' = c_1 e^{-\frac{\xi^2}{4}}$$

Integrating gives

$$f(\xi) = c_2 + c_1 \int_0^\xi e^{-\frac{s^2}{4}} ds$$

0.4 Part (d)

For initial conditions of step function

$$u(x, 0) = \begin{cases} 0 & x < 0 \\ C & x > 0 \end{cases}$$

The solution found in class was

$$u(x, t) = \frac{C}{2} + \frac{C}{2} \operatorname{erf}\left(\frac{x}{\sqrt{4vt}}\right) \quad (1)$$

Where $\operatorname{erf}\left(\frac{x}{\sqrt{4vt}}\right) = \frac{2}{\sqrt{\pi}} \int_0^{\frac{x}{\sqrt{4vt}}} e^{-z^2} dz$. The solution found in part (c) earlier is

$$f(\xi) = c_1 \int_0^\xi e^{-\frac{s^2}{4}} ds + c_2$$

Let $s = \sqrt{4}z$, then $\frac{ds}{dz} = \sqrt{4}$, when $s = 0, z = 0$ and when $s = \xi, z = \frac{\xi}{\sqrt{4}}$, therefore the integral becomes

$$f(\xi) = c_1 \sqrt{4} \int_0^{\frac{\xi}{\sqrt{4}}} e^{-z^2} dz + c_2$$

But $\frac{2}{\sqrt{\pi}} \int_0^{\frac{\xi}{\sqrt{4}}} e^{-z^2} dz = \operatorname{erf}\left(\frac{\xi}{\sqrt{4}}\right)$, hence $\int_0^{\frac{\xi}{\sqrt{4}}} e^{-z^2} dz = \frac{\sqrt{\pi}}{2} \operatorname{erf}\left(\frac{\xi}{\sqrt{4}}\right)$ and the above becomes

$$\begin{aligned} f(\xi) &= c_1 \sqrt{\pi} \operatorname{erf}\left(\frac{\xi}{\sqrt{4}}\right) + c_2 \\ &= c_3 \operatorname{erf}\left(\frac{\xi}{\sqrt{4}}\right) + c_2 \end{aligned}$$

Since $\xi = \frac{x}{\sqrt{vt}}$, then above becomes, when converting back to $u(x, t)$

$$u(x, t) = c_3 \operatorname{erf}\left(\frac{x}{\sqrt{4vt}}\right) + c_2 \quad (2)$$

1
2
3
4
5
6 Comparing (1) and (2), we see they are the same. Constants of integration are arbitrary and can be
7 found from initial conditions.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56