

Analytical solution to diffusion-convection PDE in 1D

Nasser M. Abbasi

July 8, 2025

Compiled on July 8, 2025 at 4:52pm

This is a diffusion-convection PDE.

$$\begin{aligned}\frac{\partial F(t, z)}{\partial t} &= k \frac{\partial^2 F(t, z)}{\partial z^2} + v \frac{\partial F(t, z)}{\partial z} \\ t &> 0 \\ 0 &< z < L\end{aligned}\tag{1}$$

Where k is the diffusion constant and v is the convection speed. Boundary conditions are

$$\begin{aligned}F(t, 0) &= 0 \\ F(t, L) &= 1\end{aligned}$$

Initial conditions are

$$F(0, z) = \begin{cases} 0 & 0 \leq z < L \\ 1 & z = L \end{cases}$$

The first step is to convert the PDE to pure diffusion PDE using the transformation

$$F(t, z) = A(t, z) u(t, z)$$

Substituting this back in (1) gives

$$A_t u + A u_t = k(A_{zz} u + 2A_z u_z + A u_{zz}) + v(A_z u + A u_z)$$

Dividing by A and simplifying

$$\begin{aligned}\frac{A_t u}{A} + u_t &= k \left(\frac{A_{zz} u + 2A_z u_z}{A} + u_{zz} \right) + v \left(\frac{A_z u}{A} + u_z \right) \\ u_t &= k u_{zz} + k \frac{A_{zz} - \frac{A_t}{k} + v A_z}{A} u + \left(\frac{2k A_z + v A}{A} \right) u_z\end{aligned}\tag{2}$$

To make (2) pure diffusion PDE, we want

$$k \frac{A_{zz} - \frac{A_t}{k} + v A_z}{A} u = 0\tag{3}$$

$$\left(\frac{2k A_z + v A}{A} \right) u_z = 0\tag{4}$$

From (4) $(2k A_z + v A) u_z = 0$ or $2k A_z + v A = 0$ or $\frac{\partial A}{\partial z} + \frac{v}{2k} A = 0$ which has the solution

$$A(t, z) = C(t) e^{-\frac{v}{2k} z}\tag{5}$$

From (3) we want $k(A_{zz} - \frac{A_t}{k} + vA_z) = 0$. Substituting the result just obtained for $A(t, z)$ in (3) gives

$$\begin{aligned} \frac{v^2}{4k^2}C(t)e^{-\frac{v}{2k}z} - \frac{dC(t)}{dt}\frac{1}{k}e^{-\frac{v}{2k}z} - \frac{v^2}{2k}C(t)e^{-\frac{v}{2k}z} &= 0 \\ \frac{v^2}{4k^2}C(t) - \frac{1}{k}C'(t) - \frac{v^2}{2k}C(t) &= 0 \\ C'(t) + \frac{v^2}{4k}C(t) &= 0 \end{aligned}$$

Hence

$$C(t) = C_1 e^{-\frac{v^2}{4k}t}$$

For some constant C_1 . The constant C_1 ends up canceling out at the very end. Hence we set it to 1 now instead of carrying along in all the derivation in order to simplify notations. Therefore $C(t) = e^{-\frac{v^2}{4k}t}$. Substituting this into (5) gives the transformation function

$$A(t, z) = e^{-\left(\frac{v^2 t}{4k} + \frac{vz}{2k}\right)}$$

Using this in (2) gives the pure diffusion PDE to solve

$$u_t = ku_{zz} \tag{6}$$

Converting the original boundary conditions from F to u gives

$$\begin{aligned} F(t, 0) &= 0 \\ A(t, 0)u(t, 0) &= 0 \\ e^{-\frac{v^2 t}{4k}}u(t, 0) &= 0 \\ u(t, 0) &= 0 \end{aligned}$$

And

$$\begin{aligned} F(t, L) &= 1 \\ A(t, L)u(t, L) &= 1 \\ e^{-\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)}u(t, L) &= 1 \\ u(t, L) &= e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)} \end{aligned}$$

And for the initial conditions

$$\begin{aligned} F(0, z) &= \begin{cases} 0 & 0 \leq z < L \\ 1 & z = L \end{cases} \\ A(0, z)u(0, z) &= \begin{cases} 0 & 0 \leq z < L \\ 1 & z = L \end{cases} \\ e^{-\frac{vz}{2k}}u(0, z) &= \begin{cases} 0 & 0 \leq z < L \\ 1 & z = L \end{cases} \\ u(0, z) &= \begin{cases} 0 & 0 \leq z < L \\ e^{\frac{vL}{2k}} & z = L \end{cases} \end{aligned}$$

Therefore the new PDE to solve is

$$u_t = ku_{zz}$$

With time varying boundary conditions

$$\begin{aligned} u(t, 0) &= 0 \\ u(t, L) &= e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)} \end{aligned}$$

And initial conditions

$$u(0, z) = \begin{cases} 0 & 0 \leq z < L \\ e^{\frac{vL}{2k}} & z = L \end{cases}$$

To solve this using separation of variables, the boundary conditions has to be homogenous. Therefore we use standard method to handle this as follows. Let

$$u(t, z) = \phi(t, z) + u_E(t, z) \quad (7)$$

Where $u_E(t, z)$ is the steady state solution which needs to only satisfy the boundary conditions and $\phi(t, z)$ satisfies the PDE but with homogenous boundary conditions. Therefore

$$\begin{aligned} u_E(t, z) &= u(t, 0) + z \left(\frac{u(t, L) - u(t, 0)}{L} \right) \\ u_E(t, z) &= \frac{z}{L} e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)} \end{aligned}$$

And (7) becomes

$$u(t, z) = \phi(t, z) + \frac{z}{L} e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)}$$

Substituting the above in (6) gives

$$\begin{aligned} \frac{\partial \phi}{\partial t} + \frac{z}{L} \frac{v^2}{4k} e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)} &= k \frac{\partial^2 \phi}{\partial z^2} \\ \frac{\partial \phi}{\partial t} &= k \frac{\partial^2 \phi}{\partial z^2} - \frac{z}{L} \frac{v^2}{4k} e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)} \end{aligned}$$

Or

$$\frac{\partial \phi}{\partial t} = k \frac{\partial^2 \phi}{\partial z^2} + Q(t, z) \quad (8)$$

This is diffusion PDE with homogenous B.C. with source term

$$Q(t, z) = -\frac{d}{dt} u_E(t, z)$$

Now we find $\phi(t, z)$. Since this solution needs to satisfy homogenous boundary conditions, we know the solution to pure diffusion on bounded domain with source present is by given by the following eigenfunction expansion

$$\phi(t, z) = \sum_{n=1}^{\infty} b_n(t) \sin\left(\sqrt{\lambda_n} z\right) \quad (8A)$$

Where eigenvalues are $\lambda_n = \left(\frac{n\pi}{L}\right)^2$ for $n = 1, 2, \dots$ and $\sin(\sqrt{\lambda_n}z)$ are the eigenfunction. Substituting the above in (8) in order to obtain an ODE to solve for $b_n(t)$ gives

$$\sum_{n=1}^{\infty} b'_n(t) \sin(\sqrt{\lambda_n}z) = k \sum_{n=1}^{\infty} -b_n(t) \lambda_n \sin(\sqrt{\lambda_n}z) + Q(t, z) \quad (9)$$

Expanding $Q(t, z)$ in terms of eigenfunctions

$$Q(t, z) = \sum_{n=1}^{\infty} q_n(t) \sin(\sqrt{\lambda_n}z)$$

Applying orthogonality

$$\int_0^L Q(t, z) \sin(\sqrt{\lambda_n}z) dz = q_n(t) \frac{L}{2} \quad (9A)$$

But

$$\begin{aligned} \int_0^a Q(t, z) \sin(\sqrt{\lambda_n}z) dz &= \int_0^L -\frac{z}{L} \frac{v^2}{4k} e^{\left(\frac{v^2t}{4k} + \frac{vL}{2k}\right)} \sin(\sqrt{\lambda_n}z) dz \\ &= -\frac{v^2}{4kL} e^{\left(\frac{v^2t}{4k} + \frac{vL}{2k}\right)} \int_0^L z \sin(\sqrt{\lambda_n}z) dz \\ &= -\frac{v^2}{4kL} e^{\left(\frac{v^2t}{4k} + \frac{vL}{2k}\right)} \frac{(-1)^{n+1} L^2}{n\pi} \\ &= \frac{(-1)^n L v^2 e^{\left(\frac{v^2t}{4k} + \frac{vL}{2k}\right)}}{4kn\pi} \\ &= \frac{(-1)^n v^2 e^{\left(\frac{v^2t}{4k} + \frac{vL}{2k}\right)}}{4k\sqrt{\lambda_n}} \end{aligned}$$

Hence from (9A) we find

$$q_n(t) = \frac{(-1)^n v^2 e^{\left(\frac{v^2t}{4k} + \frac{vL}{2k}\right)}}{2Lk\sqrt{\lambda_n}}$$

Using the above in (9) gives

$$\begin{aligned} \sum_{n=1}^{\infty} b'_n(t) \sin(\sqrt{\lambda_n}z) &= k \sum_{n=1}^{\infty} -b_n(t) \lambda_n \sin(\sqrt{\lambda_n}z) + \sum_{n=1}^{\infty} q_n(t) \sin(\sqrt{\lambda_n}z) \\ b'_n(t) + k\lambda_n b_n(t) &= q_n(t) \end{aligned}$$

To solve the above ODE, the integrating factor is $\mu = e^{k\lambda_n t}$, therefore

$$\begin{aligned} b_n(t) e^{k\lambda_n t} &= \int_0^t q_n(\tau) e^{k\lambda_n \tau} d\tau + C_n \\ b_n(t) &= e^{-k\lambda_n t} \int_0^t q_n(\tau) e^{k\lambda_n \tau} d\tau + C_n e^{-k\lambda_n t} \\ &= \int_0^t q_n(\tau) e^{k\lambda_n(\tau-t)} d\tau + C_n e^{-k\lambda_n t} \end{aligned}$$

Using the above in (7) gives

$$u(t, z) = \frac{z}{L} e^{\left(\frac{v^2t}{4k} + \frac{vL}{2k}\right)} + \sum_{n=1}^{\infty} \left(\int_0^t q_n(\tau) e^{k\lambda_n(\tau-t)} d\tau + C_n e^{-k\lambda_n t} \right) \sin(\sqrt{\lambda_n}z) \quad (10)$$

C_n is now found from initial conditions. At $t = 0$ the above becomes

$$u(0, z) - \frac{z}{L} e^{\frac{vL}{2k}} = \sum_{n=1}^{\infty} C_n \sin(\sqrt{\lambda_n} z)$$

Applying orthogonality

$$\begin{aligned} \int_0^L \left(u(0, z) - \frac{z}{L} e^{\frac{vL}{2k}} \right) \sin(\sqrt{\lambda_m} z) dz &= \int_0^L \sum_{n=1}^{\infty} C_n \sin(\sqrt{\lambda_n} z) \sin(\sqrt{\lambda_m} z) dz \\ \int_0^L u(0, z) \sin(\sqrt{\lambda_n} z) dz - \int_0^L \frac{z}{L} e^{\frac{vL}{2k}} \sin(\sqrt{\lambda_n} z) dz &= C_n \frac{L}{2} \end{aligned}$$

But $\int_0^L u(0, z) \sin(\sqrt{\lambda_n} z) dz = 0$ since $u(0, z)$ is zero everywhere except at the end point. And

$$- \int_0^L \frac{z}{L} e^{\frac{vL}{2k}} \sin(\sqrt{\lambda_n} z) dz = \frac{(-1)^n e^{\frac{vL}{2k}}}{\sqrt{\lambda_n}}$$

Therefore

$$\begin{aligned} \frac{(-1)^n e^{\frac{vL}{2k}}}{\sqrt{\lambda_n}} &= C_n \frac{L}{2} \\ C_n &= \frac{2}{L} \frac{(-1)^n e^{\frac{vL}{2k}}}{\sqrt{\lambda_n}} \end{aligned} \quad (10A)$$

And the solution (10) becomes

$$u(t, z) = \frac{z}{L} e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)} + \sum_{n=1}^{\infty} \left(\int_0^t q_n(\tau) e^{k\lambda_n(\tau-t)} d\tau + C_n e^{-k\lambda_n t} \right) \sin(\sqrt{\lambda_n} z) \quad (11)$$

But

$$\begin{aligned} \int_0^t q_n(\tau) e^{k\lambda_n(\tau-t)} d\tau &= \int_0^t \frac{(-1)^n v^2 e^{\left(\frac{v^2 \tau}{4k} + \frac{vL}{2k}\right)}}{2Lk\sqrt{\lambda_n}} e^{k\lambda_n(\tau-t)} d\tau \\ &= \frac{2(-1)^n v^2 e^{-k\lambda_n t + \frac{vL}{2k}} \left(e^{k\lambda_n t + \frac{tv^2}{4k}} - 1 \right)}{n\pi (4k^2 \lambda_n + v^2)} \end{aligned}$$

Hence (11) becomes

$$u(t, z) = \frac{z}{L} e^{\left(\frac{v^2 t}{4k} + \frac{vL}{2k}\right)} + \sum_{n=1}^{\infty} \left(\frac{2(-1)^n v^2 e^{-k\lambda_n t + \frac{vL}{2k}} \left(e^{k\lambda_n t + \frac{tv^2}{4k}} - 1 \right)}{n\pi (4k^2 \lambda_n + v^2)} + \frac{2}{L} \frac{(-1)^n e^{\frac{vL}{2k}}}{\sqrt{\lambda_n}} e^{-k\lambda_n t} \right) \sin(\sqrt{\lambda_n} z) \quad (12)$$

We now convert back to $F(t, z)$. Since $F(t, z) = A(t, z) u(t, z)$ and $A(t, z) = e^{-\left(\frac{v^2 t}{4} + \frac{vz}{2}\right)}$ then the final solution is

$$F(t, z) = e^{-\left(\frac{v^2 t}{4} + \frac{vz}{2}\right)} u(t, z)$$

The following is animation of the solution for 30 seconds, side-by-side with numerical solution.

The following is the code used

```

ClearAll[t, z, L0, k, lam, n, v, f]
L0 = 10;
v = 1/2;
k = 1/2;
ode = D[f[t, z], t] == k*D[f[t, z], {z, 2}] + v*D[f[t, z], z];
ic = f[0, z] == Piecewise[{{1, z == L0}, {0, True}}];
bc = {f[t, 0] == 0, f[t, L0] == 1};
sol = NDSolve[{ode, bc, ic}, f, {t, 0, 100}, {z, 0, 10}];
(*analytical*)
lam[n_] := (n^2*Pi^2)/L0^2;
max = 100;
u[t_, z_] := Module[{},
  Sum[((2*(-1)^n*v^2*Exp[(L0*v)/(2*k)] - k*lam[n]*t)*
    (Exp[(t*v^2)/(4*k)] + k*lam[n]*t) - 1))/(n*
    Pi*(4*k^2*lam[n] + v^2)) +
    (2/L0)*((( -1)^n*Exp[(L0*v)/(2*k)])/Sqrt[lam[n]])*
    Exp[(-lam[n])*k*t])*
    Sin[Sqrt[lam[n]]*z], {n, 1, max}] + (z/L0)*
    Exp[(v^2*t)/(4*k)] + (v*L0)/(2*k)];

fAnalytical[t_, z_] := Module[{}, Exp[-((v^2*t)/(4*k)] + (v*z)/(2*k))]*u[t, z]];

Manipulate[Grid[{{Row[{"t=", t}], SpanFromLeft},
  {Plot[Evaluate[f[t, z] /. sol], {z, 0, 10},
    PlotRange -> {{0, 10}, {-0.2, 1}},
    ImageSize -> 300, PlotLabel -> "NDSolve solution"},
  Plot[Evaluate[fAnalytical[t, z]], {z, 0, 10},
    PlotRange -> {{0, 10}, {-0.2, 1}},
    ImageSize -> 300, PlotLabel -> "Analytical solution",
    PlotStyle -> Red}}}],

{{t, 0, "t"}, 0, 30, 0.01},
TrackedSymbols :> {t}]

```

References

- Paper "Analytical Solution to the One-Dimensional Advection-Diffusion Equation with Temporally Dependent Coefficients". Dilip Kumar Jaiswal, Atul Kumar, Raja Ram Yadav. Journal of Water Resource and Protection, 2011, 3, 76-84
- Paper "An analytical solution of the diffusion convection equation over a finite domain". Mohammad Farrukh N. Mohsen and Mohammed H. Baluch, Appl. Math. Modelling, 1983, Vol. 7, August 285.
- Lecture 20: Heat conduction with time dependent boundary conditions using Eigenfunction Expansions. Introductory lecture notes on Partial Differential Equations By Anthony Peirce.