

A Solution Manual For

Second order enumerated odes

Nasser M. Abbasi December 31, 2024

Compiled on December 31, 2024 at 1:10pm

Contents

1	Lookup tables for all problems in current book	5
2	Book Solved Problems	11

CHAPTER **1**

LOOKUP TABLES FOR ALL PROBLEMS IN CURRENT
BOOK

1.1	section 1	6
1.2	section 2	7

1.1 section 1

Table 1.1: Lookup table for all problems in current section

ID	problem	ODE
8748	1	$y'' = 0$
8749	2	$y''^2 = 0$
8750	3	$y''^n = 0$
8751	4	$ay'' = 0$
8752	5	$ay''^2 = 0$
8753	6	$ay''^n = 0$
8754	7	$y'' = 1$
8755	8	$y''^2 = 1$
8756	9	$y'' = x$
8757	10	$y''^2 = x$
8758	11	$y''^3 = 0$
8759	12	$y'' + y' = 0$
8760	13	$y''^2 + y' = 0$
8761	14	$y'' + y'^2 = 0$
8762	15	$y'' + y' = 1$
8763	16	$y''^2 + y' = 1$
8764	17	$y'' + y'^2 = 1$
8765	18	$y'' + y' = x$
8766	19	$y''^2 + y' = x$
8767	20	$y'' + y'^2 = x$
8768	21	$y'' + y' + y = 0$
8769	22	$y''^2 + y' + y = 0$
8770	23	$y'' + y'^2 + y = 0$
8771	24	$y'' + y' + y = 1$
8772	25	$y'' + y' + y = x$
8773	26	$y'' + y' + y = 1 + x$
8774	27	$y'' + y' + y = x^2 + x + 1$
8775	28	$y'' + y' + y = x^3 + x^2 + x + 1$
8776	29	$y'' + y' + y = \sin(x)$
8777	30	$y'' + y' + y = \cos(x)$
8778	31	$y'' + y' = 1$
8779	32	$y'' + y' = x$
8780	33	$y'' + y' = 1 + x$
8781	34	$y'' + y' = x^2 + x + 1$

Continued on next page

Table 1.1 Lookup table
Continued from previous page

ID	problem	ODE
8782	35	$y'' + y' = x^3 + x^2 + x + 1$
8783	36	$y'' + y' = \sin(x)$
8784	37	$y'' + y' = \cos(x)$
8785	38	$y'' + y = 1$
8786	39	$y'' + y = x$
8787	40	$y'' + y = 1 + x$
8788	41	$y'' + y = x^2 + x + 1$
8789	42	$y'' + y = x^3 + x^2 + x + 1$
8790	43	$y'' + y = \sin(x)$
8791	44	$y'' + y = \cos(x)$
8792	45	$yy''^2 + y' = 0$
8793	46	$yy''^2 + y'^3 = 0$
8794	47	$y^2y''^2 + y' = 0$
8795	48	$yy''^4 + y'^2 = 0$
8796	49	$y^3y''^2 + yy' = 0$
8797	50	$yy'' + y'^3 = 0$
8798	51	$yy''^3 + y^3y' = 0$
8799	52	$yy''^3 + y^3y'^5 = 0$

1.2 section 2

Table 1.2: Lookup table for all problems in current section

ID	problem	ODE
8800	1	$y'' + xy' + yy'^2 = 0$
8801	2	$y'' + \sin(x)y' + yy'^2 = 0$
8802	3	$y'' + (1-x)y' + y^2y'^2 = 0$
8803	4	$y'' + (\sin(x) + 2x)y' + \cos(y)yy'^2 = 0$
8804	5	$y''y' + y^2 = 0$
8805	6	$y''y' + y^n = 0$
8806	8	$y' = (x+y)^4$
8807	9	$y'' + (x+3)y' + (3+y^2)y'^2 = 0$
8808	10	$y'' + xy' + yy'^2 = 0$
8809	11	$y'' + \sin(x)y' + y'^2 = 0$
8810	12	$3y'' + \cos(x)y' + \sin(y)y'^2 = 0$
8811	13	$10y'' + x^2y' + \frac{3y'^2}{y} = 0$

Continued on next page

Table 1.2 Lookup table

Continued from previous page

ID	problem	ODE
8812	14	$10y'' + (e^x + 3x)y' + \frac{3e^y y'^2}{\sin(y)} = 0$
8813	15	$y'' - \frac{2y}{x^2} = x e^{-\sqrt{x}}$
8814	16	$y'' - \frac{y'}{\sqrt{x}} + \frac{(x+\sqrt{x}-8)y}{4x^2} = x$
8815	17	$y'' + \frac{2y'}{x} + \frac{a^2 y}{x^4} = 0$
8816	18	$(-x^2 + 1)y'' - xy' - c^2 y = 0$
8817	19	$x^6 y'' + 3x^5 y' + a^2 y = \frac{1}{x^2}$
8818	20	$x^2 y'' - 3xy' + 3y = 2x^3 - x^2$
8819	21	$y'' + \cot(x)y' + 4y \csc(x)^2 = 0$
8820	22	$(x^2 + 1)y'' + (1 + x)y' + y = 4 \cos(\ln(1 + x))$
8821	23	$y'' + \tan(x)y' + \cos(x)^2 y = 0$
8822	24	$xy'' - y' + 4x^3 y = 8x^3 \sin(x)^2$
8823	25	$xy'' - y' + 4x^3 y = x^5$
8824	25	$\cos(x)y'' + \sin(x)y' - 2y \cos(x)^3 = 2 \cos(x)^5$
8825	26	$y'' + (1 - \frac{1}{x})y' + 4x^2 y e^{-2x} = 4(x^3 + x^2) e^{-3x}$
8826	27	$y'' - x^2 y' + yx = x^{m+1}$
8827	28	$y'' - \frac{y'}{\sqrt{x}} + \frac{(x+\sqrt{x}-8)y}{4x^2} = 0$
8828	29	$\cos(x)^2 y'' - 2 \cos(x) \sin(x) y' + \cos(x)^2 y = 0$
8829	30	$y'' - 4xy' + (4x^2 - 1)y = -3e^{x^2} \sin(x)$
8830	31	$y'' - 2bxy' + b^2 x^2 y = x$
8831	32	$y'' - 4xy' + (4x^2 - 3)y = e^{x^2}$
8832	33	$y'' - 2 \tan(x)y' + 5y = e^{x^2} \sec(x)$
8833	34	$x^2 y'' - 2xy' + 2(x^2 + 1)y = 0$
8834	35	$4x^2 y'' + 4x^5 y' + (x^8 + 6x^4 + 4)y = 0$
8835	36	$x^2 y'' + (-y + xy')^2 = 0$
8836	37	$xy'' + 2y' - yx = 0$
8837	38	$xy'' + 2y' + yx = 0$
8838	39	$y' + y \cot(x) = 2 \cos(x)$
8839	40	$2xy^2 - y + (y^2 + x + y)y' = 0$
8840	41	$y' = x - y^2$
8841	42	$y'''' - y''' - 3y'' + 5y' - 2y = x e^x + 3 e^{-2x}$
8842	43	$x^2 y'' - x(6 + x)y' + 10y = 0$
8843	44	$x^2 y'' + xy' + (x^2 - 5)y = 0$
8844	45	$x^2 y'' + xy' + (x^2 - 5)y = 0$
8845	46	$x^2 y'' - 4xy' + 6y = 0$

Continued on next page

Table 1.2 Lookup table
Continued from previous page

ID	problem	ODE
8846	47	$y''' - yx = 0$
8847	48	$y' = y^{1/3}$
8848	49	$[x'(t) = 3x(t) + y(t), y'(t) = -x(t) + y(t)]$

CHAPTER 2

BOOK SOLVED PROBLEMS

2.1	section 1	12
2.2	section 2	507

2.1 section 1

2.1.1	problem 1	13
2.1.2	problem 2	25
2.1.3	problem 3	28
2.1.4	problem 4	30
2.1.5	problem 5	42
2.1.6	problem 6	45
2.1.7	problem 7	47
2.1.8	problem 8	62
2.1.9	problem 9	65
2.1.10	problem 10	79
2.1.11	problem 11	82
2.1.12	problem 12	85
2.1.13	problem 13	97
2.1.14	problem 14	101
2.1.15	problem 15	104
2.1.16	problem 16	120
2.1.17	problem 17	124
2.1.18	problem 18	127
2.1.19	problem 19	143
2.1.20	problem 20	146
2.1.21	problem 21	149
2.1.22	problem 22	158
2.1.23	problem 23	160
2.1.24	problem 24	165
2.1.25	problem 25	177
2.1.26	problem 26	189
2.1.27	problem 27	201
2.1.28	problem 28	213
2.1.29	problem 29	225
2.1.30	problem 30	237
2.1.31	problem 31	249
2.1.32	problem 32	265
2.1.33	problem 33	281
2.1.34	problem 34	297
2.1.35	problem 35	313
2.1.36	problem 36	329
2.1.37	problem 37	345
2.1.38	problem 38	361
2.1.39	problem 39	373
2.1.40	problem 40	383
2.1.41	problem 41	393
2.1.42	problem 42	403
2.1.43	problem 43	413
2.1.44	problem 44	424
2.1.45	problem 45	435
2.1.46	problem 46	451
2.1.47	problem 47	459
2.1.48	problem 48	463
2.1.49	problem 49	488
2.1.50	problem 50	493
2.1.51	problem 51	497
2.1.52	problem 52	503

2.1.1 problem 1

Solved as second order ode quadrature	13
Solved as second order linear constant coeff ode	14
Solved as second order linear exact ode	15
Solved as second order missing y ode	16
Solved as second order integrable as is ode	17
Solved as second order integrable as is ode (ABC method)	18
Solved as second order can be made integrable	18
Solved as second order ode using Kovacic algorithm	19
Solved as second order ode adjoint method	22
Maple step by step solution	23
Maple trace	24
Maple dsolve solution	24
Mathematica DSolve solution	24

Internal problem ID [8748]

Book : Second order enumerated odes

Section : section 1

Problem number : 1

Date solved : Thursday, December 12, 2024 at 09:42:48 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y'' = 0$$

Solved as second order ode quadrature

Time used: 0.018 (sec)

Integrating twice gives the solution

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$




Figure 2.1: Slope field plot
 $y'' = 0$

Solved as second order linear constant coeff ode

Time used: 0.046 (sec)

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2 x + c_1$$




Figure 2.2: Slope field plot
 $y'' = 0$

Solved as second order linear exact ode

Time used: 0.050 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$p(x) = 1$$

$$q(x) = 0$$

$$r(x) = 0$$

$$s(x) = 0$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x) y' + (q(x) - p'(x)) y)' = s(x)$$

Integrating gives

$$p(x) y' + (q(x) - p'(x)) y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = c_1$$

We now have a first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1 x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 x + c_2$$




Figure 2.3: Slope field plot
 $y'' = 0$

Solved as second order missing y ode

Time used: 0.039 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_1$$

$$p(x) = c_1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$




Figure 2.4: Slope field plot
 $y'' = 0$

Solved as second order integrable as is ode

Time used: 0.024 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = 0$$

$$y' = c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1 x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 x + c_2$$




Figure 2.5: Slope field plot
 $y'' = 0$

Solved as second order integrable as is ode (ABC method)

Time used: 0.023 (sec)

Writing the ode as

$$y'' = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = 0$$


$$y' = c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1 x + c_2$$

Will add steps showing solving for IC soon.

Figure 2.6: Slope field plot
 $y'' = 0$ **Solved as second order can be made integrable**

Time used: 0.250 (sec)

Multiplying the ode by y' gives

$$y'y'' = 0$$

Integrating the above w.r.t x gives

$$\int y'y'' dx = 0$$

$$\frac{y'^2}{2} = c_1$$

Which is now solved for y . Solving for the derivative gives these ODE's to solve

$$y' = \sqrt{2} \sqrt{c_1} \tag{1}$$

$$y' = -\sqrt{2} \sqrt{c_1} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \sqrt{2} \sqrt{c_1} dx$$

$$y = \sqrt{2} \sqrt{c_1} x + c_2$$

Solving Eq. (2)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\sqrt{2} \sqrt{c_1} dx$$

$$y = -\sqrt{2} \sqrt{c_1} x + c_3$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\sqrt{2} \sqrt{c_1} x + c_3$$

$$y = \sqrt{2} \sqrt{c_1} x + c_2$$




Figure 2.7: Slope field plot
 $y'' = 0$

Solved as second order ode using Kovacic algorithm

Time used: 0.026 (sec)

Writing the ode as

$$y'' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0$$

$$C = 0 \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.1: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2 x + c_1$$




Figure 2.8: Slope field plot
 $y'' = 0$

Solved as second order ode adjoint method

Time used: 0.366 (sec)

In normal form the ode

$$y'' = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$p(x) = 0$$

$$q(x) = 0$$

$$r(x) = 0$$

The Lagrange adjoint ode is given by

$$\xi'' - (\xi p)' + \xi q = 0$$

$$\xi'' - (0)' + (0) = 0$$

$$\xi''(x) = 0$$

Which is solved for $\xi(x)$. Integrating twice gives the solution

$$\xi = c_1x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x)y' - y\xi'(x) + \xi(x)p(x)y &= \int \xi(x)r(x) dx \\ y' + y\left(p(x) - \frac{\xi'(x)}{\xi(x)}\right) &= \frac{\int \xi(x)r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' - \frac{yc_1}{c_1x + c_2} = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{c_1}{c_1x + c_2}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{c_1}{c_1x+c_2} dx} \\ &= \frac{1}{c_1x + c_2} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}\left(\frac{y}{c_1x + c_2}\right) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{c_1x + c_2} &= \int 0 dx + c_3 \\ &= c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1x+c_2}$ gives the final solution

$$y = (c_1x + c_2) c_3$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = (c_1x + c_2) c_3$$

The constants can be merged to give

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$




Figure 2.9: Slope field plot
 $y'' = 0$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence
 $y_2(x) = x$
- General solution of the ODE
 $y(x) = C_1 y_1(x) + C_2 y_2(x)$
- Substitute in solutions
 $y(x) = C_2 x + C_1$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.002 (sec)
Leaf size : 9

```
dsolve(diff(diff(y(x),x),x) = 0,
        y(x),singsol=all)
```

$$y = c_1 x + c_2$$

Mathematica DSolve solution

Solving time : 0.002 (sec)
Leaf size : 12

```
DSolve[{D[y[x],{x,2}]==0,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2 x + c_1$$

2.1.2 problem 2

Solved as second order missing y ode	25
Maple step by step solution	26
Maple trace	27
Maple dsolve solution	27
Mathematica DSolve solution	27

Internal problem ID [8749]

Book : Second order enumerated odes

Section : section 1

Problem number : 2

Date solved : Thursday, December 12, 2024 at 09:42:50 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y''^2 = 0$$

Solved as second order missing y ode

Time used: 0.147 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. Solving for the derivative gives these ODE's to solve

$$p'(x) = 0 \tag{1}$$

$$p'(x) = 0 \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_1$$

$$p(x) = c_1$$

Solving Eq. (2)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_2$$

$$p(x) = c_2$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1x + c_3$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_2$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_2 dx$$

$$y = c_2x + c_4$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_3$$

$$y = c_2x + c_4$$

Maple step by step solution

Let's solve

$$\left(\frac{d^2}{dx^2}y(x)\right)^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2}y(x) = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x)$$

- Substitute in solutions

$$y(x) = C2x + C1$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.024 (sec)

Leaf size : 9

```
dsolve(diff(diff(y(x),x),x)^2 = 0,
        y(x),singsol=all)
```

$$y = c_1x + c_2$$

Mathematica DSolve solution

Solving time : 0.002 (sec)

Leaf size : 12

```
DSolve[{(D[y[x],{x,2}])^2==0,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2x + c_1$$

2.1.3 problem 3

Solved as second order missing y ode	28
Maple step by step solution	29
Maple trace	29
Maple dsolve solution	29
Mathematica DSolve solution	29

Internal problem ID [8750]

Book : Second order enumerated odes

Section : section 1

Problem number : 3

Date solved : Thursday, December 12, 2024 at 09:42:50 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y'''' = 0$$

Solved as second order missing y ode

Time used: 0.078 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^n = 0$$

Which is now solve for $p(x)$ as first order ode. Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_1$$

$$p(x) = c_1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$

Maple step by step solution

Let's solve

$$\left(\frac{d^2}{dx^2}y(x)\right)^n = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2}y(x) = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x)$$

- Substitute in solutions

$$y(x) = C_2 x + C_1$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.007 (sec)

Leaf size : 9

```
dsolve(diff(diff(y(x),x),x)^n = 0,
        y(x),singsol=all)
```

$$y = c_1 x + c_2$$

Mathematica DSolve solution

Solving time : 0.003 (sec)

Leaf size : 24

```
DSolve[{(D[y[x],{x,2}])^n==0,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{2} 0^{\frac{1}{n}} x^2 + c_2 x + c_1$$

2.1.4 problem 4

Solved as second order ode quadrature	30
Solved as second order linear constant coeff ode	31
Solved as second order linear exact ode	32
Solved as second order missing y ode	33
Solved as second order integrable as is ode	34
Solved as second order integrable as is ode (ABC method)	35
Solved as second order can be made integrable	35
Solved as second order ode using Kovacic algorithm	36
Solved as second order ode adjoint method	39
Maple step by step solution	40
Maple trace	41
Maple dsolve solution	41
Mathematica DSolve solution	41

Internal problem ID [8751]

Book : Second order enumerated odes

Section : section 1

Problem number : 4

Date solved : Thursday, December 12, 2024 at 09:42:51 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$ay'' = 0$$

Solved as second order ode quadrature

Time used: 0.021 (sec)

The ODE simplifies to

$$y'' = 0$$

Integrating twice gives the solution

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$




Figure 2.10: Slope field plot
 $ay'' = 0$

Solved as second order linear constant coeff ode

Time used: 0.025 (sec)

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = a, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$a \lambda^2 e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$a \lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = a, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(a)} \pm \frac{1}{(2)(a)} \sqrt{(0)^2 - (4)(a)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2 x + c_1$$




Figure 2.11: Slope field plot
 $ay'' = 0$

Solved as second order linear exact ode

Time used: 0.053 (sec)

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$p(x) = a$$

$$q(x) = 0$$

$$r(x) = 0$$

$$s(x) = 0$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$ay' = c_1$$

We now have a first order ode to solve which is

$$ay' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{c_1}{a} dx$$

$$y = \frac{c_1 x}{a} + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 x}{a} + c_2$$




Figure 2.12: Slope field plot
 $ay'' = 0$

Solved as second order missing y ode

Time used: 0.045 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$ap'(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_1$$

$$p(x) = c_1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$




Figure 2.13: Slope field plot
 $ay'' = 0$

Solved as second order integrable as is ode

Time used: 0.030 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int ay'' dx = 0$$

$$ay' = c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{c_1}{a} dx$$

$$y = \frac{c_1 x}{a} + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 x}{a} + c_2$$




Figure 2.14: Slope field plot
 $ay'' = 0$

Solved as second order integrable as is ode (ABC method)

Time used: 0.030 (sec)

Writing the ode as

$$ay'' = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int ay'' dx = 0$$


$$ay' = c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{c_1}{a} dx$$

$$y = \frac{c_1 x}{a} + c_2$$

Will add steps showing solving for IC soon.

Figure 2.15: Slope field plot
 $ay'' = 0$ **Solved as second order can be made integrable**

Time used: 0.270 (sec)

Multiplying the ode by y' gives

$$ay'y'' = 0$$

Integrating the above w.r.t x gives

$$\int ay'y'' dx = 0$$

$$\frac{ay'^2}{2} = c_1$$

Which is now solved for y . Solving for the derivative gives these ODE's to solve

$$y' = \frac{\sqrt{2} \sqrt{ac_1}}{a} \tag{1}$$

$$y' = -\frac{\sqrt{2} \sqrt{ac_1}}{a} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{\sqrt{2} \sqrt{ac_1}}{a} dx$$

$$y = \frac{\sqrt{2} \sqrt{ac_1} x}{a} + c_2$$

Solving Eq. (2)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\frac{\sqrt{2} \sqrt{ac_1}}{a} dx$$

$$y = -\frac{\sqrt{2} \sqrt{ac_1} x}{a} + c_3$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\sqrt{2} \sqrt{ac_1} x}{a} + c_3$$

$$y = \frac{\sqrt{2} \sqrt{ac_1} x}{a} + c_2$$




Figure 2.16: Slope field plot
 $ay'' = 0$

Solved as second order ode using Kovacic algorithm

Time used: 0.029 (sec)

Writing the ode as

$$ay'' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = a$$

$$B = 0$$

$$C = 0 \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.5: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2 x + c_1$$




Figure 2.17: Slope field plot
 $ay'' = 0$

Solved as second order ode adjoint method

Time used: 0.306 (sec)

In normal form the ode

$$ay'' = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$p(x) = 0$$

$$q(x) = 0$$

$$r(x) = 0$$

The Lagrange adjoint ode is given by

$$\xi'' - (\xi p)' + \xi q = 0$$

$$\xi'' - (0)' + (0) = 0$$

$$\xi''(x) = 0$$

Which is solved for $\xi(x)$. Integrating twice gives the solution

$$\xi = c_1x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x)y' - y\xi'(x) + \xi(x)p(x)y &= \int \xi(x)r(x) dx \\ y' + y\left(p(x) - \frac{\xi'(x)}{\xi(x)}\right) &= \frac{\int \xi(x)r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' - \frac{yc_1}{c_1x + c_2} = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{c_1}{c_1x + c_2}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{c_1}{c_1x + c_2} dx} \\ &= \frac{1}{c_1x + c_2} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{y}{c_1x + c_2} \right) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{c_1x + c_2} &= \int 0 dx + c_3 \\ &= c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1x+c_2}$ gives the final solution

$$y = (c_1x + c_2) c_3$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = (c_1x + c_2) c_3$$

The constants can be merged to give

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$




Figure 2.18: Slope field plot
 $ay'' = 0$

Maple step by step solution

Let's solve

$$a\left(\frac{d^2}{dx^2}y(x)\right) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2}y(x) = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE
 $y_1(x) = 1$
- Repeated root, multiply $y_1(x)$ by x to ensure linear independence
 $y_2(x) = x$
- General solution of the ODE
 $y(x) = C_1 y_1(x) + C_2 y_2(x)$
- Substitute in solutions
 $y(x) = C_2 x + C_1$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.002 (sec)
Leaf size : 9

```
dsolve(a*diff(diff(y(x),x),x) = 0,
       y(x),singsol=all)
```

$$y = c_1 x + c_2$$

Mathematica DSolve solution

Solving time : 0.002 (sec)
Leaf size : 12

```
DSolve[{a*D[y[x],{x,2}]==0,{}},
       y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2 x + c_1$$

2.1.5 problem 5

Solved as second order missing y ode	42
Maple step by step solution	43
Maple trace	44
Maple dsolve solution	44
Mathematica DSolve solution	44

Internal problem ID [8752]

Book : Second order enumerated odes

Section : section 1

Problem number : 5

Date solved : Thursday, December 12, 2024 at 09:42:52 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$ay''^2 = 0$$

Solved as second order missing y ode

Time used: 0.187 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$ap'(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. Solving for the derivative gives these ODE's to solve

$$p'(x) = 0 \tag{1}$$

$$p'(x) = 0 \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_1$$

$$p(x) = c_1$$

Solving Eq. (2)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_2$$

$$p(x) = c_2$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1x + c_3$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_2$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_2 dx$$

$$y = c_2x + c_4$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_3$$

$$y = c_2x + c_4$$

Maple step by step solution

Let's solve

$$a \left(\frac{d^2}{dx^2} y(x) \right)^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x)$$

- Substitute in solutions

$$y(x) = C2x + C1$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.029 (sec)

Leaf size : 9

```
dsolve(a*diff(diff(y(x),x),x)^2 = 0,  
       y(x),singsol=all)
```

$$y = c_1x + c_2$$

Mathematica DSolve solution

Solving time : 0.002 (sec)

Leaf size : 12

```
DSolve[{a*(D[y[x],{x,2}])^2==0,{}},  
       y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2x + c_1$$

2.1.6 problem 6

Solved as second order missing y ode	45
Maple step by step solution	46
Maple trace	46
Maple dsolve solution	46
Mathematica DSolve solution	46

Internal problem ID [8753]

Book : Second order enumerated odes

Section : section 1

Problem number : 6

Date solved : Thursday, December 12, 2024 at 09:42:53 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$ay''^m = 0$$

Solved as second order missing y ode

Time used: 0.093 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$ap'(x)^n = 0$$

Which is now solve for $p(x)$ as first order ode. Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_1$$

$$p(x) = c_1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_2$$

Maple step by step solution

Let's solve

$$a \left(\frac{d^2}{dx^2} y(x) \right)^n = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x)$$

- Substitute in solutions

$$y(x) = C_2 x + C_1$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.006 (sec)

Leaf size : 9

```
dsolve(a*diff(diff(y(x),x),x)^n = 0,
       y(x),singsol=all)
```

$$y = c_1 x + c_2$$

Mathematica DSolve solution

Solving time : 0.003 (sec)

Leaf size : 24

```
DSolve[{a*(D[y[x],{x,2}])^n==0,{}},
       y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{2} 0^{\frac{1}{n}} x^2 + c_2 x + c_1$$

2.1.7 problem 7

Solved as second order ode quadrature	47
Solved as second order linear constant coeff ode	48
Solved as second order linear exact ode	50
Solved as second order missing y ode	51
Solved as second order integrable as is ode	52
Solved as second order integrable as is ode (ABC method)	53
Solved as second order can be made integrable	53
Solved as second order ode using Kovacic algorithm	55
Solved as second order ode adjoint method	58
Maple step by step solution	60
Maple trace	61
Maple dsolve solution	61
Mathematica DSolve solution	61

Internal problem ID [8754]

Book : Second order enumerated odes

Section : section 1

Problem number : 7

Date solved : Thursday, December 12, 2024 at 09:42:54 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y'' = 1$$

Solved as second order ode quadrature

Time used: 0.026 (sec)

The ODE can be written as

$$y'' = 1$$

Integrating once gives

$$y' = x + c_1$$

Integrating again gives

$$y = \frac{x^2}{2} + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{2}x^2 + c_1x + c_2$$




Figure 2.19: Slope field plot
 $y'' = 1$

Solved as second order linear constant coeff ode

Time used: 0.086 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 0, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \tag{1}$$

Therefore the homogeneous solution y_h is

$$y_h = c_2x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1x^2$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{x^2}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{2}x^2 + c_2x + c_1$$




Figure 2.20: Slope field plot
 $y'' = 1$

Solved as second order linear exact ode

Time used: 0.053 (sec)

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= 1 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = \int 1 dx$$

We now have a first order ode to solve which is

$$y' = x + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int x + c_1 dx$$

$$y = \frac{1}{2}x^2 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{2}x^2 + c_1x + c_2$$




Figure 2.21: Slope field plot
 $y'' = 1$

Solved as second order missing y ode

Time used: 0.043 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 1 dx$$

$$p(x) = x + c_1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int x + c_1 dx$$

$$y = \frac{1}{2}x^2 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{2}x^2 + c_1x + c_2$$




Figure 2.22: Slope field plot
 $y'' = 1$

Solved as second order integrable as is ode

Time used: 0.030 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int 1 dx$$

$$y' = x + c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int x + c_1 dx$$

$$y = \frac{1}{2}x^2 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{2}x^2 + c_1x + c_2$$




Figure 2.23: Slope field plot
 $y'' = 1$

Solved as second order integrable as is ode (ABC method)

Time used: 0.029 (sec)

Writing the ode as

$$y'' = 1$$

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int 1 dx$$


$$y' = x + c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int x + c_1 dx$$

$$y = \frac{1}{2}x^2 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Figure 2.24: Slope field plot
 $y'' = 1$ **Solved as second order can be made integrable**

Time used: 0.421 (sec)

Multiplying the ode by y' gives

$$y'y'' - y' = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' - y') dx = 0$$

$$\frac{y'^2}{2} - y = c_1$$

Which is now solved for y . Solving for the derivative gives these ODE's to solve

$$y' = \sqrt{2y + 2c_1} \tag{1}$$

$$y' = -\sqrt{2y + 2c_1} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Integrating gives

$$\int \frac{1}{\sqrt{2y + 2c_1}} dy = dx$$

$$\sqrt{2y + 2c_1} = x + c_2$$

Singular solutions are found by solving

$$\sqrt{2y + 2c_1} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -c_1$$

Solving for y gives

$$y = \frac{1}{2}c_2^2 + c_2x + \frac{1}{2}x^2 - c_1$$

Solving Eq. (2)

Integrating gives

$$\int -\frac{1}{\sqrt{2y + 2c_1}} dy = dx$$

$$-\sqrt{2y + 2c_1} = x + c_3$$

Singular solutions are found by solving

$$-\sqrt{2y + 2c_1} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -c_1$$

Solving for y gives

$$y = \frac{1}{2}c_3^2 + c_3x + \frac{1}{2}x^2 - c_1$$

Will add steps showing solving for IC soon.

The solution

$$y = -c_1$$

was found not to satisfy the ode or the IC. Hence it is removed.

Summary of solutions found

$$y = \frac{1}{2}c_2^2 + c_2x + \frac{1}{2}x^2 - c_1$$

$$y = \frac{1}{2}c_3^2 + c_3x + \frac{1}{2}x^2 - c_1$$




Figure 2.25: Slope field plot
 $y'' = 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.046 (sec)

Writing the ode as

$$y'' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.9: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{x^2}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{2}x^2 + c_2x + c_1$$




Figure 2.26: Slope field plot
 $y'' = 1$

Solved as second order ode adjoint method

Time used: 0.421 (sec)

In normal form the ode

$$y'' = 1 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$p(x) = 0$$

$$q(x) = 0$$

$$r(x) = 1$$

The Lagrange adjoint ode is given by

$$\begin{aligned}\xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + (0) &= 0 \\ \xi''(x) &= 0\end{aligned}$$

Which is solved for $\xi(x)$. Integrating twice gives the solution

$$\xi = c_1x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y\xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y\left(p(x) - \frac{\xi'(x)}{\xi(x)}\right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' - \frac{yc_1}{c_1x + c_2} = \frac{\frac{1}{2}c_1x^2 + c_2x}{c_1x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{c_1}{c_1x + c_2} \\ p(x) &= \frac{x(c_1x + 2c_2)}{2c_1x + 2c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{c_1}{c_1x + c_2} dx} \\ &= \frac{1}{c_1x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{x(c_1x + 2c_2)}{2c_1x + 2c_2} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1x + c_2} \right) &= \left(\frac{1}{c_1x + c_2} \right) \left(\frac{x(c_1x + 2c_2)}{2c_1x + 2c_2} \right) \\ d \left(\frac{y}{c_1x + c_2} \right) &= \left(\frac{x(c_1x + 2c_2)}{(2c_1x + 2c_2)(c_1x + c_2)} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{c_1x + c_2} &= \int \frac{x(c_1x + 2c_2)}{(2c_1x + 2c_2)(c_1x + c_2)} dx \\ &= \frac{x}{2c_1} + \frac{c_2^2}{2c_1^2(c_1x + c_2)} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1x+c_2}$ gives the final solution

$$y = \frac{2c_1^3c_3x + (2c_2c_3 + x^2)c_1^2 + c_1c_2x + c_2^2}{2c_1^2}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{2c_1^3c_3x + (2c_2c_3 + x^2)c_1^2 + c_1c_2x + c_2^2}{2c_1^2}$$

The constants can be merged to give

$$y = \frac{2c_1^3x + (x^2 + 2c_2)c_1^2 + c_1c_2x + c_2^2}{2c_1^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{2c_1^3x + (x^2 + 2c_2)c_1^2 + c_1c_2x + c_2^2}{2c_1^2}$$




Figure 2.27: Slope field plot
 $y'' = 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) = 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

- $y_2(x) = x$
- General solution of the ODE
 $y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$
 - Substitute in solutions of the homogeneous ODE
 $y(x) = C_1 + C_2 x + y_p(x)$
 - Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 1 \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$
 - Compute Wronskian
 $W(y_1(x), y_2(x)) = 1$
 - Substitute functions into equation for $y_p(x)$
 $y_p(x) = -\left(\int x dx \right) + \left(\int 1 dx \right) x$
 - Compute integrals
 $y_p(x) = \frac{x^2}{2}$
 - Substitute particular solution into general solution to ODE
 $y(x) = C_1 + C_2 x + \frac{1}{2} x^2$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.002 (sec)
 Leaf size : 14

```
dsolve(diff(diff(y(x),x),x) = 1,
        y(x),singsol=all)
```

$$y = \frac{1}{2}x^2 + c_1x + c_2$$

Mathematica DSolve solution

Solving time : 0.002 (sec)
 Leaf size : 19

```
DSolve[{D[y[x],{x,2}]==1,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{x^2}{2} + c_2x + c_1$$

2.1.8 problem 8

Solved as second order missing y ode	62
Maple step by step solution	63
Maple trace	63
Maple dsolve solution	63
Mathematica DSolve solution	64

Internal problem ID [8755]

Book : Second order enumerated odes

Section : section 1

Problem number : 8

Date solved : Thursday, December 12, 2024 at 09:42:56 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y''^2 = 1$$

Solved as second order missing y ode

Time used: 0.201 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^2 - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Solving for the derivative gives these ODE's to solve

$$p'(x) = 1 \tag{1}$$

$$p'(x) = -1 \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 1 dx$$

$$p(x) = x + c_1$$

Solving Eq. (2)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int -1 dx$$

$$p(x) = -x + c_2$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -x + c_2$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -x + c_2 dx$$

$$y = -\frac{1}{2}x^2 + c_2x + c_3$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int x + c_1 dx$$

$$y = \frac{1}{2}x^2 + c_1x + c_4$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{1}{2}x^2 + c_2x + c_3$$

$$y = \frac{1}{2}x^2 + c_1x + c_4$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful
Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.005 (sec)

Leaf size : 27

```
dsolve(diff(diff(y(x),x),x)^2 = 1,
        y(x),singsol=all)
```

$$y = \frac{1}{2}x^2 + c_1x + c_2$$

$$y = -\frac{1}{2}x^2 + c_1x + c_2$$

Mathematica DSolve solution

Solving time : 0.003 (sec)

Leaf size : 37

```
DSolve[{(D[y[x],{x,2}])^2==1,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{x^2}{2} + c_2x + c_1$$

$$y(x) \rightarrow \frac{x^2}{2} + c_2x + c_1$$

2.1.9 problem 9

Solved as second order ode quadrature	65
Solved as second order linear constant coeff ode	66
Solved as second order linear exact ode	68
Solved as second order missing y ode	69
Solved as second order integrable as is ode	70
Solved as second order integrable as is ode (ABC method)	71
Solved as second order ode using Kovacic algorithm	71
Solved as second order ode adjoint method	75
Maple step by step solution	77
Maple trace	78
Maple dsolve solution	78
Mathematica DSolve solution	78

Internal problem ID [8756]

Book : Second order enumerated odes

Section : section 1

Problem number : 9

Date solved : Thursday, December 12, 2024 at 09:42:56 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y'' = x$$

Solved as second order ode quadrature

Time used: 0.026 (sec)

Integrating once gives

$$y' = \frac{x^2}{2} + c_1$$

Integrating again gives

$$y = \frac{x^3}{6} + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{6}x^3 + c_1x + c_2$$




Figure 2.28: Slope field plot
 $y'' = x$

Solved as second order linear constant coeff ode

Time used: 0.098 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 0, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6xA_2 + 2A_1 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^3}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{x^3}{6} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{6}x^3 + c_2x + c_1$$




Figure 2.29: Slope field plot
 $y'' = x$

Solved as second order linear exact ode

Time used: 0.056 (sec)

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$p(x) = 1$$

$$q(x) = 0$$

$$r(x) = 0$$

$$s(x) = x$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = \int x dx$$

We now have a first order ode to solve which is

$$y' = \frac{x^2}{2} + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{x^2}{2} + c_1 dx$$

$$y = \frac{1}{6}x^3 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{6}x^3 + c_1x + c_2$$




Figure 2.30: Slope field plot
 $y'' = x$

Solved as second order missing y ode

Time used: 0.052 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - x = 0$$

Which is now solve for $p(x)$ as first order ode. Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int x dx$$

$$p(x) = \frac{x^2}{2} + c_1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{x^2}{2} + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{x^2}{2} + c_1 dx$$

$$y = \frac{1}{6}x^3 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{6}x^3 + c_1x + c_2$$




Figure 2.31: Slope field plot
 $y'' = x$

Solved as second order integrable as is ode

Time used: 0.034 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int x dx$$

$$y' = \frac{x^2}{2} + c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{x^2}{2} + c_1 dx$$

$$y = \frac{1}{6}x^3 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{6}x^3 + c_1x + c_2$$




Figure 2.32: Slope field plot
 $y'' = x$

Solved as second order integrable as is ode (ABC method)

Time used: 0.036 (sec)

Writing the ode as

$$y'' = x$$

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int x dx$$


$$y' = \frac{x^2}{2} + c_1$$

Which is now solved for y . Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{x^2}{2} + c_1 dx$$

$$y = \frac{1}{6}x^3 + c_1x + c_2$$

Will add steps showing solving for IC soon.

Figure 2.33: Slope field plot
 $y'' = x$ **Solved as second order ode using Kovacic algorithm**

Time used: 0.140 (sec)

Writing the ode as

$$y'' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0$$

$$C = 0 \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.11: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6xA_2 + 2A_1 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^3}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{x^3}{6} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{6}x^3 + c_2x + c_1$$




Figure 2.34: Slope field plot
 $y'' = x$

Solved as second order ode adjoint method

Time used: 0.351 (sec)

In normal form the ode

$$y'' = x \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$p(x) = 0$$

$$q(x) = 0$$

$$r(x) = x$$

The Lagrange adjoint ode is given by

$$\xi'' - (\xi p)' + \xi q = 0$$

$$\xi'' - (0)' + (0) = 0$$

$$\xi''(x) = 0$$

Which is solved for $\xi(x)$. Integrating twice gives the solution

$$\xi = c_1x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\xi(x)y' - y\xi'(x) + \xi(x)p(x)y = \int \xi(x)r(x)dx$$

$$y' + y\left(p(x) - \frac{\xi'(x)}{\xi(x)}\right) = \frac{\int \xi(x)r(x)dx}{\xi(x)}$$

Or

$$y' - \frac{yc_1}{c_1x + c_2} = \frac{\frac{1}{3}c_1x^3 + \frac{1}{2}c_2x^2}{c_1x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{c_1}{c_1x + c_2}$$

$$p(x) = \frac{2c_1x^3 + 3c_2x^2}{6c_1x + 6c_2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{c_1}{c_1x+c_2} dx} \\ &= \frac{1}{c_1x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{2c_1x^3 + 3c_2x^2}{6c_1x + 6c_2} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1x + c_2} \right) &= \left(\frac{1}{c_1x + c_2} \right) \left(\frac{2c_1x^3 + 3c_2x^2}{6c_1x + 6c_2} \right) \\ d \left(\frac{y}{c_1x + c_2} \right) &= \left(\frac{2c_1x^3 + 3c_2x^2}{(6c_1x + 6c_2)(c_1x + c_2)} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{c_1x + c_2} &= \int \frac{2c_1x^3 + 3c_2x^2}{(6c_1x + 6c_2)(c_1x + c_2)} dx \\ &= \frac{c_1x^2 - c_2x}{6c_1^2} - \frac{c_2^3}{6c_1^3(c_1x + c_2)} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1x+c_2}$ gives the final solution

$$y = \frac{6c_1^4c_3x + (x^3 + 6c_2c_3)c_1^3 - c_1c_2^2x - c_2^3}{6c_1^3}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{6c_1^4c_3x + (x^3 + 6c_2c_3)c_1^3 - c_1c_2^2x - c_2^3}{6c_1^3}$$

The constants can be merged to give

$$y = \frac{6c_1^4x + (x^3 + 6c_2)c_1^3 - c_1c_2^2x - c_2^3}{6c_1^3}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{6c_1^4x + (x^3 + 6c_2)c_1^3 - c_1c_2^2x - c_2^3}{6c_1^3}$$




Figure 2.35: Slope field plot
 $y'' = x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) = x$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 + C_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\left(\int x^2 dx \right) + \left(\int x dx \right) x$$

- Compute integrals

$$y_p(x) = \frac{x^3}{6}$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 + C_2x + \frac{1}{6}x^3$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.003 (sec)
Leaf size : 14

```
dsolve(diff(diff(y(x),x),x) = x,
        y(x),singsol=all)
```

$$y = \frac{1}{6}x^3 + c_1x + c_2$$

Mathematica DSolve solution

Solving time : 0.002 (sec)
Leaf size : 19

```
DSolve[{D[y[x],{x,2}]==x,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{x^3}{6} + c_2x + c_1$$

2.1.10 problem 10

Solved as second order missing y ode	79
Maple step by step solution	80
Maple trace	80
Maple dsolve solution	81
Mathematica DSolve solution	81

Internal problem ID [8757]

Book : Second order enumerated odes

Section : section 1

Problem number : 10

Date solved : Thursday, December 12, 2024 at 09:42:58 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y''^2 = x$$

Solved as second order missing y ode

Time used: 0.272 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^2 - x = 0$$

Which is now solve for $p(x)$ as first order ode. Solving for the derivative gives these ODE's to solve

$$p'(x) = \sqrt{x} \tag{1}$$

$$p'(x) = -\sqrt{x} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int \sqrt{x} dx$$

$$p(x) = \frac{2x^{3/2}}{3} + c_1$$

Solving Eq. (2)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int -\sqrt{x} dx$$

$$p(x) = -\frac{2x^{3/2}}{3} + c_2$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{2x^{3/2}}{3} + c_2$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\frac{2x^{3/2}}{3} + c_2 dx$$

$$y = c_2x - \frac{4x^{5/2}}{15} + c_3$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{2x^{3/2}}{3} + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{2x^{3/2}}{3} + c_1 dx$$

$$y = c_1x + \frac{4x^{5/2}}{15} + c_4$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + \frac{4x^{5/2}}{15} + c_4$$

$$y = c_2x - \frac{4x^{5/2}}{15} + c_3$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each resultin
*** Sublevel 2 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful
-----
* Tackling next ODE.
*** Sublevel 2 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```


Maple dsolve solution

Solving time : 0.031 (sec)

Leaf size : 27

```
dsolve(diff(diff(y(x),x),x)^2 = x,
        y(x),singsol=all)
```

$$y = \frac{4x^{5/2}}{15} + c_1x + c_2$$

$$y = -\frac{4x^{5/2}}{15} + c_1x + c_2$$

Mathematica DSolve solution

Solving time : 0.004 (sec)

Leaf size : 41

```
DSolve[{(D[y[x],{x,2}])^2==x,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{4x^{5/2}}{15} + c_2x + c_1$$

$$y(x) \rightarrow \frac{4x^{5/2}}{15} + c_2x + c_1$$

2.1.11 problem 11

Solved as second order missing y ode	82
Maple step by step solution	83
Maple trace	84
Maple dsolve solution	84
Mathematica DSolve solution	84

Internal problem ID [8758]

Book : Second order enumerated odes

Section : section 1

Problem number : 11

Date solved : Thursday, December 12, 2024 at 09:42:58 AM

CAS classification : [[_2nd_order, _quadrature]]

Solve

$$y''^3 = 0$$

Solved as second order missing y ode

Time used: 0.179 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^3 = 0$$

Which is now solve for $p(x)$ as first order ode. Solving for the derivative gives these ODE's to solve

$$p'(x) = 0 \tag{1}$$

$$p'(x) = 0 \tag{2}$$

$$p'(x) = 0 \tag{3}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_1$$

$$p(x) = c_1$$

Solving Eq. (2)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_2$$

$$p(x) = c_2$$

Solving Eq. (3)

Since the ode has the form $p'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\int dp = \int 0 dx + c_3$$

$$p(x) = c_3$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_1 dx$$

$$y = c_1x + c_4$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_2$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_2 dx$$

$$y = c_2x + c_5$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_3$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int c_3 dx$$

$$y = c_3x + c_6$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + c_4$$

$$y = c_2x + c_5$$

$$y = c_3x + c_6$$

Maple step by step solution

Let's solve

$$\left(\frac{d^2}{dx^2}y(x)\right)^3 = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2}y(x) = 0$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$
- Roots of the characteristic polynomial

$$r = 0$$
- 1st solution of the ODE

$$y_1(x) = 1$$
- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$
- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x)$$
- Substitute in solutions

$$y(x) = C_2 x + C_1$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

Maple dsolve solution

Solving time : 0.024 (sec)
 Leaf size : 9

```
dsolve(diff(diff(y(x),x),x)^3 = 0,
        y(x),singsol=all)
```

$$y = c_1 x + c_2$$

Mathematica DSolve solution

Solving time : 0.002 (sec)
 Leaf size : 12

```
DSolve[{(D[y[x],{x,2}])^3==0,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2 x + c_1$$

2.1.12 problem 12

Solved as second order linear constant coeff ode	85
Solved as second order linear exact ode	86
Solved as second order missing y ode	88
Solved as second order integrable as is ode	89
Solved as second order integrable as is ode (ABC method)	90
Solved as second order ode using Kovacic algorithm	91
Solved as second order ode adjoint method	93
Maple step by step solution	96
Maple trace	96
Maple dsolve solution	96
Mathematica DSolve solution	96

Internal problem ID [8759]

Book : Second order enumerated odes

Section : section 1

Problem number : 12

Date solved : Thursday, December 12, 2024 at 09:42:59 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y'' + y' = 0$$

Solved as second order linear constant coeff ode

Time used: 0.069 (sec)

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\lambda_1 = 0$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(0)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 + c_2 e^{-x}$$




Figure 2.36: Slope field plot
 $y'' + y' = 0$

Solved as second order linear exact ode

Time used: 0.135 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$p(x) = 1$$

$$q(x) = 1$$

$$r(x) = 0$$

$$s(x) = 0$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = c_1$$

We now have a first order ode to solve which is

$$y' + y = c_1$$

Integrating gives

$$\int \frac{1}{-y + c_1} dy = dx$$

$$-\ln(-y + c_1) = x + c_2$$

Singular solutions are found by solving

$$-y + c_1 = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = c_1$$

Solving for y gives

$$y = -e^{-x-c_2} + c_1$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1$$

$$y = -e^{-x-c_2} + c_1$$




Figure 2.37: Slope field plot

$$y'' + y' = 0$$

Solved as second order missing y ode

Time used: 0.099 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating gives

$$\int -\frac{1}{p} dp = dx$$

$$-\ln(p) = x + c_1$$

Singular solutions are found by solving

$$-p = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 0$$

Solving for $p(x)$ gives

$$p(x) = e^{-x-c_1}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 0$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_2$$

$$y = c_2$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = e^{-x-c_1}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int e^{-x-c_1} dx$$

$$y = -e^{-x-c_1} + c_3$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2$$

$$y = -e^{-x-c_1} + c_3$$




Figure 2.38: Slope field plot
 $y'' + y' = 0$

Solved as second order integrable as is ode

Time used: 0.106 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = 0$$

$$y' + y = c_1$$

Which is now solved for y . Integrating gives

$$\int \frac{1}{-y + c_1} dy = dx$$

$$-\ln(-y + c_1) = x + c_2$$

Singular solutions are found by solving

$$-y + c_1 = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = c_1$$

Solving for y gives

$$y = -e^{-x-c_2} + c_1$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1$$

$$y = -e^{-x-c_2} + c_1$$




Figure 2.39: Slope field plot
 $y'' + y' = 0$

Solved as second order integrable as is ode (ABC method)

Time used: 0.044 (sec)

Writing the ode as

$$y'' + y' = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = 0$$

$$y' + y = c_1$$

Which is now solved for y . Integrating gives

$$\int \frac{1}{-y + c_1} dy = dx$$

$$-\ln(-y + c_1) = x + c_2$$

Singular solutions are found by solving

$$-y + c_1 = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = c_1$$

Solving for y gives

$$y = -e^{-x-c_2} + c_1$$

Will add steps showing solving for IC soon.




Figure 2.40: Slope field plot
 $y'' + y' = 0$

Solved as second order ode using Kovacic algorithm

Time used: 0.037 (sec)

Writing the ode as

$$y'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.14: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2$$




Figure 2.41: Slope field plot
 $y'' + y' = 0$

Solved as second order ode adjoint method

Time used: 0.411 (sec)

In normal form the ode

$$y'' + y' = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = \frac{c_2}{c_1 e^x + c_2}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2} \end{aligned}$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$

$$\frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) = 0$$

Integrating gives

$$\begin{aligned} \frac{y e^x}{c_1 e^x + c_2} &= \int 0 dx + c_3 \\ &= c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = c_3 (c_1 + c_2 e^{-x})$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_3 (c_1 + c_2 e^{-x})$$

The constants can be merged to give

$$y = c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 + c_2 e^{-x}$$




Figure 2.42: Slope field plot
 $y'' + y' = 0$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x)$$

- Substitute in solutions

$$y(x) = C1 e^{-x} + C2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`
```

Maple dsolve solution

Solving time : 0.004 (sec)

Leaf size : 12

```
dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = 0,
        y(x),singsol=all)
```

$$y = c_1 + e^{-x}c_2$$

Mathematica DSolve solution

Solving time : 0.01 (sec)

Leaf size : 17

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]==0,{x},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2 - c_1 e^{-x}$$

2.1.13 problem 13

Solved as second order missing y ode	97
Maple step by step solution	99
Maple trace	99
Maple dsolve solution	99
Mathematica DSolve solution	100

Internal problem ID [8760]

Book : Second order enumerated odes

Section : section 1

Problem number : 13

Date solved : Thursday, December 12, 2024 at 09:43:01 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y''^2 + y' = 0$$

Solved as second order missing y ode

Time used: 0.596 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^2 + p(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Solving for the derivative gives these ODE's to solve

$$p'(x) = \sqrt{-p(x)} \tag{1}$$

$$p'(x) = -\sqrt{-p(x)} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Integrating gives

$$\int \frac{1}{\sqrt{-p}} dp = dx$$

$$-2\sqrt{-p} = x + c_1$$

Singular solutions are found by solving

$$\sqrt{-p} = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 0$$

Solving Eq. (2)

Integrating gives

$$\int -\frac{1}{\sqrt{-p}} dp = dx$$

$$2\sqrt{-p} = x + c_2$$

Singular solutions are found by solving

$$-\sqrt{-p} = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$-2\sqrt{-y'} = x + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\frac{1}{4}x^2 - \frac{1}{2}c_1x - \frac{1}{4}c_1^2 dx$$

$$y = -\frac{(x + c_1)^3}{12} + c_3$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$2\sqrt{-y'} = x + c_2$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\frac{1}{4}x^2 - \frac{1}{2}c_2x - \frac{1}{4}c_2^2 dx$$

$$y = -\frac{(x + c_2)^3}{12} + c_4$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 0$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_5$$

$$y = c_5$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_5$$

$$y = -\frac{(x + c_1)^3}{12} + c_3$$

$$y = -\frac{(x + c_2)^3}{12} + c_4$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each res
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
-> Calling odsolve with the ODE`, diff(diff(diff(y(x), x), x), x)+1/2, y(x)`
Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful
<- 2nd order ODE linearizable_by_differentiation successful
-----
* Tackling next ODE.
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
<- 2nd order ODE linearizable_by_differentiation successful`

```

Maple dsolve solution

Solving time : 0.305 (sec)

Leaf size : 27

```

dsolve(diff(diff(y(x),x),x)^2+diff(y(x),x) = 0,
y(x),singsol=all)

```

$$y = c_1$$

$$y = -\frac{1}{12}x^3 + \frac{1}{2}c_1x^2 - c_1^2x + c_2$$

Mathematica DSolve solution

Solving time : 0.022 (sec)

Leaf size : 69

```
DSolve[{(D[y[x],{x,2}])^2+D[y[x],x]==0,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{x^3}{12} - \frac{1}{4}ic_1x^2 + \frac{c_1^2x}{4} + c_2$$

$$y(x) \rightarrow -\frac{x^3}{12} + \frac{1}{4}ic_1x^2 + \frac{c_1^2x}{4} + c_2$$

2.1.14 problem 14

Solved as second order missing y ode	101
Maple step by step solution	102
Maple trace	103
Maple dsolve solution	103
Mathematica DSolve solution	103

Internal problem ID [8761]

Book : Second order enumerated odes

Section : section 1

Problem number : 14

Date solved : Thursday, December 12, 2024 at 09:43:02 AM

CAS classification :

[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + y'^2 = 0$$

Solved as second order missing y ode

Time used: 0.093 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating gives

$$\int -\frac{1}{p^2} dp = dx$$

$$\frac{1}{p} = x + c_1$$

Singular solutions are found by solving

$$-p^2 = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 0$$

Solving for $p(x)$ gives

$$p(x) = \frac{1}{x + c_1}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 0$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_2$$

$$y = c_2$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{1}{x + c_1}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{1}{x + c_1} dx$$

$$y = \ln(x + c_1) + c_3$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2$$

$$y = \ln(x + c_1) + c_3$$

Maple step by step solution

Let's solve

$$\left(\frac{d}{dx}y(x)\right)^2 + \frac{d^2}{dx^2}y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Make substitution $u = \frac{d}{dx}y(x)$ to reduce order of ODE

$$u(x)^2 + \frac{d}{dx}u(x) = 0$$

- Solve for the highest derivative

$$\frac{d}{dx}u(x) = -u(x)^2$$

- Separate variables

$$\frac{\frac{d}{dx}u(x)}{u(x)^2} = -1$$

- Integrate both sides with respect to x

$$\int \frac{\frac{d}{dx}u(x)}{u(x)^2} dx = \int (-1) dx + C1$$

- Evaluate integral

$$-\frac{1}{u(x)} = C1 - x$$

- Solve for $u(x)$

$$u(x) = -\frac{1}{C1-x}$$

- Solve 1st ODE for $u(x)$

$$u(x) = -\frac{1}{C1-x}$$

- Make substitution $u = \frac{d}{dx}y(x)$

$$\frac{d}{dx}y(x) = -\frac{1}{C1-x}$$

- Integrate both sides to solve for $y(x)$

$$\int \left(\frac{d}{dx}y(x)\right) dx = \int -\frac{1}{C1-x} dx + C2$$

- Compute integrals

$$y(x) = \ln(C1 - x) + C2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.016 (sec)

Leaf size : 10

```
dsolve(diff(diff(y(x),x),x)+diff(y(x),x)^2 = 0,  
        y(x),singsol=all)
```

$$y = \ln(c_1x + c_2)$$

Mathematica DSolve solution

Solving time : 0.193 (sec)

Leaf size : 15

```
DSolve[{D[y[x],{x,2}]+(D[y[x],x])^2==0,{}},  
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \log(x - c_1) + c_2$$

2.1.15 problem 15

Solved as second order linear constant coeff ode	104
Solved as second order linear exact ode	106
Solved as second order missing y ode	108
Solved as second order integrable as is ode	109
Solved as second order integrable as is ode (ABC method)	110
Solved as second order ode using Kovacic algorithm	112
Solved as second order ode adjoint method	115
Maple step by step solution	118
Maple trace	118
Maple dsolve solution	119
Mathematica DSolve solution	119

Internal problem ID [8762]

Book : Second order enumerated odes

Section : section 1

Problem number : 15

Date solved : Thursday, December 12, 2024 at 09:43:02 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y'' + y' = 1$$

Solved as second order linear constant coeff ode

Time used: 0.109 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + (x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_1 + c_2 e^{-x}$$




Figure 2.43: Slope field plot
 $y'' + y' = 1$

Solved as second order linear exact ode

Time used: 0.102 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= 1 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int 1 dx$$

We now have a first order ode to solve which is

$$y' + y = x + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= x + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= ((x + c_1) e^x) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int (x + c_1) e^x dx \\ &= (x + c_1 - 1) e^x + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = c_1 + x - 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 + x - 1 + c_2 e^{-x}$$




Figure 2.44: Slope field plot
 $y'' + y' = 1$

Solved as second order missing y ode

Time used: 0.152 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating gives

$$\int \frac{1}{1-p} dp = dx$$

$$-\ln(-1+p) = x + c_1$$

Singular solutions are found by solving

$$1 - p = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 1$$

Solving for $p(x)$ gives

$$p(x) = e^{-x-c_1} + 1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 1 dx$$

$$y = x + c_2$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = e^{-x-c_1} + 1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int e^{-x-c_1} + 1 dx$$

$$y = x - e^{-x-c_1} + c_3$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_2$$

$$y = x - e^{-x-c_1} + c_3$$




Figure 2.45: Slope field plot
 $y'' + y' = 1$

Solved as second order integrable as is ode

Time used: 0.063 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int 1 dx$$

$$y' + y = x + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = x + c_1$$

The integrating factor μ is

$$\mu = e^{\int q dx}$$

$$= e^{\int 1 dx}$$

$$= e^x$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= ((x + c_1) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int (x + c_1) e^x dx \\ &= (x + c_1 - 1) e^x + c_2\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = c_1 + x - 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 + x - 1 + c_2 e^{-x}$$




Figure 2.46: Slope field plot
 $y'' + y' = 1$

Solved as second order integrable as is ode (ABC method)

Time used: 0.047 (sec)

Writing the ode as

$$y'' + y' = 1$$

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned}\int (y'' + y') dx &= \int 1 dx \\ y' + y &= x + c_1\end{aligned}$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= 1 \\p(x) &= x + c_1\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\&= e^{\int 1 dx} \\&= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= ((x + c_1) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int (x + c_1) e^x dx \\ &= (x + c_1 - 1) e^x + c_2\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = c_1 + x - 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.47: Slope field plot
 $y'' + y' = 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.061 (sec)

Writing the ode as

$$y'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.17: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 (e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 (e^{-x} (e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + (x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 + x$$




Figure 2.48: Slope field plot
 $y'' + y' = 1$

Solved as second order ode adjoint method

Time used: 0.457 (sec)

In normal form the ode

$$y'' + y' = 1 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$p(x) = 1$$

$$q(x) = 0$$

$$r(x) = 1$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 1 \\ \lambda_2 &= 0\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}\xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x}\end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = \frac{c_2 x + c_1 e^x}{c_1 e^x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= \frac{c_2}{c_1 e^x + c_2} \\ p(x) &= \frac{c_2 x + c_1 e^x}{c_1 e^x + c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{c_2 x + c_1 e^x}{c_1 e^x + c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{e^x}{c_1 e^x + c_2} \right) \left(\frac{c_2 x + c_1 e^x}{c_1 e^x + c_2} \right) \\ d \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{(c_2 x + c_1 e^x) e^x}{(c_1 e^x + c_2)^2} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y e^x}{c_1 e^x + c_2} &= \int \frac{(c_2 x + c_1 e^x) e^x}{(c_1 e^x + c_2)^2} dx \\ &= \frac{c_2}{c_1 (c_1 e^x + c_2)} + \frac{x e^x}{c_1 e^x + c_2} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = \frac{c_2(c_3 c_1 + 1) e^{-x} + c_1(c_3 c_1 + x)}{c_1}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{c_2(c_3 c_1 + 1) e^{-x} + c_1(c_3 c_1 + x)}{c_1}$$

The constants can be merged to give

$$y = \frac{c_2(c_1 + 1) e^{-x} + c_1(x + c_1)}{c_1}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_2(c_1 + 1) e^{-x} + c_1(x + c_1)}{c_1}$$




Figure 2.49: Slope field plot
 $y'' + y' = 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x dx \right) + \int 1 dx$$

- Compute integrals

$$y_p(x) = x - 1$$

- Substitute particular solution into general solution to ODE

$$y(x) = C1 e^{-x} + C2 + x - 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 14

```
dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = 1,  
        y(x),singsol=all)
```

$$y = -c_1 e^{-x} + x + c_2$$

Mathematica DSolve solution

Solving time : 0.012 (sec)

Leaf size : 18

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]==1,{}},  
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x - c_1 e^{-x} + c_2$$

2.1.16 problem 16

Solved as second order missing y ode	120
Maple step by step solution	122
Maple trace	122
Maple dsolve solution	122
Mathematica DSolve solution	123

Internal problem ID [8763]

Book : Second order enumerated odes

Section : section 1

Problem number : 16

Date solved : Thursday, December 12, 2024 at 09:43:04 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y''^2 + y' = 1$$

Solved as second order missing y ode

Time used: 0.595 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^2 + p(x) - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Solving for the derivative gives these ODE's to solve

$$p'(x) = \sqrt{1 - p(x)} \quad (1)$$

$$p'(x) = -\sqrt{1 - p(x)} \quad (2)$$

Now each of the above is solved separately.

Solving Eq. (1)

Integrating gives

$$\int \frac{1}{\sqrt{1-p}} dp = dx$$

$$-2\sqrt{1-p} = x + c_1$$

Singular solutions are found by solving

$$\sqrt{1-p} = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 1$$

Solving Eq. (2)

Integrating gives

$$\int -\frac{1}{\sqrt{1-p}} dp = dx$$

$$2\sqrt{1-p} = x + c_2$$

Singular solutions are found by solving

$$-\sqrt{1-p} = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$-2\sqrt{1-y'} = x + c_1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\frac{1}{4}c_1^2 - \frac{1}{2}c_1x - \frac{1}{4}x^2 + 1 dx$$

$$y = -\frac{x^3}{12} - \frac{c_1 x^2}{4} - \frac{(c_1 + 2)(c_1 - 2)x}{4} + c_3$$

$$y = -\frac{1}{12}x^3 - \frac{1}{4}c_1 x^2 - \frac{1}{4}c_1^2 x + x + c_3$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$2\sqrt{1-y'} = x + c_2$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\frac{1}{4}c_2^2 - \frac{1}{2}c_2x - \frac{1}{4}x^2 + 1 dx$$

$$y = -\frac{x^3}{12} - \frac{c_2 x^2}{4} - \frac{(c_2 + 2)(c_2 - 2)x}{4} + c_4$$

$$y = -\frac{1}{12}x^3 - \frac{1}{4}c_2 x^2 - \frac{1}{4}c_2^2 x + x + c_4$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 1 dx$$

$$y = x + c_5$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_5$$

$$y = -\frac{1}{12}x^3 - \frac{1}{4}c_1 x^2 - \frac{1}{4}c_1^2 x + x + c_3$$

$$y = -\frac{1}{12}x^3 - \frac{1}{4}c_2 x^2 - \frac{1}{4}c_2^2 x + x + c_4$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each result
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
<- 2nd order ODE linearizable_by_differentiation successful
-----
* Tackling next ODE.
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
<- 2nd order ODE linearizable_by_differentiation successful
-> Calling odsolve with the ODE`, diff(y(x), x) = 1, y(x), singsol = none`
Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

Maple dsolve solution

Solving time : 0.306 (sec)

Leaf size : 30

```

dsolve(diff(diff(y(x),x),x)^2+diff(y(x),x) = 1,
        y(x),singsol=all)

```

$$y = x + c_1$$

$$y = -\frac{1}{12}x^3 + \frac{1}{2}c_1x^2 - c_1^2x + x + c_2$$

Mathematica DSolve solution

Solving time : 0.024 (sec)

Leaf size : 67

```
DSolve[{(D[y[x],{x,2}])^2+D[y[x],x]==1,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{x^3}{12} - \frac{c_1 x^2}{4} + x - \frac{c_1^2 x}{4} + c_2$$

$$y(x) \rightarrow -\frac{x^3}{12} + \frac{c_1 x^2}{4} + x - \frac{c_1^2 x}{4} + c_2$$

2.1.17 problem 17

Solved as second order missing y ode	124
Maple step by step solution	125
Maple trace	126
Maple dsolve solution	126
Mathematica DSolve solution	126

Internal problem ID [8764]

Book : Second order enumerated odes

Section : section 1

Problem number : 17

Date solved : Thursday, December 12, 2024 at 09:43:05 AM

CAS classification :

[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + y'^2 = 1$$

Solved as second order missing y ode

Time used: 0.200 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x)^2 - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating gives

$$\int \frac{1}{-p^2 + 1} dp = dx$$

$$\operatorname{arctanh}(p) = x + c_1$$

Singular solutions are found by solving

$$-p^2 + 1 = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = -1$$

$$p(x) = 1$$

Solving for $p(x)$ gives

$$p(x) = \tanh(x + c_1)$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -1 dx$$

$$y = -x + c_2$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 1 dx$$

$$y = x + c_3$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \tanh(x + c_1)$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \tanh(x + c_1) dx$$

$$y = \ln(\cosh(x + c_1)) + c_4$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -x + c_2$$

$$y = x + c_3$$

$$y = \ln(\cosh(x + c_1)) + c_4$$

Maple step by step solution

Let's solve

$$\left(\frac{d}{dx}y(x)\right)^2 + \frac{d^2}{dx^2}y(x) = 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Make substitution $u = \frac{d}{dx}y(x)$ to reduce order of ODE

$$u(x)^2 + \frac{d}{dx}u(x) = 1$$

- Solve for the highest derivative

$$\frac{d}{dx}u(x) = -u(x)^2 + 1$$

- Separate variables

$$\frac{\frac{d}{dx}u(x)}{-u(x)^2+1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{\frac{d}{dx}u(x)}{-u(x)^2+1} dx = \int 1 dx + C1$$

- Evaluate integral

$$\arctanh(u(x)) = x + C1$$

- Solve for $u(x)$

- $$u(x) = \tanh(x + C1)$$
- Solve 1st ODE for $u(x)$
 $u(x) = \tanh(x + C1)$
 - Make substitution $u = \frac{d}{dx}y(x)$
 $\frac{d}{dx}y(x) = \tanh(x + C1)$
 - Integrate both sides to solve for $y(x)$
 $\int \left(\frac{d}{dx}y(x)\right) dx = \int \tanh(x + C1) dx + C2$
 - Compute integrals
 $y(x) = \ln(\cosh(x + C1)) + C2$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
<- 2nd order, 2 integrating factors of the form mu(x,y) successful`

```

Maple dsolve solution

Solving time : 0.017 (sec)

Leaf size : 22

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)^2 = 1,
        y(x),singsol=all)

```

$$y = -x - \ln(2) + \ln(-e^{2x}c_1 + c_2)$$

Mathematica DSolve solution

Solving time : 0.307 (sec)

Leaf size : 48

```

DSolve[{D[y[x],{x,2}]+(D[y[x],x])^2==1,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow -\frac{1}{2} \log(e^{2x}) + \log(e^{2x} + e^{2c_1}) + c_2$$

$$y(x) \rightarrow \frac{1}{2} \log(e^{2x}) + c_2$$

2.1.18 problem 18

Solved as second order linear constant coeff ode	127
Solved as second order linear exact ode	129
Solved as second order missing y ode	131
Solved as second order integrable as is ode	133
Solved as second order integrable as is ode (ABC method)	134
Solved as second order ode using Kovacic algorithm	135
Solved as second order ode adjoint method	138
Maple step by step solution	141
Maple trace	142
Maple dsolve solution	142
Mathematica DSolve solution	142

Internal problem ID [8765]

Book : Second order enumerated odes

Section : section 1

Problem number : 18

Date solved : Thursday, December 12, 2024 at 09:43:06 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y'' + y' = x$$

Solved as second order linear constant coeff ode

Time used: 0.123 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2xA_2 + A_1 + 2A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -1, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{2}x^2 - x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + \left(\frac{1}{2}x^2 - x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} - x + c_1 + c_2 e^{-x}$$




Figure 2.50: Slope field plot
 $y'' + y' = x$

Solved as second order linear exact ode

Time used: 0.105 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= x \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int x dx$$

We now have a first order ode to solve which is

$$y' + y = \frac{x^2}{2} + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= \frac{x^2}{2} + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{x^2}{2} + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{x^2}{2} + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{x^2}{2} + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{x^2}{2} + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1 - 2x + 2) e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$




Figure 2.51: Slope field plot
 $y'' + y' = x$

Solved as second order missing y ode

Time used: 0.092 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - x = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = x$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= \mu p \\ \frac{d}{dx}(\mu p) &= (\mu)(x) \\ \frac{d}{dx}(p e^x) &= (e^x)(x) \\ d(p e^x) &= (x e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}p e^x &= \int x e^x dx \\ &= (x - 1) e^x + c_1\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$p(x) = x - 1 + c_1 e^{-x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x - 1 + c_1 e^{-x}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned}\int dy &= \int x - 1 + c_1 e^{-x} dx \\ y &= -x + \frac{x^2}{2} - c_1 e^{-x} + c_2\end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -x + \frac{x^2}{2} - c_1 e^{-x} + c_2$$




Figure 2.52: Slope field plot
 $y'' + y' = x$

Solved as second order integrable as is ode

Time used: 0.072 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int x dx$$

$$y' + y = \frac{x^2}{2} + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{x^2}{2} + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{x^2}{2} + c_1 \right)$$

$$\frac{d}{dx}(y e^x) = (e^x) \left(\frac{x^2}{2} + c_1 \right)$$

$$d(y e^x) = \left(\left(\frac{x^2}{2} + c_1 \right) e^x \right) dx$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{x^2}{2} + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1 - 2x + 2) e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$




Figure 2.53: Slope field plot
 $y'' + y' = x$

Solved as second order integrable as is ode (ABC method)

Time used: 0.049 (sec)

Writing the ode as

$$y'' + y' = x$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int x dx$$

$$y' + y = \frac{x^2}{2} + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{x^2}{2} + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{x^2}{2} + c_1 \right)$$

$$\frac{d}{dx}(y e^x) = (e^x) \left(\frac{x^2}{2} + c_1 \right)$$

$$d(y e^x) = \left(\left(\frac{x^2}{2} + c_1 \right) e^x \right) dx$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{x^2}{2} + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1 - 2x + 2) e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.54: Slope field plot
 $y'' + y' = x$

Solved as second order ode using Kovacic algorithm

Time used: 0.068 (sec)

Writing the ode as

$$y'' + y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.20: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2xA_2 + A_1 + 2A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -1, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{2}x^2 - x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + \left(\frac{1}{2}x^2 - x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 + \frac{x^2}{2} - x$$




Figure 2.55: Slope field plot
 $y'' + y' = x$

Solved as second order ode adjoint method

Time used: 0.503 (sec)

In normal form the ode

$$y'' + y' = x \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= x \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = \frac{\frac{c_2 x^2}{2} + c_1 (x e^x - e^x)}{c_1 e^x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= \frac{c_2}{c_1 e^x + c_2} \\ p(x) &= \frac{2c_1(x-1)e^x + c_2 x^2}{2c_1 e^x + 2c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{2c_1(x-1)e^x + c_2 x^2}{2c_1 e^x + 2c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{e^x}{c_1 e^x + c_2} \right) \left(\frac{2c_1(x-1)e^x + c_2 x^2}{2c_1 e^x + 2c_2} \right) \\ d \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{(2c_1(x-1)e^x + c_2 x^2) e^x}{(2c_1 e^x + 2c_2)(c_1 e^x + c_2)} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y e^x}{c_1 e^x + c_2} &= \int \frac{(2c_1(x-1)e^x + c_2 x^2) e^x}{(2c_1 e^x + 2c_2)(c_1 e^x + c_2)} dx \\ &= \frac{e^x - x e^x + \frac{e^x x^2}{2}}{c_1 e^x + c_2} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = c_1 c_3 + \frac{x^2}{2} - x + c_2 c_3 e^{-x} + 1$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_1 c_3 + \frac{x^2}{2} - x + c_2 c_3 e^{-x} + 1$$

The constants can be merged to give

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$




Figure 2.56: Slope field plot
 $y'' + y' = x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = x$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 e^{-x} + C_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x x dx \right) + \int x dx$$
- Compute integrals
$$y_p(x) = 1 - x + \frac{1}{2}x^2$$
- Substitute particular solution into general solution to ODE
$$y(x) = C_1 e^{-x} + C_2 + 1 - x + \frac{x^2}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)+_a, _b(_a)` *** Sublevel
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)
Leaf size : 21

```

dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = x,
        y(x),singsol=all)

```

$$y = \frac{x^2}{2} - c_1 e^{-x} - x + c_2$$

Mathematica DSolve solution

Solving time : 0.039 (sec)
Leaf size : 27

```

DSolve[{D[y[x] ,{x,2}]+D[y[x] ,x]==x,{}},
        y[x] ,x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{x^2}{2} - x - c_1 e^{-x} + c_2$$

2.1.19 problem 19

Solved as second order missing y ode	143
Maple step by step solution	145
Maple trace	145
Maple dsolve solution	145
Mathematica DSolve solution	145

Internal problem ID [8766]

Book : Second order enumerated odes

Section : section 1

Problem number : 19

Date solved : Thursday, December 12, 2024 at 09:43:08 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y''^2 + y' = x$$

Solved as second order missing y ode

Time used: 0.561 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)^2 + p(x) - x = 0$$

Which is now solve for $p(x)$ as first order ode. Let $p = p'(x)$ the ode becomes

$$p^2 + p - x = 0$$

Solving for $p(x)$ from the above results in

$$p(x) = -p^2 + x \tag{1}$$

This has the form

$$p = xf(p) + g(p) \tag{*}$$

Where f, g are functions of $p = p'(x)$. The above ode is dAlembert ode which is now solved.

Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \tag{2}$$

Comparing the form $p(x) = xf + g$ to (1A) shows that

$$\begin{aligned} f &= 1 \\ g &= -p^2 \end{aligned}$$

Hence (2) becomes

$$p - 1 = -2pp'(x) \quad (2A)$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = -\frac{p(x) - 1}{2p(x)} \quad (3)$$

This ODE is now solved for $p(x)$. No inversion is needed. Integrating gives

$$\int -\frac{2p}{p-1} dp = dx$$

$$-2p - 2 \ln(p-1) = x + c_1$$

Singular solutions are found by solving

$$-\frac{p-1}{2p} = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 1$$

Solving for $p(x)$ gives

$$p(x) = \text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right) + 1$$

Substituting the above solution for p in (2A) gives

$$p(x) = x - 1$$

$$p(x) = -\left(\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right) + 1\right)^2 + x$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x - 1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int x - 1 dx$$

$$y = \frac{1}{2}x^2 - x + c_2$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\left(\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right) + 1\right)^2 + x$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int -\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right)^2 - 2\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right) + x - 1 dx$$

$$y = \frac{2\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right)^3}{3} + 3\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right)^2 + 4\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right) + \frac{x^2}{2} - x + c_3$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{2}x^2 - x + c_2$$

$$y = \frac{2\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right)^3}{3} + 3\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right)^2$$

$$+ 4\text{LambertW}\left(e^{-1-\frac{x}{2}-\frac{c_1}{2}}\right) + \frac{x^2}{2} - x + c_3$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each result
*** Sublevel 2 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = (-_b(_a)+_a)^(1/2), _b(_a), HI
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries:`[1, 1]

```

Maple dsolve solution

Solving time : 0.113 (sec)

Leaf size : 122

```

dsolve(diff(diff(y(x),x),x)^2+diff(y(x),x) = x,
        y(x),singsol=all)

```

$$\begin{aligned}
 y &= \int \left(-e^{2\text{RootOf}(-Z-x-2e^{-Z}+2+c_1-\ln(e^{-Z}(e^{-Z}-2)^2))} \right. \\
 &\quad \left. + 2e^{\text{RootOf}(-Z-x-2e^{-Z}+2+c_1-\ln(e^{-Z}(e^{-Z}-2)^2))} + x \right) dx - x + c_2 \\
 y &= \frac{2\text{LambertW}(-c_1e^{-\frac{x}{2}-1})^3}{3} + 3\text{LambertW}(-c_1e^{-\frac{x}{2}-1})^2 \\
 &\quad + 4\text{LambertW}(-c_1e^{-\frac{x}{2}-1}) + \frac{x^2}{2} - x + c_2
 \end{aligned}$$

Mathematica DSolve solution

Solving time : 16.306 (sec)

Leaf size : 172

```

DSolve[{(D[y[x],{x,2}])^2+D[y[x],x]==x,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$\begin{aligned}
 y(x) &\rightarrow \frac{2}{3}W\left(e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right)^3 + 3W\left(e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right)^2 + 4W\left(e^{-\frac{x}{2}-1-\frac{c_1}{2}}\right) + \frac{x^2}{2} - x + c_2 \\
 y(x) &\rightarrow \frac{2}{3}W\left(-e^{\frac{1}{2}(-x-2+c_1)}\right)^3 + 3W\left(-e^{\frac{1}{2}(-x-2+c_1)}\right)^2 + 4W\left(-e^{\frac{1}{2}(-x-2+c_1)}\right) + \frac{x^2}{2} - x + c_2 \\
 y(x) &\rightarrow \frac{x^2}{2} - x + c_2
 \end{aligned}$$

2.1.20 problem 20

Solved as second order missing y ode 146
 Maple step by step solution 147
 Maple trace 147
 Maple dsolve solution 148
 Mathematica DSolve solution 148

Internal problem ID [8767]

Book : Second order enumerated odes

Section : section 1

Problem number : 20

Date solved : Thursday, December 12, 2024 at 09:43:09 AM

CAS classification :

[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + y'^2 = x$$

Solved as second order missing y ode

Time used: 0.741 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x)^2 - x = 0$$

Which is now solve for $p(x)$ as first order ode. This is reduced Riccati ode of the form

$$p'(x) = ax^n + bp(x)^2$$

Comparing the given ode to the above shows that

$$\begin{aligned} a &= 1 \\ b &= -1 \\ n &= 1 \end{aligned}$$

Since $n \neq -2$ then the solution of the reduced Riccati ode is given by

$$w = \sqrt{x} \begin{cases} c_1 \text{BesselJ}\left(\frac{1}{2k}, \frac{1}{k}\sqrt{abx^k}\right) + c_2 \text{BesselY}\left(\frac{1}{2k}, \frac{1}{k}\sqrt{abx^k}\right) & ab > 0 \\ c_1 \text{BesselI}\left(\frac{1}{2k}, \frac{1}{k}\sqrt{-abx^k}\right) + c_2 \text{BesselK}\left(\frac{1}{2k}, \frac{1}{k}\sqrt{-abx^k}\right) & ab < 0 \end{cases} \quad (1)$$

$$p(x) = -\frac{1}{b} \frac{w'}{w}$$

$$k = 1 + \frac{n}{2}$$

Since $ab < 0$ then EQ(1) gives

$$k = \frac{3}{2}$$

$$w = \sqrt{x} \left(c_1 \text{BesselI}\left(\frac{1}{3}, \frac{2x^{3/2}}{3}\right) + c_2 \text{BesselK}\left(\frac{1}{3}, \frac{2x^{3/2}}{3}\right) \right)$$

Therefore the solution becomes

$$p(x) = -\frac{1}{b} \frac{w'}{w}$$

Substituting the value of b, w found above and simplifying gives

$$p(x) = \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_2 \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + c_2 \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}$$

Letting $c_2 = 1$ the above becomes

$$p(x) = \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned} \int dy &= \int \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)} dx \\ y &= \int \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)} dx + c_3 \\ y &= \int \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)} dx + c_3 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = - \int \frac{\sqrt{x} \left(-\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 + \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)} dx + c_3$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying a quadrature

```

```

checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  <- Bessel successful
<- special function solution successful
<- 2nd order, 2 integrating factors of the form mu(x,y) successful`

```

Maple dsolve solution

Solving time : 0.016 (sec)

Leaf size : 18

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)^2 = x,
        y(x),singsol=all)

```

$$y = \ln(\pi) + \ln(c_1 \text{AiryAi}(x) - c_2 \text{AiryBi}(x))$$

Mathematica DSolve solution

Solving time : 0.106 (sec)

Leaf size : 15

```

DSolve[{D[y[x],{x,2}]+(D[y[x],x])^2==0,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \log(x - c_1) + c_2$$

2.1.21 problem 21

Solved as second order linear constant coeff ode	149
Solved as second order ode using Kovacic algorithm	150
Solved as second order ode adjoint method	153
Maple step by step solution	156
Maple trace	157
Maple dsolve solution	157
Mathematica DSolve solution	157

Internal problem ID [8768]

Book : Second order enumerated odes

Section : section 1

Problem number : 21

Date solved : Thursday, December 12, 2024 at 09:43:10 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y'' + y' + y = 0$$

Solved as second order linear constant coeff ode

Time used: 0.107 (sec)

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$




Figure 2.57: Slope field plot
 $y'' + y' + y = 0$

Solved as second order ode using Kovacic algorithm

Time used: 0.162 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1 \tag{3}$$

$$B = 1$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.22: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin \left(\frac{\sqrt{3} x}{2} \right)}{3}$$




Figure 2.58: Slope field plot
 $y'' + y' + y = 0$

Solved as second order ode adjoint method

Time used: 1.568 (sec)

In normal form the ode

$$y'' + y' + y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$p(x) = 1$$

$$q(x) = 1$$

$$r(x) = 0$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (\xi(x)) &= 0 \\ \xi''(x) - \xi'(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(\frac{e^{\frac{x}{2}} (c_1 \cos(\frac{\sqrt{3}x}{2}) + c_2 \sin(\frac{\sqrt{3}x}{2}))}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin(\frac{\sqrt{3}x}{2})}{2} + \frac{c_2 \sqrt{3} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \\ p(x) &= 0\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2\sqrt{3}-c_1)\cos\left(\frac{\sqrt{3}x}{2}\right)-\sin\left(\frac{\sqrt{3}x}{2}\right)(c_1\sqrt{3}+c_2)}{2c_1\cos\left(\frac{\sqrt{3}x}{2}\right)+2c_2\sin\left(\frac{\sqrt{3}x}{2}\right)} dx} \\ &= \frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1}\end{aligned}$$

The ode becomes

$$\frac{d}{dx}\left(\frac{y\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1}\right) = 0$$

Integrating gives

$$\begin{aligned}\frac{y\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1} &= \int 0 dx + c_3 \\ &= c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1}$ gives the final solution

$$y = \frac{\left(\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1\right) e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} c_3}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{\left(\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1\right) e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} c_3}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

The constants can be merged to give

$$y = \frac{\left(\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1\right) e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{\left(\tan\left(\frac{\sqrt{3}x}{2}\right)c_2+c_1\right) e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$




Figure 2.59: Slope field plot
 $y'' + y' + y = 0$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x)$$

- Substitute in solutions

$$y(x) = C1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

Maple dsolve solution

Solving time : 0.002 (sec)

Leaf size : 28

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = 0,
        y(x),singsol=all)

```

$$y = e^{-\frac{x}{2}} \left(c_1 \sin \left(\frac{\sqrt{3}x}{2} \right) + c_2 \cos \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Mathematica DSolve solution

Solving time : 0.023 (sec)

Leaf size : 42

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==0,{]},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow e^{-x/2} \left(c_2 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_1 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

2.1.22 problem 22

Maple step by step solution	158
Maple trace	158
Maple dsolve solution	159
Mathematica DSolve solution	159

Internal problem ID [8769]

Book : Second order enumerated odes

Section : section 1

Problem number : 22

Date solved : Thursday, December 12, 2024 at 09:43:12 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y''^2 + y' + y = 0$$

Does not support ODE with y''^n where $n \neq 1$ unless 1 is missing which is not the case here.

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each result
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-(-_b(_a)-_a)^(1/2) = 0
Methods for first order ODEs:
--- Trying classification methods ---
trying homogeneous types:
trying exact
Looking for potential symmetries
trying an equivalence to an Abel ODE
trying 1st order ODE linearizable_by_differentiation
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one int
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular
trying differential order: 2; exact nonlinear
trying 2nd order, integrating factor of the form mu(x,y)
-> trying 2nd order, the S-function method
-> trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods
-> trying 2nd order, the S-function method
-> trying 2nd order, No Point Symmetries Class V

```

```

-> trying 2nd order, No Point Symmetries Class V
-> trying 2nd order, No Point Symmetries Class V
trying 2nd order, integrating factor of the form  $\mu(x,y)/(y)^n$ , only the general
-> trying 2nd order, dynamical_symmetries, only a reduction of order through one
solving 2nd order ODE of high degree, Lie methods
`, `2nd order, trying reduction of order with given symmetries: `[1, 0]

```

Maple dsolve solution

Solving time : 0.097 (sec)
 Leaf size : maple_leaf_size

```

dsolve(diff(diff(y(x),x),x)^2+diff(y(x),x)+y(x) = 0,
        y(x),singsol=all)

```

No solution found

Mathematica DSolve solution

Solving time : 0.0 (sec)
 Leaf size : 0

```

DSolve[{(D[y[x],{x,2}])^2+D[y[x],x]+y[x]==0,{}},
        y[x],x,IncludeSingularSolutions->True]

```

Not solved

2.1.23 problem 23

Solved as second order missing x ode	160
Maple step by step solution	163
Maple trace	163
Maple dsolve solution	163
Mathematica DSolve solution	164

Internal problem ID [8770]

Book : Second order enumerated odes

Section : section 1

Problem number : 23

Date solved : Thursday, December 12, 2024 at 09:43:13 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y'' + y'^2 + y = 0$$

Solved as second order missing x ode

Time used: 0.788 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) + p(y)^2 + y = 0$$

Which is now solved as first order ode for $p(y)$.

In canonical form, the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= -\frac{p^2 + y}{p} \end{aligned}$$

This is a Bernoulli ODE.

$$p' = (-1)p + (-y) \frac{1}{p} \tag{1}$$

The standard Bernoulli ODE has the form

$$p' = f_0(y)p + f_1(y)p^n \tag{2}$$

Comparing this to (1) shows that

$$\begin{aligned} f_0 &= -1 \\ f_1 &= -y \end{aligned}$$

The first step is to divide the above equation by p^n which gives

$$\frac{p'}{p^n} = f_0(y)p^{1-n} + f_1(y) \quad (3)$$

The next step is use the substitution $v = p^{1-n}$ in equation (3) which generates a new ODE in $v(y)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $p(y)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(y) &= -1 \\ f_1(y) &= -y \\ n &= -1 \end{aligned}$$

Dividing both sides of ODE (1) by $p^n = \frac{1}{p}$ gives

$$p'p = -p^2 - y \quad (4)$$

Let

$$\begin{aligned} v &= p^{1-n} \\ &= p^2 \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t y gives

$$v' = 2pp' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} \frac{v'(y)}{2} &= -v(y) - y \\ v' &= -2v - 2y \end{aligned} \quad (7)$$

The above now is a linear ODE in $v(y)$ which is now solved.

In canonical form a linear first order is

$$v'(y) + q(y)v(y) = p(y)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(y) &= 2 \\ p(y) &= -2y \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q \, dy} \\ &= e^{\int 2 \, dy} \\ &= e^{2y} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dy}(\mu v) &= \mu p \\ \frac{d}{dy}(\mu v) &= (\mu)(-2y) \\ \frac{d}{dy}(v e^{2y}) &= (e^{2y})(-2y) \\ d(v e^{2y}) &= (-2y e^{2y}) \, dy \end{aligned}$$

Integrating gives

$$\begin{aligned} v e^{2y} &= \int -2y e^{2y} dy \\ &= -\frac{(2y-1)e^{2y}}{2} + c_1 \end{aligned}$$

Dividing throughout by the integrating factor e^{2y} gives the final solution

$$v(y) = -y + \frac{1}{2} + c_1 e^{-2y}$$

The substitution $v = p^{1-n}$ is now used to convert the above solution back to p which results in

$$p^2 = -y + \frac{1}{2} + c_1 e^{-2y}$$

Solving for p gives

$$\begin{aligned} p &= -\frac{\sqrt{2 + 4c_1 e^{-2y} - 4y}}{2} \\ p &= \frac{\sqrt{2 + 4c_1 e^{-2y} - 4y}}{2} \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{\sqrt{2 + 4c_1 e^{-2y} - 4y}}{2}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y -\frac{2}{\sqrt{2 + 4c_1 e^{-2\tau} - 4\tau}} d\tau = x + c_2$$

Singular solutions are found by solving

$$-\frac{\sqrt{2 + 4c_1 e^{-2y} - 4y}}{2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{\text{LambertW}(2c_1 e^{-1})}{2} + \frac{1}{2}$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{\sqrt{2 + 4c_1 e^{-2y} - 4y}}{2}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{2}{\sqrt{2 + 4c_1 e^{-2\tau} - 4\tau}} d\tau = x + c_3$$

Singular solutions are found by solving

$$\frac{\sqrt{2 + 4c_1 e^{-2y} - 4y}}{2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{\text{LambertW}(2c_1 e^{-1})}{2} + \frac{1}{2}$$

Will add steps showing solving for IC soon.

The solution

$$y = \frac{\text{LambertW}(2c_1 e^{-1})}{2} + \frac{1}{2}$$

was found not to satisfy the ode or the IC. Hence it is removed.

Summary of solutions found

$$\int^y -\frac{2}{\sqrt{2 + 4c_1 e^{-2\tau} - 4\tau}} d\tau = x + c_2$$

$$\int^y \frac{2}{\sqrt{2 + 4c_1 e^{-2\tau} - 4\tau}} d\tau = x + c_3$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(`_b(_a), _a))*_b(_a)+_b(_a)^2+_a = 0, `_b(_a)`
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`

```

Maple dsolve solution

Solving time : 0.028 (sec)

Leaf size : 61

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)^2+y(x) = 0,
        y(x),singsol=all)

```

$$-2 \left(\int^y \frac{1}{\sqrt{2 + 4e^{-2a}c_1 - 4a}} da \right) - x - c_2 = 0$$

$$2 \left(\int^y \frac{1}{\sqrt{2 + 4e^{-2a}c_1 - 4a}} da \right) - x - c_2 = 0$$

Mathematica DSolve solution

Solving time : 0.746 (sec)

Leaf size : 272

```
DSolve[{D[y[x], {x, 2}] + (D[y[x], x])^2 + y[x] == 0, {}},
  y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} -\frac{\sqrt{2}}{\sqrt{2e^{-2K[1]}c_1 - 2K[1] + 1}} dK[1] \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{\sqrt{2}}{\sqrt{2e^{-2K[2]}c_1 - 2K[2] + 1}} dK[2] \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} -\frac{\sqrt{2}}{\sqrt{2e^{-2K[1]}(-c_1) - 2K[1] + 1}} dK[1] \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} -\frac{\sqrt{2}}{\sqrt{2e^{-2K[1]}c_1 - 2K[1] + 1}} dK[1] \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{\sqrt{2}}{\sqrt{2e^{-2K[2]}(-c_1) - 2K[2] + 1}} dK[2] \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{\sqrt{2}}{\sqrt{2e^{-2K[2]}c_1 - 2K[2] + 1}} dK[2] \& \right] [x + c_2]$$

2.1.24 problem 24

Solved as second order linear constant coeff ode	165
Solved as second order ode using Kovacic algorithm	167
Solved as second order ode adjoint method	171
Maple step by step solution	174
Maple trace	175
Maple dsolve solution	175
Mathematica DSolve solution	176

Internal problem ID [8771]

Book : Second order enumerated odes

Section : section 1

Problem number : 24

Date solved : Thursday, December 12, 2024 at 09:43:14 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y'' + y' + y = 1$$

Solved as second order linear constant coeff ode

Time used: 0.155 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$




Figure 2.60: Slope field plot
 $y'' + y' + y = 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.204 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.24: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} + 1$$




Figure 2.61: Slope field plot

$$y'' + y' + y = 1$$

Solved as second order ode adjoint method

Time used: 19.541 (sec)

In normal form the ode

$$y'' + y' + y = 1 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$p(x) = 1$$

$$q(x) = 1$$

$$r(x) = 1$$

The Lagrange adjoint ode is given by

$$\xi'' - (\xi p)' + \xi q = 0$$

$$\xi'' - (\xi(x))' + (\xi(x)) = 0$$

$$\xi''(x) - \xi'(x) + \xi(x) = 0$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = \frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(\frac{e^{\frac{x}{2}} (c_1 \cos(\frac{\sqrt{3}x}{2}) + c_2 \sin(\frac{\sqrt{3}x}{2}))}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin(\frac{\sqrt{3}x}{2})}{2} + \frac{c_2 \sqrt{3} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \right) = \frac{e^{-\frac{x}{2}} \left(c_1 \left(\frac{e^{\frac{x}{2}} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right)}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \\ p(x) &= \frac{(-c_2 \sqrt{3} + c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) + \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} dx} \\ &= \frac{\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2} e^{\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}}}{c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{(-c_2\sqrt{3} + c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} + c_2)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$\begin{aligned} & \frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \\ &= \left(\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \left(\frac{(-c_2\sqrt{3} + c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} + c_2)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right) \\ & \frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \\ &= \left(\frac{\left((-c_2\sqrt{3} + c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} + c_2) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} &= \int \frac{\left((-c_2\sqrt{3} + c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} + c_2) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \\ &= \int \frac{\left((-c_2\sqrt{3} + c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} + c_2) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \end{aligned}$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}$ gives the final solution

$$y = \frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2\sqrt{3} + c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} + c_2) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2\sqrt{3} + c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} + c_2) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

The constants can be merged to give

y

$$\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2\sqrt{3}+c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3}+c_2) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)$$

$$\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

y

$$\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2\sqrt{3}+c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3}+c_2) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)$$

$$\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}$$




Figure 2.62: Slope field plot
 $y'' + y' + y = 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{-1 \pm \sqrt{-3}}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = 1$$

- Substitute particular solution into general solution to ODE

$$y(x) = C1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.005 (sec)

Leaf size : 32

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = 1,
        y(x),singsol=all)

```

$$y = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + 1$$

Mathematica DSolve solution

Solving time : 0.021 (sec)

Leaf size : 49

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==1,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{-x/2} \left(e^{x/2} + c_2 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_1 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

2.1.25 problem 25

Solved as second order linear constant coeff ode	177
Solved as second order ode using Kovacic algorithm	179
Solved as second order ode adjoint method	183
Maple step by step solution	186
Maple trace	187
Maple dsolve solution	187
Mathematica DSolve solution	188

Internal problem ID [8772]

Book : Second order enumerated odes

Section : section 1

Problem number : 25

Date solved : Thursday, December 12, 2024 at 09:43:34 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' + y' + y = x$$

Solved as second order linear constant coeff ode

Time used: 0.162 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2 x + A_1 + A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x - 1$$

Therefore the general solution is

$$y = y_h + y_p = \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (x - 1)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x - 1 + e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$




Figure 2.63: Slope field plot
 $y'' + y' + y = x$

Solved as second order ode using Kovacic algorithm

Time used: 0.210 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.26: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin \left(\frac{\sqrt{3} x}{2} \right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2x + A_1 + A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x - 1$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (x - 1)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} + x - 1$$




Figure 2.64: Slope field plot
 $y'' + y' + y = x$

Solved as second order ode adjoint method

Time used: 16.848 (sec)

In normal form the ode

$$y'' + y' + y = x \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$p(x) = 1$$

$$q(x) = 1$$

$$r(x) = x$$

The Lagrange adjoint ode is given by

$$\xi'' - (\xi p)' + \xi q = 0$$

$$\xi'' - (\xi(x))' + (\xi(x)) = 0$$

$$\xi''(x) - \xi'(x) + \xi(x) = 0$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = \frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(e^{\frac{x}{2}} \left(\frac{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} \right) + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin \left(\frac{\sqrt{3}x}{2} \right)}{2} + \frac{c_2 \sqrt{3} \cos \left(\frac{\sqrt{3}x}{2} \right)}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \right) = \frac{e^{-\frac{x}{2}} \left(c_1 \left(\frac{x}{2} + \frac{1}{2} \right) e^{\frac{x}{2}} \right)}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \\ p(x) &= \frac{(-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos \left(\frac{\sqrt{3}x}{2} \right) + \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1(x-1)\sqrt{3} + c_2(x+1))}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} dx} \\ &= \frac{\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2} e^{\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}}}{c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y)$$

$$= (\mu) \left(\frac{((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)))}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$\frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \left(\frac{((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)))}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$d \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1))) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} \right)$$

Integrating gives

$$\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} = \int \frac{((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

$$= \int \frac{((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}$ gives the final solution

$$y = \frac{(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

y

$$= \frac{(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

The constants can be merged to give

y

$$\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)$$

$$\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

y

$$\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2(x-1)\sqrt{3} + c_1(x+1)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1(x-1)\sqrt{3} + c_2(x+1)) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)$$

$$\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}$$




Figure 2.65: Slope field plot
 $y'' + y' + y = x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = x$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int x e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int x e^{\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = x - 1$$

- Substitute particular solution into general solution to ODE

$$y(x) = C1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + x - 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)

Leaf size : 33

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = x,
        y(x),singsol=all)

```

$$y = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + x - 1$$

Mathematica DSolve solution

Solving time : 0.022 (sec)

Leaf size : 50

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==x,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x + c_2 e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right) - 1$$

2.1.26 problem 26

Solved as second order linear constant coeff ode	189
Solved as second order ode using Kovacic algorithm	191
Solved as second order ode adjoint method	195
Maple step by step solution	198
Maple trace	199
Maple dsolve solution	199
Mathematica DSolve solution	200

Internal problem ID [8773]

Book : Second order enumerated odes

Section : section 1

Problem number : 26

Date solved : Thursday, December 12, 2024 at 09:43:52 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' + y' + y = 1 + x$$

Solved as second order linear constant coeff ode

Time used: 0.161 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = 1 + x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2x + A_1 + A_2 = 1 + x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$




Figure 2.66: Slope field plot
 $y'' + y' + y = 1 + x$

Solved as second order ode using Kovacic algorithm

Time used: 0.212 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.28: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin \left(\frac{\sqrt{3} x}{2} \right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2x + A_1 + A_2 = 1 + x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} + x$$




Figure 2.67: Slope field plot
 $y'' + y' + y = 1 + x$

Solved as second order ode adjoint method

Time used: 22.069 (sec)

In normal form the ode

$$y'' + y' + y = 1 + x \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 1 + x \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (\xi(x)) &= 0 \\ \xi''(x) - \xi'(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(\frac{e^{\frac{x}{2}} (c_1 \cos(\frac{\sqrt{3}x}{2}) + c_2 \sin(\frac{\sqrt{3}x}{2}))}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin(\frac{\sqrt{3}x}{2})}{2} + \frac{c_2 \sqrt{3} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \right) = \frac{e^{-\frac{x}{2}} \left(c_1 \left(\frac{e^{\frac{x}{2}} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right)}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \\ p(x) &= \frac{(-xc_2 \sqrt{3} + c_1(x+2)) \cos \left(\frac{\sqrt{3}x}{2} \right) + (xc_1 \sqrt{3} + c_2(x+2)) \sin \left(\frac{\sqrt{3}x}{2} \right)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} dx} \\ &= \frac{\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2} e^{\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}}}{c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{(-xc_2\sqrt{3} + c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3} + c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$\frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \left(\frac{(-xc_2\sqrt{3} + c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3} + c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$d \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\left((-xc_2\sqrt{3} + c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3} + c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} \right)$$

Integrating gives

$$\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} = \int \frac{\left((-xc_2\sqrt{3} + c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3} + c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx$$

$$= \int \frac{\left((-xc_2\sqrt{3} + c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3} + c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}$ gives the final solution

$$y = \frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-xc_2\sqrt{3} + c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3} + c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-xc_2\sqrt{3} + c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3} + c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

The constants can be merged to give

y

$$\frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-xc_2\sqrt{3}+c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3}+c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right)\right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)\right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)\right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

y

$$\frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)\right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-xc_2\sqrt{3}+c_1(x+2)) \cos\left(\frac{\sqrt{3}x}{2}\right) + (xc_1\sqrt{3}+c_2(x+2)) \sin\left(\frac{\sqrt{3}x}{2}\right)\right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)\right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)\right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$




Figure 2.68: Slope field plot
 $y'' + y' + y = 1 + x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int (x+1)e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int (x+1)e^{\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = x$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 32

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = x+1,
        y(x),singsol=all)

```

$$y = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + x$$

Mathematica DSolve solution

Solving time : 0.021 (sec)

Leaf size : 49

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==1+x,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x + c_2 e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

2.1.27 problem 27

Solved as second order linear constant coeff ode	201
Solved as second order ode using Kovacic algorithm	203
Solved as second order ode adjoint method	207
Maple step by step solution	211
Maple trace	212
Maple dsolve solution	212
Mathematica DSolve solution	212

Internal problem ID [8774]

Book : Second order enumerated odes

Section : section 1

Problem number : 27

Date solved : Thursday, December 12, 2024 at 09:44:15 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' + y' + y = x^2 + x + 1$$

Solved as second order linear constant coeff ode

Time used: 0.166 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = x^2 + x + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_3 x^2 + A_2 x + 2x A_3 + A_1 + A_2 + 2A_3 = x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = -1, A_3 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^2 - x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (x^2 - x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^2 - x + e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$




Figure 2.69: Slope field plot
 $y'' + y' + y = x^2 + x + 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.280 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.30: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}}\right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3}\right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3}\right)\right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3x^2 + A_2x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_3x^2 + A_2x + 2xA_3 + A_1 + A_2 + 2A_3 = x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = -1, A_3 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^2 - x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (x^2 - x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} + x^2 - x$$




Figure 2.70: Slope field plot
 $y'' + y' + y = x^2 + x + 1$

Solved as second order ode adjoint method

Time used: 32.022 (sec)

In normal form the ode

$$y'' + y' + y = x^2 + x + 1 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= x^2 + x + 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (\xi(x)) &= 0 \\ \xi''(x) - \xi'(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(\frac{e^{\frac{x}{2}} (c_1 \cos(\frac{\sqrt{3}x}{2}) + c_2 \sin(\frac{\sqrt{3}x}{2}))}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin(\frac{\sqrt{3}x}{2})}{2} + \frac{c_2 \sqrt{3} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \right) = \frac{e^{-\frac{x}{2}} \left(c_1 \left(\frac{1}{2} x^2 + x \right) \right)}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \\ p(x) &= \frac{(-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2)) \cos \left(\frac{\sqrt{3}x}{2} \right) + \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2))}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2\sqrt{3}-c_1)\cos\left(\frac{\sqrt{3}x}{2}\right)-\sin\left(\frac{\sqrt{3}x}{2}\right)(c_1\sqrt{3}+c_2)}{2c_1\cos\left(\frac{\sqrt{3}x}{2}\right)+2c_2\sin\left(\frac{\sqrt{3}x}{2}\right)} dx} \\ &= \frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}\end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y)$$

$$= (\mu) \left(\frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2))\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)(c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2)))}{2c_1\cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2\sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$\frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \left(\frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2))\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)(c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2)))}{2c_1\cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2\sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$d \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2))\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)(c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2)))}{(2c_1\cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2\sin\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} \right) \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}$$

Integrating gives

$$\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} = \int \frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2))\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)(c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2)))}{(2c_1\cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2\sin\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

$$= \int \frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2))\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)(c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2)))}{(2c_1\cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2\sin\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}$ gives the final solution

$$y = \frac{(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2))\cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right)(c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2)))}{(2c_1\cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2\sin\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{(c_1 + c_2 \tan(\frac{\sqrt{3}x}{2})) e^{-\frac{\sqrt{3} \arctan(\tan(\frac{\sqrt{3}x}{2}))}{3}} \left(\int \frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2)) \cos(\frac{\sqrt{3}x}{2}) + \sin(\frac{\sqrt{3}x}{2})) (c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2))}{(2c_1 \cos(\frac{\sqrt{3}x}{2}) + 2c_2 \sin(\frac{\sqrt{3}x}{2})) (c_1 + c_2 \tan(\frac{\sqrt{3}x}{2}))} dx \right)}{\sqrt{\sec(\frac{\sqrt{3}x}{2})^2}}$$

The constants can be merged to give

$$y = \frac{(c_1 + c_2 \tan(\frac{\sqrt{3}x}{2})) e^{-\frac{\sqrt{3} \arctan(\tan(\frac{\sqrt{3}x}{2}))}{3}} \left(\int \frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2)) \cos(\frac{\sqrt{3}x}{2}) + \sin(\frac{\sqrt{3}x}{2})) (c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2))}{(2c_1 \cos(\frac{\sqrt{3}x}{2}) + 2c_2 \sin(\frac{\sqrt{3}x}{2})) (c_1 + c_2 \tan(\frac{\sqrt{3}x}{2}))} dx \right)}{\sqrt{\sec(\frac{\sqrt{3}x}{2})^2}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{(c_1 + c_2 \tan(\frac{\sqrt{3}x}{2})) e^{-\frac{\sqrt{3} \arctan(\tan(\frac{\sqrt{3}x}{2}))}{3}} \left(\int \frac{((-xc_2(x-1)\sqrt{3} + c_1(x^2 + 3x - 2)) \cos(\frac{\sqrt{3}x}{2}) + \sin(\frac{\sqrt{3}x}{2})) (c_1x(x-1)\sqrt{3} + c_2(x^2 + 3x - 2))}{(2c_1 \cos(\frac{\sqrt{3}x}{2}) + 2c_2 \sin(\frac{\sqrt{3}x}{2})) (c_1 + c_2 \tan(\frac{\sqrt{3}x}{2}))} dx \right)}{\sqrt{\sec(\frac{\sqrt{3}x}{2})^2}}$$




Figure 2.71: Slope field plot
 $y'' + y' + y = x^2 + x + 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = x^2 + x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^2 + x + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} (x^2+x+1) \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} (x^2+x+1) \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = x(x-1)$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + x(x-1)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)

Leaf size : 37

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = x^2+x+1,
        y(x),singsol=all)

```

$$y = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + x^2 - x$$

Mathematica DSolve solution

Solving time : 0.022 (sec)

Leaf size : 54

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==1+x+x^2,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow e^{-x/2} \left(e^{x/2} (x-1)x + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

2.1.28 problem 28

Solved as second order linear constant coeff ode	213
Solved as second order ode using Kovacic algorithm	215
Solved as second order ode adjoint method	219
Maple step by step solution	223
Maple trace	224
Maple dsolve solution	224
Mathematica DSolve solution	224

Internal problem ID [8775]

Book : Second order enumerated odes

Section : section 1

Problem number : 28

Date solved : Thursday, December 12, 2024 at 09:44:48 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' + y' + y = x^3 + x^2 + x + 1$$

Solved as second order linear constant coeff ode

Time used: 0.173 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = x^3 + x^2 + x + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^3 + x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_4 x^3 + A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_4 x^3 + A_3 x^2 + 3x^2 A_4 + A_2 x + 2x A_3 + 6x A_4 + A_1 + A_2 + 2A_3 = x^3 + x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 6, A_2 = -1, A_3 = -2, A_4 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^3 - 2x^2 - x + 6$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (x^3 - 2x^2 - x + 6) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^3 - 2x^2 - x + 6 + e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$




Figure 2.72: Slope field plot
 $y'' + y' + y = x^3 + x^2 + x + 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.268 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1$$

$$C = 1$$

(3)

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.32: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^3 + x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[1, x, x^2, x^3]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_4x^3 + A_3x^2 + A_2x + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_4x^3 + A_3x^2 + 3x^2A_4 + A_2x + 2xA_3 + 6xA_4 + A_1 + A_2 + 2A_3 = x^3 + x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 6, A_2 = -1, A_3 = -2, A_4 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^3 - 2x^2 - x + 6$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (x^3 - 2x^2 - x + 6) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} + x^3 - 2x^2 - x + 6$$




Figure 2.73: Slope field plot
 $y'' + y' + y = x^3 + x^2 + x + 1$

Solved as second order ode adjoint method

Time used: 32.128 (sec)

In normal form the ode

$$y'' + y' + y = x^3 + x^2 + x + 1 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= x^3 + x^2 + x + 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (\xi(x)) &= 0 \\ \xi''(x) - \xi'(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(\frac{e^{\frac{x}{2}} (c_1 \cos(\frac{\sqrt{3}x}{2}) + c_2 \sin(\frac{\sqrt{3}x}{2}))}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sin(\frac{\sqrt{3}x}{2}) \sqrt{3}}{2} + \frac{c_2 \sqrt{3} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \right) = \frac{e^{-\frac{x}{2}} \left(c_1 \left(\left(\frac{1}{2} x^3 + \frac{3}{2} x \right) \right) \right)}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) + 2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right)} \\ p(x) &= \frac{(-c_2(x^3 - 2x^2 - x + 6) \sqrt{3} + c_1(x-1)(x^2 + 5x - 4)) \cos \left(\frac{\sqrt{3}x}{2} \right) + (c_1(x^3 - 2x^2 - x + 6) \sqrt{3} + c_2(x^3 - 2x^2 - x + 6)) \sin \left(\frac{\sqrt{3}x}{2} \right)}{2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) + 2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right)}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2\sqrt{3}-c_1)\cos\left(\frac{\sqrt{3}x}{2}\right)-\sin\left(\frac{\sqrt{3}x}{2}\right)(c_1\sqrt{3}+c_2)}{2c_2\sin\left(\frac{\sqrt{3}x}{2}\right)+2c_1\cos\left(\frac{\sqrt{3}x}{2}\right)} dx} \\ &= \frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}\end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y)$$

$$= (\mu) \left(\frac{(-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4))\cos\left(\frac{\sqrt{3}x}{2}\right) + (c_1(x^3 - 2x^2 - x + 6)\sqrt{3} + c_2(x^3 - 2x^2 - x + 6))\sin\left(\frac{\sqrt{3}x}{2}\right)}{2c_2\sin\left(\frac{\sqrt{3}x}{2}\right) + 2c_1\cos\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$\frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \left(\frac{(-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4))\cos\left(\frac{\sqrt{3}x}{2}\right) + (c_1(x^3 - 2x^2 - x + 6)\sqrt{3} + c_2(x^3 - 2x^2 - x + 6))\sin\left(\frac{\sqrt{3}x}{2}\right)}{2c_2\sin\left(\frac{\sqrt{3}x}{2}\right) + 2c_1\cos\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$d \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{((-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4))\cos\left(\frac{\sqrt{3}x}{2}\right) + (c_1(x^3 - 2x^2 - x + 6)\sqrt{3} + c_2(x^3 - 2x^2 - x + 6))\sin\left(\frac{\sqrt{3}x}{2}\right))}{(2c_2\sin\left(\frac{\sqrt{3}x}{2}\right) + 2c_1\cos\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} \right)$$

Integrating gives

$$\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} = \int \frac{((-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4))\cos\left(\frac{\sqrt{3}x}{2}\right) + (c_1(x^3 - 2x^2 - x + 6)\sqrt{3} + c_2(x^3 - 2x^2 - x + 6))\sin\left(\frac{\sqrt{3}x}{2}\right))}{(2c_2\sin\left(\frac{\sqrt{3}x}{2}\right) + 2c_1\cos\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

$$= \int \frac{((-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4))\cos\left(\frac{\sqrt{3}x}{2}\right) + (c_1(x^3 - 2x^2 - x + 6)\sqrt{3} + c_2(x^3 - 2x^2 - x + 6))\sin\left(\frac{\sqrt{3}x}{2}\right))}{(2c_2\sin\left(\frac{\sqrt{3}x}{2}\right) + 2c_1\cos\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}$ gives the final solution

$$y = \frac{(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{\sqrt{3}\arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{((-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4))\cos\left(\frac{\sqrt{3}x}{2}\right) + (c_1(x^3 - 2x^2 - x + 6)\sqrt{3} + c_2(x^3 - 2x^2 - x + 6))\sin\left(\frac{\sqrt{3}x}{2}\right))}{(2c_2\sin\left(\frac{\sqrt{3}x}{2}\right) + 2c_1\cos\left(\frac{\sqrt{3}x}{2}\right))(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

y

$$\left(c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right) \right) e^{-\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}} \left(\int \frac{\left((-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4)) \cos \left(\frac{\sqrt{3}x}{2} \right) + c_1(x^3 - 2x^2 - x + 6)\sqrt{3} \right)}{\left(2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) + 2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) \right)} dx \right)$$

$$\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2}$$

The constants can be merged to give

y

$$\left(c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right) \right) e^{-\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}} \left(\int \frac{\left((-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4)) \cos \left(\frac{\sqrt{3}x}{2} \right) + c_1(x^3 - 2x^2 - x + 6)\sqrt{3} \right)}{\left(2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) + 2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) \right)} dx \right)$$

$$\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

y

$$\left(c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right) \right) e^{-\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}} \left(\int \frac{\left((-c_2(x^3 - 2x^2 - x + 6)\sqrt{3} + c_1(x-1)(x^2 + 5x - 4)) \cos \left(\frac{\sqrt{3}x}{2} \right) + c_1(x^3 - 2x^2 - x + 6)\sqrt{3} \right)}{\left(2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) + 2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) \right)} dx \right)$$

$$\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2}$$




Figure 2.74: Slope field plot
 $y'' + y' + y = x^3 + x^2 + x + 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = x^3 + x^2 + x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^3 + x^2 + x + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}}(x+1)(x^2+1) \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}}(x+1)(x^2+1) \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = x^3 - 2x^2 - x + 6$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + x^3 - 2x^2 - x + 6$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)

Leaf size : 43

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = x^3+x^2+x+1,
        y(x),singsol=all)

```

$$y = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + x^3 - 2x^2 - x + 6$$

Mathematica DSolve solution

Solving time : 0.024 (sec)

Leaf size : 60

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==1+x+x^2+x^3,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow x^3 - 2x^2 - x + c_2 e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right) + 6$$

2.1.29 problem 29

Solved as second order linear constant coeff ode	225
Solved as second order ode using Kovacic algorithm	227
Solved as second order ode adjoint method	231
Maple step by step solution	234
Maple trace	235
Maple dsolve solution	235
Mathematica DSolve solution	236

Internal problem ID [8776]

Book : Second order enumerated odes

Section : section 1

Problem number : 29

Date solved : Thursday, December 12, 2024 at 09:45:22 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' + y' + y = \sin(x)$$

Solved as second order linear constant coeff ode

Time used: 0.171 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \right) + (-\cos(x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\cos(x) + e^{-\frac{x}{2}} \left(c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$




Figure 2.75: Slope field plot
 $y'' + y' + y = \sin(x)$

Solved as second order ode using Kovacic algorithm

Time used: 0.267 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.34: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin \left(\frac{\sqrt{3} x}{2} \right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (-\cos(x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} - \cos(x)$$




Figure 2.76: Slope field plot
 $y'' + y' + y = \sin(x)$

Solved as second order ode adjoint method

Time used: 40.572 (sec)

In normal form the ode

$$y'' + y' + y = \sin(x) \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= \sin(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (\xi(x)) &= 0 \\ \xi''(x) - \xi'(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(\frac{e^{\frac{x}{2}} (c_1 \cos(\frac{\sqrt{3}x}{2}) + c_2 \sin(\frac{\sqrt{3}x}{2}))}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin(\frac{\sqrt{3}x}{2})}{2} + \frac{c_2 \sqrt{3} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right) = \frac{e^{-\frac{x}{2}} \left(c_1 \frac{\left((-\frac{\sqrt{3}}{2} - 1) e^{\frac{x}{2}} \right)}{\frac{1}{4} + \dots} \right)}{\dots}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \\ p(x) &= \frac{(\cos(x) \sqrt{3} c_2 - c_1 (\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2 (\cos(x) - 2 \sin(x)))}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2 \sqrt{3} - c_1) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} dx} \\ &= \frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y)$$

$$= (\mu) \left(\frac{((\cos(x) \sqrt{3} c_2 - c_1(\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2(\cos(x) - 2 \sin(x))))}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$\frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \left(\frac{((\cos(x) \sqrt{3} c_2 - c_1(\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2(\cos(x) - 2 \sin(x))))}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$d \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{((\cos(x) \sqrt{3} c_2 - c_1(\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2(\cos(x) - 2 \sin(x))))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} \right)$$

Integrating gives

$$\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} = \int \frac{((\cos(x) \sqrt{3} c_2 - c_1(\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2(\cos(x) - 2 \sin(x))))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

$$= \int \frac{((\cos(x) \sqrt{3} c_2 - c_1(\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2(\cos(x) - 2 \sin(x))))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}$ gives the final solution

$$y = \frac{(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{((\cos(x) \sqrt{3} c_2 - c_1(\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2(\cos(x) - 2 \sin(x))))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

y

$$= \frac{(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{((\cos(x) \sqrt{3} c_2 - c_1(\cos(x) - 2 \sin(x))) \cos\left(\frac{\sqrt{3}x}{2}\right) - \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1 \sqrt{3} \cos(x) + c_2(\cos(x) - 2 \sin(x))))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

The constants can be merged to give

y

$$\left(c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right) \right) e^{-\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}} \left(\int \frac{\left((\cos(x)\sqrt{3}c_2 - c_1(\cos(x) - 2\sin(x))) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1\sqrt{3} \cos(x) + c_2(\cos(x) - 2\sin(x))) \right)}{(2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)) (c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right))} dx \right)$$

$$\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

y

$$\left(c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right) \right) e^{-\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}} \left(\int \frac{\left((\cos(x)\sqrt{3}c_2 - c_1(\cos(x) - 2\sin(x))) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1\sqrt{3} \cos(x) + c_2(\cos(x) - 2\sin(x))) \right)}{(2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)) (c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right))} dx \right)$$

$$\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2}$$




Figure 2.77: Slope field plot $y'' + y' + y = \sin(x)$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = \sin(x)$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = -\cos(x)$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) - \cos(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.006 (sec)

Leaf size : 35

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = sin(x),
        y(x),singsol=all)

```

$$y = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 - \cos(x)$$

Mathematica DSolve solution

Solving time : 0.297 (sec)

Leaf size : 53

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==Sin[x],{}}],  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{-x/2} \left(-e^{x/2} \cos(x) + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

2.1.30 problem 30

Solved as second order linear constant coeff ode	237
Solved as second order ode using Kovacic algorithm	239
Solved as second order ode adjoint method	243
Maple step by step solution	246
Maple trace	247
Maple dsolve solution	247
Mathematica DSolve solution	248

Internal problem ID [8777]

Book : Second order enumerated odes

Section : section 1

Problem number : 30

Date solved : Thursday, December 12, 2024 at 09:46:03 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' + y' + y = \cos(x)$$

Solved as second order linear constant coeff ode

Time used: 0.171 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \right) + (\sin(x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \sin(x) + e^{-\frac{x}{2}} \left(c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$




Figure 2.78: Slope field plot
 $y'' + y' + y = \cos(x)$

Solved as second order ode using Kovacic algorithm

Time used: 0.267 (sec)

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.36: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3} x}{2} \right)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3} x}{2} \right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin \left(\frac{\sqrt{3} x}{2} \right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (\sin(x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}}\sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} + \sin(x)$$




Figure 2.79: Slope field plot
 $y'' + y' + y = \cos(x)$

Solved as second order ode adjoint method

Time used: 71.785 (sec)

In normal form the ode

$$y'' + y' + y = \cos(x) \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= \cos(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (\xi(x)) &= 0 \\ \xi''(x) - \xi'(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(1)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{\left(\frac{e^{\frac{x}{2}} (c_1 \cos(\frac{\sqrt{3}x}{2}) + c_2 \sin(\frac{\sqrt{3}x}{2}))}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{3} \sin(\frac{\sqrt{3}x}{2})}{2} + \frac{c_2 \sqrt{3} \cos(\frac{\sqrt{3}x}{2})}{2} \right) \right) e^{-\frac{x}{2}}}{c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \right) = \frac{e^{-\frac{x}{2}} \left(c_1 \frac{e^{\frac{x}{2}} \cos(x) (-\frac{1}{2} + 2(-1 + \dots))}{\frac{1}{2} + 2(-1 + \dots)} \right)}{\dots}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \\ p(x) &= \frac{\left(-c_2 \sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \cos \left(\frac{\sqrt{3}x}{2} \right) + \sin \left(\frac{\sqrt{3}x}{2} \right) \left(c_1 \sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2 \sqrt{3} - c_1) \cos \left(\frac{\sqrt{3}x}{2} \right) - \sin \left(\frac{\sqrt{3}x}{2} \right) (c_1 \sqrt{3} + c_2)}{2c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + 2c_2 \sin \left(\frac{\sqrt{3}x}{2} \right)} dx} \\ &= \frac{\sqrt{\sec \left(\frac{\sqrt{3}x}{2} \right)^2} e^{\frac{\sqrt{3} \arctan \left(\tan \left(\frac{\sqrt{3}x}{2} \right) \right)}{3}}}{c_1 + c_2 \tan \left(\frac{\sqrt{3}x}{2} \right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y)$$

$$= (\mu) \left(\frac{\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(c_1\sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \right)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$\frac{d}{dx} \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right) \left(\frac{\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(c_1\sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \right)}{2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$d \left(\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} \right)$$

$$= \left(\frac{\left(\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(c_1\sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \right)}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} \right)$$

Integrating gives

$$\frac{y \sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)} = \int \frac{\left(\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(c_1\sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \right)}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx$$

$$= \int \frac{\left(\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(c_1\sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \right)}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx$$

Dividing throughout by the integrating factor $\frac{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2} e^{\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}}}{c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right)}$ gives the final solution

$$y = \frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left(\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(c_1\sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \right)}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left(\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2} \right)) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) \left(c_1\sqrt{3} \sin(x) + 2c_2 \left(\cos(x) + \frac{\sin(x)}{2} \right) \right) \right)}{\left(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right)} dx \right)}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

The constants can be merged to give

y

$$\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2}\right)) \right) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} \sin(x) + 2c_2 \cos(x))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right) \frac{1}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

y

$$\left(c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right) \right) e^{-\frac{\sqrt{3} \arctan\left(\tan\left(\frac{\sqrt{3}x}{2}\right)\right)}{3}} \left(\int \frac{\left((-c_2\sqrt{3} \sin(x) + 2c_1 \left(\cos(x) + \frac{\sin(x)}{2}\right)) \right) \cos\left(\frac{\sqrt{3}x}{2}\right) + \sin\left(\frac{\sqrt{3}x}{2}\right) (c_1\sqrt{3} \sin(x) + 2c_2 \cos(x))}{(2c_1 \cos\left(\frac{\sqrt{3}x}{2}\right) + 2c_2 \sin\left(\frac{\sqrt{3}x}{2}\right)) (c_1 + c_2 \tan\left(\frac{\sqrt{3}x}{2}\right))} dx \right) \frac{1}{\sqrt{\sec\left(\frac{\sqrt{3}x}{2}\right)^2}}$$




Figure 2.80: Slope field plot
 $y'' + y' + y = \cos(x)$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) + y(x) = \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \cos(x) \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \cos(x) \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = \sin(x)$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + C_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + \sin(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.008 (sec)

Leaf size : 33

```

dsolve(diff(diff(y(x),x),x)+diff(y(x),x)+y(x) = cos(x),
        y(x),singsol=all)

```

$$y = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + \sin(x)$$

Mathematica DSolve solution

Solving time : 0.588 (sec)

Leaf size : 50

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]+y[x]==Cos[x],{}}],  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \sin(x) + c_2 e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

2.1.31 problem 31

Solved as second order linear constant coeff ode	249
Solved as second order linear exact ode	251
Solved as second order missing y ode	253
Solved as second order integrable as is ode	254
Solved as second order integrable as is ode (ABC method)	255
Solved as second order ode using Kovacic algorithm	257
Solved as second order ode adjoint method	260
Maple step by step solution	263
Maple trace	263
Maple dsolve solution	264
Mathematica DSolve solution	264

Internal problem ID [8778]

Book : Second order enumerated odes

Section : section 1

Problem number : 31

Date solved : Thursday, December 12, 2024 at 09:47:16 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y'' + y' = 1$$

Solved as second order linear constant coeff ode

Time used: 0.115 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + (x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_1 + c_2 e^{-x}$$




Figure 2.81: Slope field plot
 $y'' + y' = 1$

Solved as second order linear exact ode

Time used: 0.102 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= 1 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int 1 dx$$

We now have a first order ode to solve which is

$$y' + y = x + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= x + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= ((x + c_1) e^x) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int (x + c_1) e^x dx \\ &= (x + c_1 - 1) e^x + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = c_1 + x - 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 + x - 1 + c_2 e^{-x}$$




Figure 2.82: Slope field plot
 $y'' + y' = 1$

Solved as second order missing y ode

Time used: 0.149 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating gives

$$\int \frac{1}{1-p} dp = dx$$

$$-\ln(-1+p) = x + c_1$$

Singular solutions are found by solving

$$1 - p = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 1$$

Solving for $p(x)$ gives

$$p(x) = e^{-x-c_1} + 1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 1 dx$$

$$y = x + c_2$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = e^{-x-c_1} + 1$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int e^{-x-c_1} + 1 dx$$

$$y = x - e^{-x-c_1} + c_3$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_2$$

$$y = x - e^{-x-c_1} + c_3$$




Figure 2.83: Slope field plot
 $y'' + y' = 1$

Solved as second order integrable as is ode

Time used: 0.049 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int 1 dx$$

$$y' + y = x + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = x + c_1$$

The integrating factor μ is

$$\mu = e^{\int q dx}$$

$$= e^{\int 1 dx}$$

$$= e^x$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= ((x + c_1) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int (x + c_1) e^x dx \\ &= (x + c_1 - 1) e^x + c_2\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = c_1 + x - 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 + x - 1 + c_2 e^{-x}$$




Figure 2.84: Slope field plot
 $y'' + y' = 1$

Solved as second order integrable as is ode (ABC method)

Time used: 0.045 (sec)

Writing the ode as

$$y'' + y' = 1$$

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned}\int (y'' + y') dx &= \int 1 dx \\ y' + y &= x + c_1\end{aligned}$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= 1 \\p(x) &= x + c_1\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\&= e^{\int 1 dx} \\&= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= ((x + c_1) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int (x + c_1) e^x dx \\ &= (x + c_1 - 1) e^x + c_2\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = c_1 + x - 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.85: Slope field plot
 $y'' + y' = 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.063 (sec)

Writing the ode as

$$y'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.38: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 (e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 (e^{-x} (e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + (x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 + x$$




Figure 2.86: Slope field plot
 $y'' + y' = 1$

Solved as second order ode adjoint method

Time used: 0.456 (sec)

In normal form the ode

$$y'' + y' = 1 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 1 \\ \lambda_2 &= 0\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}\xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x}\end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = \frac{c_2 x + c_1 e^x}{c_1 e^x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= \frac{c_2}{c_1 e^x + c_2} \\ p(x) &= \frac{c_2 x + c_1 e^x}{c_1 e^x + c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{c_2 x + c_1 e^x}{c_1 e^x + c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{e^x}{c_1 e^x + c_2} \right) \left(\frac{c_2 x + c_1 e^x}{c_1 e^x + c_2} \right) \\ d \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{(c_2 x + c_1 e^x) e^x}{(c_1 e^x + c_2)^2} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y e^x}{c_1 e^x + c_2} &= \int \frac{(c_2 x + c_1 e^x) e^x}{(c_1 e^x + c_2)^2} dx \\ &= \frac{c_2}{c_1 (c_1 e^x + c_2)} + \frac{x e^x}{c_1 e^x + c_2} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = \frac{c_2(c_3 c_1 + 1) e^{-x} + c_1(c_3 c_1 + x)}{c_1}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{c_2(c_3 c_1 + 1) e^{-x} + c_1(c_3 c_1 + x)}{c_1}$$

The constants can be merged to give

$$y = \frac{c_2(c_1 + 1) e^{-x} + c_1(x + c_1)}{c_1}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_2(c_1 + 1) e^{-x} + c_1(x + c_1)}{c_1}$$




Figure 2.87: Slope field plot
 $y'' + y' = 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x dx \right) + \int 1 dx$$

- Compute integrals

$$y_p(x) = x - 1$$

- Substitute particular solution into general solution to ODE

$$y(x) = C1 e^{-x} + C2 + x - 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 14

```
dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = 1,  
        y(x),singsol=all)
```

$$y = -c_1 e^{-x} + x + c_2$$

Mathematica DSolve solution

Solving time : 0.013 (sec)

Leaf size : 18

```
DSolve[{D[y[x],{x,2}]+D[y[x],x]==1,{}},  
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x - c_1 e^{-x} + c_2$$

2.1.32 problem 32

Solved as second order linear constant coeff ode	265
Solved as second order linear exact ode	267
Solved as second order missing y ode	269
Solved as second order integrable as is ode	271
Solved as second order integrable as is ode (ABC method)	272
Solved as second order ode using Kovacic algorithm	273
Solved as second order ode adjoint method	276
Maple step by step solution	279
Maple trace	280
Maple dsolve solution	280
Mathematica DSolve solution	280

Internal problem ID [8779]

Book : Second order enumerated odes

Section : section 1

Problem number : 32

Date solved : Thursday, December 12, 2024 at 09:47:18 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y'' + y' = x$$

Solved as second order linear constant coeff ode

Time used: 0.126 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2xA_2 + A_1 + 2A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -1, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{2}x^2 - x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + \left(\frac{1}{2}x^2 - x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} - x + c_1 + c_2 e^{-x}$$




Figure 2.88: Slope field plot
 $y'' + y' = x$

Solved as second order linear exact ode

Time used: 0.110 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= x \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int x dx$$

We now have a first order ode to solve which is

$$y' + y = \frac{x^2}{2} + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= \frac{x^2}{2} + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{x^2}{2} + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{x^2}{2} + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{x^2}{2} + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{x^2}{2} + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1 - 2x + 2) e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$




Figure 2.89: Slope field plot
 $y'' + y' = x$

Solved as second order missing y ode

Time used: 0.099 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - x = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = x$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= \mu p \\ \frac{d}{dx}(\mu p) &= (\mu)(x) \\ \frac{d}{dx}(p e^x) &= (e^x)(x) \\ d(p e^x) &= (x e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}p e^x &= \int x e^x dx \\ &= (x - 1) e^x + c_1\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$p(x) = x - 1 + c_1 e^{-x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x - 1 + c_1 e^{-x}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned}\int dy &= \int x - 1 + c_1 e^{-x} dx \\ y &= -x + \frac{x^2}{2} - c_1 e^{-x} + c_2\end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -x + \frac{x^2}{2} - c_1 e^{-x} + c_2$$




Figure 2.90: Slope field plot
 $y'' + y' = x$

Solved as second order integrable as is ode

Time used: 0.128 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int x dx$$

$$y' + y = \frac{x^2}{2} + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{x^2}{2} + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{x^2}{2} + c_1 \right)$$

$$\frac{d}{dx}(y e^x) = (e^x) \left(\frac{x^2}{2} + c_1 \right)$$

$$d(y e^x) = \left(\left(\frac{x^2}{2} + c_1 \right) e^x \right) dx$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{x^2}{2} + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1 - 2x + 2) e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$




Figure 2.91: Slope field plot
 $y'' + y' = x$

Solved as second order integrable as is ode (ABC method)

Time used: 0.046 (sec)

Writing the ode as

$$y'' + y' = x$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int x dx$$

$$y' + y = \frac{x^2}{2} + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{x^2}{2} + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{x^2}{2} + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{x^2}{2} + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{x^2}{2} + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{x^2}{2} + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1 - 2x + 2) e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.92: Slope field plot
 $y'' + y' = x$

Solved as second order ode using Kovacic algorithm

Time used: 0.061 (sec)

Writing the ode as

$$y'' + y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = y e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = r z(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.40: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2xA_2 + A_1 + 2A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -1, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{2}x^2 - x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + \left(\frac{1}{2}x^2 - x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 + \frac{x^2}{2} - x$$




Figure 2.93: Slope field plot
 $y'' + y' = x$

Solved as second order ode adjoint method

Time used: 0.508 (sec)

In normal form the ode

$$y'' + y' = x \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= x \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = \frac{\frac{c_2 x^2}{2} + c_1 (x e^x - e^x)}{c_1 e^x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= \frac{c_2}{c_1 e^x + c_2} \\ p(x) &= \frac{2c_1(x-1)e^x + c_2 x^2}{2c_1 e^x + 2c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{2c_1(x-1)e^x + c_2 x^2}{2c_1 e^x + 2c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{e^x}{c_1 e^x + c_2} \right) \left(\frac{2c_1(x-1)e^x + c_2 x^2}{2c_1 e^x + 2c_2} \right) \\ d \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{(2c_1(x-1)e^x + c_2 x^2) e^x}{(2c_1 e^x + 2c_2)(c_1 e^x + c_2)} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y e^x}{c_1 e^x + c_2} &= \int \frac{(2c_1(x-1)e^x + c_2 x^2) e^x}{(2c_1 e^x + 2c_2)(c_1 e^x + c_2)} dx \\ &= \frac{e^x - x e^x + \frac{e^x x^2}{2}}{c_1 e^x + c_2} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = c_1 c_3 + \frac{x^2}{2} - x + c_2 c_3 e^{-x} + 1$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_1 c_3 + \frac{x^2}{2} - x + c_2 c_3 e^{-x} + 1$$

The constants can be merged to give

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 - x + 1 + c_2 e^{-x}$$




Figure 2.94: Slope field plot
 $y'' + y' = x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = x$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x x dx \right) + \int x dx$$
- Compute integrals
$$y_p(x) = 1 - x + \frac{1}{2}x^2$$
- Substitute particular solution into general solution to ODE
$$y(x) = C_1 e^{-x} + C_2 + 1 - x + \frac{x^2}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)
 Leaf size : 21

```

dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = x,
        y(x),singsol=all)

```

$$y = \frac{x^2}{2} - c_1 e^{-x} - x + c_2$$

Mathematica DSolve solution

Solving time : 0.038 (sec)
 Leaf size : 27

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]==x,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{x^2}{2} - x - c_1 e^{-x} + c_2$$

2.1.33 problem 33

Solved as second order linear constant coeff ode	281
Solved as second order linear exact ode	283
Solved as second order missing y ode	285
Solved as second order integrable as is ode	287
Solved as second order integrable as is ode (ABC method)	288
Solved as second order ode using Kovacic algorithm	289
Solved as second order ode adjoint method	292
Maple step by step solution	295
Maple trace	296
Maple dsolve solution	296
Mathematica DSolve solution	296

Internal problem ID [8780]

Book : Second order enumerated odes

Section : section 1

Problem number : 33

Date solved : Thursday, December 12, 2024 at 09:47:20 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y'' + y' = 1 + x$$

Solved as second order linear constant coeff ode

Time used: 0.129 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = 1 + x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2xA_2 + A_1 + 2A_2 = 1 + x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + \left(\frac{x^2}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$




Figure 2.95: Slope field plot
 $y'' + y' = 1 + x$

Solved as second order linear exact ode

Time used: 0.110 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= 1 + x \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int 1 + x dx$$

We now have a first order ode to solve which is

$$y' + y = x + \frac{1}{2}x^2 + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= x + \frac{1}{2}x^2 + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(x + \frac{1}{2}x^2 + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(x + \frac{1}{2}x^2 + c_1 \right) \\ d(y e^x) &= \left(\left(x + \frac{1}{2}x^2 + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(x + \frac{1}{2}x^2 + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1)e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$




Figure 2.96: Slope field plot
 $y'' + y' = 1 + x$

Solved as second order missing y ode

Time used: 0.101 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - 1 - x = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = 1 + x$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= \mu p \\ \frac{d}{dx}(\mu p) &= (\mu)(1+x) \\ \frac{d}{dx}(p e^x) &= (e^x)(1+x) \\ d(p e^x) &= ((1+x) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}p e^x &= \int (1+x) e^x dx \\ &= x e^x + c_1\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$p(x) = x + c_1 e^{-x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x + c_1 e^{-x}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned}\int dy &= \int x + c_1 e^{-x} dx \\ y &= \frac{x^2}{2} - c_1 e^{-x} + c_2\end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} - c_1 e^{-x} + c_2$$




Figure 2.97: Slope field plot
 $y'' + y' = 1 + x$

Solved as second order integrable as is ode

Time used: 0.119 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (1 + x) dx$$

$$y' + y = x + \frac{1}{2}x^2 + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = x + \frac{1}{2}x^2 + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(x + \frac{1}{2}x^2 + c_1 \right)$$

$$\frac{d}{dx}(y e^x) = (e^x) \left(x + \frac{1}{2}x^2 + c_1 \right)$$

$$d(y e^x) = \left(\left(x + \frac{1}{2}x^2 + c_1 \right) e^x \right) dx$$

Integrating gives

$$\begin{aligned}y e^x &= \int \left(x + \frac{1}{2}x^2 + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1) e^x}{2} + c_2\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$




Figure 2.98: Slope field plot

$$y'' + y' = 1 + x$$

Solved as second order integrable as is ode (ABC method)

Time used: 0.050 (sec)

Writing the ode as

$$y'' + y' = 1 + x$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (1 + x) dx$$

$$y' + y = x + \frac{1}{2}x^2 + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = x + \frac{1}{2}x^2 + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(x + \frac{1}{2}x^2 + c_1 \right)$$

$$\frac{d}{dx}(y e^x) = (e^x) \left(x + \frac{1}{2}x^2 + c_1 \right)$$

$$d(y e^x) = \left(\left(x + \frac{1}{2}x^2 + c_1 \right) e^x \right) dx$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(x + \frac{1}{2}x^2 + c_1 \right) e^x dx \\ &= \frac{(x^2 + 2c_1) e^x}{2} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.99: Slope field plot
 $y'' + y' = 1 + x$

Solved as second order ode using Kovacic algorithm

Time used: 0.068 (sec)

Writing the ode as

$$y'' + y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 1 \\ t &= 4\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.42: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2xA_2 + A_1 + 2A_2 = 1 + x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + \left(\frac{x^2}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 + \frac{x^2}{2}$$




Figure 2.100: Slope field plot
 $y'' + y' = 1 + x$

Solved as second order ode adjoint method

Time used: 0.498 (sec)

In normal form the ode

$$y'' + y' = 1 + x \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 1 + x \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = \frac{c_2 x + c_1 e^x + \frac{c_2 x^2}{2} + c_1 (x e^x - e^x)}{c_1 e^x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= \frac{c_2}{c_1 e^x + c_2} \\ p(x) &= \frac{x(2c_1 e^x + c_2 x + 2c_2)}{2c_1 e^x + 2c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{x(2c_1 e^x + c_2 x + 2c_2)}{2c_1 e^x + 2c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{e^x}{c_1 e^x + c_2} \right) \left(\frac{x(2c_1 e^x + c_2 x + 2c_2)}{2c_1 e^x + 2c_2} \right) \\ d \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{x(2c_1 e^x + c_2 x + 2c_2) e^x}{(2c_1 e^x + 2c_2)(c_1 e^x + c_2)} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y e^x}{c_1 e^x + c_2} &= \int \frac{x(2c_1 e^x + c_2 x + 2c_2) e^x}{(2c_1 e^x + 2c_2)(c_1 e^x + c_2)} dx \\ &= \frac{x^2 e^x}{2c_1 e^x + 2c_2} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = c_1 c_3 + \frac{x^2}{2} + c_2 c_3 e^{-x}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_1 c_3 + \frac{x^2}{2} + c_2 c_3 e^{-x}$$

The constants can be merged to give

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{2} + c_1 + c_2 e^{-x}$$




Figure 2.101: Slope field plot
 $y'' + y' = 1 + x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = x + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x(x+1) dx \right) + \int (x+1) dx$$
- Compute integrals
$$y_p(x) = \frac{x^2}{2}$$
- Substitute particular solution into general solution to ODE
$$y(x) = C_1 e^{-x} + C_2 + \frac{x^2}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)+_a+1, _b(_a)`
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

*** Subleve

Maple dsolve solution

Solving time : 0.001 (sec)
Leaf size : 18

```

dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = x+1,
        y(x),singsol=all)

```

$$y = \frac{x^2}{2} - c_1 e^{-x} + c_2$$

Mathematica DSolve solution

Solving time : 0.032 (sec)
Leaf size : 24

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]==1+x,{]},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{x^2}{2} - c_1 e^{-x} + c_2$$

2.1.34 problem 34

Solved as second order linear constant coeff ode 297
 Solved as second order linear exact ode 299
 Solved as second order missing y ode 301
 Solved as second order integrable as is ode 303
 Solved as second order integrable as is ode (ABC method) 304
 Solved as second order ode using Kovacic algorithm 305
 Solved as second order ode adjoint method 308
 Maple step by step solution 311
 Maple trace 312
 Maple dsolve solution 312
 Mathematica DSolve solution 312

Internal problem ID [8781]

Book : Second order enumerated odes

Section : section 1

Problem number : 34

Date solved : Thursday, December 12, 2024 at 09:47:22 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y'' + y' = x^2 + x + 1$$

Solved as second order linear constant coeff ode

Time used: 0.135 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = x^2 + x + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_3 x^3 + A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3x^2 A_3 + 2x A_2 + 6x A_3 + A_1 + 2A_2 = x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 2, A_2 = -\frac{1}{2}, A_3 = \frac{1}{3} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + \left(\frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^3}{3} - \frac{x^2}{2} + 2x + c_1 + c_2 e^{-x}$$




Figure 2.102: Slope field plot
 $y'' + y' = x^2 + x + 1$

Solved as second order linear exact ode

Time used: 0.113 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= x^2 + x + 1 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int x^2 + x + 1 dx$$

We now have a first order ode to solve which is

$$y' + y = \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x dx \\ &= \frac{(2x^3 - 3x^2 + 6c_1 + 12x - 12) e^x}{6} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^3}{3} - \frac{x^2}{2} + c_1 + 2x - 2 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^3}{3} - \frac{x^2}{2} + c_1 + 2x - 2 + c_2 e^{-x}$$




Figure 2.103: Slope field plot
 $y'' + y' = x^2 + x + 1$

Solved as second order missing y ode

Time used: 0.105 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - x^2 - x - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= x^2 + x + 1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= \mu p \\ \frac{d}{dx}(\mu p) &= (\mu)(x^2 + x + 1) \\ \frac{d}{dx}(p e^x) &= (e^x)(x^2 + x + 1) \\ d(p e^x) &= ((x^2 + x + 1) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}p e^x &= \int (x^2 + x + 1) e^x dx \\ &= (x^2 - x + 2) e^x + c_1\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$p(x) = x^2 - x + 2 + c_1 e^{-x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x^2 - x + 2 + c_1 e^{-x}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned}\int dy &= \int x^2 - x + 2 + c_1 e^{-x} dx \\ y &= 2x + \frac{x^3}{3} - c_1 e^{-x} - \frac{x^2}{2} + c_2\end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 2x + \frac{x^3}{3} - c_1 e^{-x} - \frac{x^2}{2} + c_2$$




Figure 2.104: Slope field plot

$$y'' + y' = x^2 + x + 1$$

Solved as second order integrable as is ode

Time used: 0.122 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (x^2 + x + 1) dx$$

$$y' + y = \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x dx \\ &= \frac{(2x^3 - 3x^2 + 6c_1 + 12x - 12) e^x}{6} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^3}{3} - \frac{x^2}{2} + c_1 + 2x - 2 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^3}{3} - \frac{x^2}{2} + c_1 + 2x - 2 + c_2 e^{-x}$$




Figure 2.105: Slope field plot

$$y'' + y' = x^2 + x + 1$$

Solved as second order integrable as is ode (ABC method)

Time used: 0.054 (sec)

Writing the ode as

$$y'' + y' = x^2 + x + 1$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (x^2 + x + 1) dx$$

$$y' + y = \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x dx \\ &= \frac{(2x^3 - 3x^2 + 6c_1 + 12x - 12) e^x}{6} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^3}{3} - \frac{x^2}{2} + c_1 + 2x - 2 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.106: Slope field plot
 $y'' + y' = x^2 + x + 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.073 (sec)

Writing the ode as

$$y'' + y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 1 \\ t &= 4\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.44: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_3 x^3 + A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3x^2 A_3 + 2xA_2 + 6xA_3 + A_1 + 2A_2 = x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 2, A_2 = -\frac{1}{2}, A_3 = \frac{1}{3} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + \left(\frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 + \frac{x^3}{3} - \frac{x^2}{2} + 2x$$




Figure 2.107: Slope field plot
 $y'' + y' = x^2 + x + 1$

Solved as second order ode adjoint method

Time used: 0.541 (sec)

In normal form the ode

$$y'' + y' = x^2 + x + 1 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= x^2 + x + 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = \frac{c_2 x + c_1 e^x + \frac{c_2 x^2}{2} + \frac{c_2 x^3}{3} + c_1 (e^x x - e^x) + c_1 (e^x x^2 - 2 e^x x + 2 e^x)}{c_1 e^x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= \frac{c_2}{c_1 e^x + c_2} \\ p(x) &= \frac{6c_1(x^2 - x + 2) e^x + 2c_2 x(x^2 + \frac{3}{2}x + 3)}{6c_1 e^x + 6c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{6c_1(x^2 - x + 2) e^x + 2c_2 x(x^2 + \frac{3}{2}x + 3)}{6c_1 e^x + 6c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{e^x}{c_1 e^x + c_2} \right) \left(\frac{6c_1(x^2 - x + 2) e^x + 2c_2 x(x^2 + \frac{3}{2}x + 3)}{6c_1 e^x + 6c_2} \right) \\ d \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{(6c_1(x^2 - x + 2) e^x + 2c_2 x(x^2 + \frac{3}{2}x + 3)) e^x}{(6c_1 e^x + 6c_2)(c_1 e^x + c_2)} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y e^x}{c_1 e^x + c_2} &= \int \frac{(6c_1(x^2 - x + 2) e^x + 2c_2 x(x^2 + \frac{3}{2}x + 3)) e^x}{(6c_1 e^x + 6c_2)(c_1 e^x + c_2)} dx \\ &= \frac{-2e^x + \frac{x^3 e^x}{3} + 2e^x x - \frac{e^x x^2}{2}}{c_1 e^x + c_2} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = \frac{x^3}{3} + c_1 c_3 - \frac{x^2}{2} + 2x + c_2 c_3 e^{-x} - 2$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{x^3}{3} + c_1 c_3 - \frac{x^2}{2} + 2x + c_2 c_3 e^{-x} - 2$$

The constants can be merged to give

$$y = \frac{x^3}{3} - \frac{x^2}{2} + c_1 + 2x - 2 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^3}{3} - \frac{x^2}{2} + c_1 + 2x - 2 + c_2 e^{-x}$$




Figure 2.108: Slope field plot
 $y'' + y' = x^2 + x + 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = x^2 + x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^2 + x + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int (x^2 + x + 1) e^x dx \right) + \int (x^2 + x + 1) dx$$

- Compute integrals

$$y_p(x) = -\frac{1}{2}x^2 + 2x - 2 + \frac{1}{3}x^3$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-x} + C_2 - \frac{x^2}{2} + 2x - 2 + \frac{x^3}{3}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = _a^2-_b(_a)+_a+1, _b(_a)` *** Sub
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)

Leaf size : 26

```

dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = x^2+x+1,
        y(x),singsol=all)

```

$$y = \frac{x^3}{3} - c_1 e^{-x} - \frac{x^2}{2} + 2x + c_2$$

Mathematica DSolve solution

Solving time : 0.101 (sec)

Leaf size : 34

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]==1+x+x^2,{x}],
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{x^3}{3} - \frac{x^2}{2} + 2x - c_1 e^{-x} + c_2$$

2.1.35 problem 35

Solved as second order linear constant coeff ode	313
Solved as second order linear exact ode	315
Solved as second order missing y ode	317
Solved as second order integrable as is ode	319
Solved as second order integrable as is ode (ABC method)	320
Solved as second order ode using Kovacic algorithm	321
Solved as second order ode adjoint method	324
Maple step by step solution	327
Maple trace	328
Maple dsolve solution	328
Mathematica DSolve solution	328

Internal problem ID [8782]

Book : Second order enumerated odes

Section : section 1

Problem number : 35

Date solved : Thursday, December 12, 2024 at 09:47:23 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y'' + y' = x^3 + x^2 + x + 1$$

Solved as second order linear constant coeff ode

Time used: 0.138 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = x^3 + x^2 + x + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{1}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{1}{2}$$

Which simplifies to

$$\lambda_1 = 0$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(0)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^3 + x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3, x^4\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_4 x^4 + A_3 x^3 + A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4x^3 A_4 + 3x^2 A_3 + 12x^2 A_4 + 2xA_2 + 6xA_3 + A_1 + 2A_2 = x^3 + x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -4, A_2 = \frac{5}{2}, A_3 = -\frac{2}{3}, A_4 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{4}x^4 - \frac{2}{3}x^3 + \frac{5}{2}x^2 - 4x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + \left(\frac{1}{4}x^4 - \frac{2}{3}x^3 + \frac{5}{2}x^2 - 4x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} - 4x + c_1 + c_2 e^{-x}$$




Figure 2.109: Slope field plot
 $y'' + y' = x^3 + x^2 + x + 1$

Solved as second order linear exact ode

Time used: 0.130 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= x^3 + x^2 + x + 1 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int x^3 + x^2 + x + 1 dx$$

We now have a first order ode to solve which is

$$y' + y = \frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= \frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x dx \\ &= \frac{(3x^4 - 8x^3 + 30x^2 + 12c_1 - 48x + 48) e^x}{12} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} + c_1 - 4x + 4 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} + c_1 - 4x + 4 + c_2 e^{-x}$$




Figure 2.110: Slope field plot
 $y'' + y' = x^3 + x^2 + x + 1$

Solved as second order missing y ode

Time used: 0.112 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - x^3 - x^2 - x - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = x^3 + x^2 + x + 1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu p) = \mu p$$

$$\frac{d}{dx}(\mu p) = (\mu) (x^3 + x^2 + x + 1)$$

$$\frac{d}{dx}(p e^x) = (e^x) (x^3 + x^2 + x + 1)$$

$$d(p e^x) = ((x^3 + x^2 + x + 1) e^x) dx$$

Integrating gives

$$\begin{aligned} p e^x &= \int (x^3 + x^2 + x + 1) e^x dx \\ &= (x^3 - 2x^2 + 5x - 4) e^x + c_1 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$p(x) = x^3 - 2x^2 + 5x - 4 + c_1 e^{-x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = x^3 - 2x^2 + 5x - 4 + c_1 e^{-x}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned} \int dy &= \int x^3 - 2x^2 + 5x - 4 + c_1 e^{-x} dx \\ y &= -4x + \frac{x^4}{4} - c_1 e^{-x} + \frac{5x^2}{2} - \frac{2x^3}{3} + c_2 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -4x + \frac{x^4}{4} - c_1 e^{-x} + \frac{5x^2}{2} - \frac{2x^3}{3} + c_2$$




Figure 2.111: Slope field plot

$$y'' + y' = x^3 + x^2 + x + 1$$

Solved as second order integrable as is ode

Time used: 0.125 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (x^3 + x^2 + x + 1) dx$$

$$y' + y = \frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) \\ d(y e^x) &= \left(\left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x dx \\ &= \frac{(3x^4 - 8x^3 + 30x^2 + 12c_1 - 48x + 48) e^x}{12} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} + c_1 - 4x + 4 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} + c_1 - 4x + 4 + c_2 e^{-x}$$




Figure 2.112: Slope field plot

$$y'' + y' = x^3 + x^2 + x + 1$$

Solved as second order integrable as is ode (ABC method)

Time used: 0.061 (sec)

Writing the ode as

$$y'' + y' = x^3 + x^2 + x + 1$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (x^3 + x^2 + x + 1) dx$$

$$y' + y = \frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right)$$

$$\frac{d}{dx}(y e^x) = (e^x) \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right)$$

$$d(y e^x) = \left(\left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x \right) dx$$

Integrating gives

$$\begin{aligned} y e^x &= \int \left(\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + x + c_1 \right) e^x dx \\ &= \frac{(3x^4 - 8x^3 + 30x^2 + 12c_1 - 48x + 48) e^x}{12} + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} + c_1 - 4x + 4 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.113: Slope field plot
 $y'' + y' = x^3 + x^2 + x + 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.078 (sec)

Writing the ode as

$$y'' + y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = y e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = r z(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 1 \\ t &= 4\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.46: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^3 + x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3, x^4\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_4 x^4 + A_3 x^3 + A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4x^3A_4 + 3x^2A_3 + 12x^2A_4 + 2xA_2 + 6xA_3 + A_1 + 2A_2 = x^3 + x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -4, A_2 = \frac{5}{2}, A_3 = -\frac{2}{3}, A_4 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{4}x^4 - \frac{2}{3}x^3 + \frac{5}{2}x^2 - 4x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + \left(\frac{1}{4}x^4 - \frac{2}{3}x^3 + \frac{5}{2}x^2 - 4x \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 + \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} - 4x$$




Figure 2.114: Slope field plot
 $y'' + y' = x^3 + x^2 + x + 1$

Solved as second order ode adjoint method

Time used: 0.579 (sec)

In normal form the ode

$$y'' + y' = x^3 + x^2 + x + 1 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= x^3 + x^2 + x + 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = c_1 e^x + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(1 - \frac{c_1 e^x}{c_1 e^x + c_2} \right) = \frac{c_2 x + c_1 e^x + \frac{c_2 x^2}{2} + \frac{c_2 x^3}{3} + \frac{c_2 x^4}{4} + c_1 (e^x x - e^x) + c_1 (e^x x^2 - 2 e^x x + 2 e^x) + c_1 (e^x x^3 - 3 e^x x^2 + 6 e^x x - 6 e^x)}{c_1 e^x + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= \frac{c_2}{c_1 e^x + c_2} \\ p(x) &= \frac{12c_1(x-1)(x^2-x+4)e^x + 3(x^3 + \frac{4}{3}x^2 + 2x + 4)xc_2}{12c_1 e^x + 12c_2}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{c_1 e^x + c_2} dx} \\ &= \frac{e^x}{c_1 e^x + c_2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{12c_1(x-1)(x^2-x+4)e^x + 3(x^3 + \frac{4}{3}x^2 + 2x + 4)xc_2}{12c_1 e^x + 12c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{e^x}{c_1 e^x + c_2} \right) \left(\frac{12c_1(x-1)(x^2-x+4)e^x + 3(x^3 + \frac{4}{3}x^2 + 2x + 4)xc_2}{12c_1 e^x + 12c_2} \right) \\ d \left(\frac{y e^x}{c_1 e^x + c_2} \right) &= \left(\frac{(12c_1(x-1)(x^2-x+4)e^x + 3(x^3 + \frac{4}{3}x^2 + 2x + 4)xc_2) e^x}{(12c_1 e^x + 12c_2)(c_1 e^x + c_2)} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y e^x}{c_1 e^x + c_2} &= \int \frac{(12c_1(x-1)(x^2-x+4)e^x + 3(x^3 + \frac{4}{3}x^2 + 2x + 4)xc_2) e^x}{(12c_1 e^x + 12c_2)(c_1 e^x + c_2)} dx \\ &= \frac{4e^x + \frac{x^4 e^x}{4} - 4e^x x + \frac{5e^x x^2}{2} - \frac{2e^x x^3}{3}}{c_1 e^x + c_2} + c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{c_1 e^x + c_2}$ gives the final solution

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + c_1 c_3 + \frac{5x^2}{2} - 4x + c_2 c_3 e^{-x} + 4$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + c_1 c_3 + \frac{5x^2}{2} - 4x + c_2 c_3 e^{-x} + 4$$

The constants can be merged to give

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} + c_1 - 4x + 4 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} + c_1 - 4x + 4 + c_2 e^{-x}$$




Figure 2.115: Slope field plot
 $y'' + y' = x^3 + x^2 + x + 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = x^3 + x^2 + x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right) \right], f(x) = x^3 + x^2 + x + 1$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x(x+1)(x^2+1) dx \right) + \int (x^3+x^2+x+1) dx$$

- Compute integrals

$$y_p(x) = -\frac{2}{3}x^3 + \frac{5}{2}x^2 - 4x + 4 + \frac{1}{4}x^4$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-x} + C_2 - \frac{2x^3}{3} + \frac{5x^2}{2} - 4x + 4 + \frac{x^4}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = _a^3+_a^2-_b(_a)+_a+1, _b(_a)`
Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 31

```

dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = x^3+x^2+x+1,
y(x),singsol=all)

```

$$y = \frac{x^4}{4} - c_1 e^{-x} + \frac{5x^2}{2} - \frac{2x^3}{3} - 4x + c_2$$

Mathematica DSolve solution

Solving time : 0.132 (sec)

Leaf size : 41

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]==1+x+x^2+x^3,{x]},
y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{x^4}{4} - \frac{2x^3}{3} + \frac{5x^2}{2} - 4x - c_1 e^{-x} + c_2$$

2.1.36 problem 36

Solved as second order linear constant coeff ode	329
Solved as second order linear exact ode	331
Solved as second order missing y ode	333
Solved as second order integrable as is ode	334
Solved as second order integrable as is ode (ABC method)	336
Solved as second order ode using Kovacic algorithm	337
Solved as second order ode adjoint method	340
Maple step by step solution	343
Maple trace	344
Maple dsolve solution	344
Mathematica DSolve solution	344

Internal problem ID [8783]

Book : Second order enumerated odes

Section : section 1

Problem number : 36

Date solved : Thursday, December 12, 2024 at 09:47:25 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y'' + y' = \sin(x)$$

Solved as second order linear constant coeff ode

Time used: 0.146 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \cos(x) - A_2 \sin(x) - A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = -\frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + \left(-\frac{\cos(x)}{2} - \frac{\sin(x)}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$




Figure 2.116: Slope field plot
 $y'' + y' = \sin(x)$

Solved as second order linear exact ode

Time used: 0.157 (sec)

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= \sin(x) \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int \sin(x) dx$$

We now have a first order ode to solve which is

$$y' + y = -\cos(x) + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= -\cos(x) + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(-\cos(x) + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(-\cos(x) + c_1) \\ d(y e^x) &= ((-\cos(x) + c_1) e^x) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int (-\cos(x) + c_1) e^x dx \\ &= -\frac{e^x \cos(x)}{2} - \frac{\sin(x) e^x}{2} + e^x c_1 + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$




Figure 2.117: Slope field plot
 $y'' + y' = \sin(x)$

Solved as second order missing y ode

Time used: 0.196 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - \sin(x) = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= \sin(x) \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu p) &= \mu p \\ \frac{d}{dx}(\mu p) &= (\mu) (\sin(x)) \\ \frac{d}{dx}(p e^x) &= (e^x) (\sin(x)) \\ d(p e^x) &= (\sin(x) e^x) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} p e^x &= \int \sin(x) e^x dx \\ &= -\frac{e^x \cos(x)}{2} + \frac{\sin(x) e^x}{2} + c_1 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$p(x) = \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_1 e^{-x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_1 e^{-x}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned} \int dy &= \int \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_1 e^{-x} dx \\ y &= -c_1 e^{-x} - \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_2 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -c_1 e^{-x} - \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_2$$




Figure 2.118: Slope field plot
 $y'' + y' = \sin(x)$

Solved as second order integrable as is ode

Time used: 0.056 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned} \int (y'' + y') dx &= \int \sin(x) dx \\ y' + y &= -\cos(x) + c_1 \end{aligned}$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= -\cos(x) + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(-\cos(x) + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(-\cos(x) + c_1) \\ d(y e^x) &= ((-\cos(x) + c_1) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int (-\cos(x) + c_1) e^x dx \\ &= -\frac{e^x \cos(x)}{2} - \frac{\sin(x) e^x}{2} + e^x c_1 + c_2\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$




Figure 2.119: Slope field plot
 $y'' + y' = \sin(x)$

Solved as second order integrable as is ode (ABC method)

Time used: 0.055 (sec)

Writing the ode as

$$y'' + y' = \sin(x)$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int \sin(x) dx$$

$$y' + y = -\cos(x) + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = -\cos(x) + c_1$$

The integrating factor μ is

$$\mu = e^{\int q dx}$$

$$= e^{\int 1 dx}$$

$$= e^x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu)(-\cos(x) + c_1)$$

$$\frac{d}{dx}(y e^x) = (e^x)(-\cos(x) + c_1)$$

$$d(y e^x) = ((-\cos(x) + c_1) e^x) dx$$

Integrating gives

$$y e^x = \int (-\cos(x) + c_1) e^x dx$$

$$= -\frac{e^x \cos(x)}{2} - \frac{\sin(x) e^x}{2} + e^x c_1 + c_2$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.120: Slope field plot
 $y'' + y' = \sin(x)$

Solved as second order ode using Kovacic algorithm

Time used: 0.138 (sec)

Writing the ode as

$$y'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.48: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{y_1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \cos(x) - A_2 \sin(x) - A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = -\frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{2} - \frac{\sin(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + \left(-\frac{\cos(x)}{2} - \frac{\sin(x)}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 - \frac{\cos(x)}{2} - \frac{\sin(x)}{2}$$




Figure 2.121: Slope field plot
 $y'' + y' = \sin(x)$

Solved as second order ode adjoint method

Time used: 0.603 (sec)

In normal form the ode

$$y'' + y' = \sin(x) \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= \sin(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = e^x c_1 + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{e^x c_1}{e^x c_1 + c_2} \right) = \frac{c_1 \left(-\frac{e^x \cos(x)}{2} + \frac{\sin(x) e^x}{2} \right) - \cos(x) c_2}{e^x c_1 + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = \frac{c_2}{e^x c_1 + c_2}$$

$$p(x) = \frac{-c_1(-\sin(x) + \cos(x)) e^x - 2 \cos(x) c_2}{2 e^x c_1 + 2 c_2}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{e^x c_1 + c_2} dx} \\ &= \frac{e^x}{e^x c_1 + c_2} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{-c_1(-\sin(x) + \cos(x)) e^x - 2 \cos(x) c_2}{2 e^x c_1 + 2 c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{e^x c_1 + c_2} \right) &= \left(\frac{e^x}{e^x c_1 + c_2} \right) \left(\frac{-c_1(-\sin(x) + \cos(x)) e^x - 2 \cos(x) c_2}{2 e^x c_1 + 2 c_2} \right) \\ d \left(\frac{y e^x}{e^x c_1 + c_2} \right) &= \left(\frac{(-c_1(-\sin(x) + \cos(x)) e^x - 2 \cos(x) c_2) e^x}{(2 e^x c_1 + 2 c_2) (e^x c_1 + c_2)} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y e^x}{e^x c_1 + c_2} &= \int \frac{(-c_1(-\sin(x) + \cos(x)) e^x - 2 \cos(x) c_2) e^x}{(2 e^x c_1 + 2 c_2) (e^x c_1 + c_2)} dx \\ &= \frac{-e^x - e^x \tan\left(\frac{x}{2}\right) - \frac{c_2}{2c_1} - \frac{c_2 \tan\left(\frac{x}{2}\right)^2}{2c_1}}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right) (e^x c_1 + c_2)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{e^x c_1 + c_2}$ gives the final solution

$$y = \frac{c_2(2c_3c_1 - 1) e^{-x} + 2 \left(c_3c_1 - \frac{\cos(x)}{2} - \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{c_2(2c_3c_1 - 1) e^{-x} + 2 \left(c_3c_1 - \frac{\cos(x)}{2} - \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$

The constants can be merged to give

$$y = \frac{c_2(2c_1 - 1) e^{-x} + 2 \left(c_1 - \frac{\cos(x)}{2} - \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_2(2c_1 - 1) e^{-x} + 2 \left(c_1 - \frac{\cos(x)}{2} - \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$




Figure 2.122: Slope field plot
 $y'' + y' = \sin(x)$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = \sin(x)$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x \sin(x) dx \right) + \int \sin(x) dx$$

- Compute integrals

$$y_p(x) = -\frac{\sin(x)}{2} - \frac{\cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-x} + C_2 - \frac{\sin(x)}{2} - \frac{\cos(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)+sin(_a), _b(_a)` *** Subl
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 21

```

dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = sin(x),
        y(x),singsol=all)

```

$$y = -c_1 e^{-x} - \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_2$$

Mathematica DSolve solution

Solving time : 0.105 (sec)

Leaf size : 29

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]==Sin[x],{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow -\frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_1(-e^{-x}) + c_2$$

2.1.37 problem 37

Solved as second order linear constant coeff ode	345
Solved as second order linear exact ode	347
Solved as second order missing y ode	349
Solved as second order integrable as is ode	350
Solved as second order integrable as is ode (ABC method)	352
Solved as second order ode using Kovacic algorithm	353
Solved as second order ode adjoint method	356
Maple step by step solution	359
Maple trace	360
Maple dsolve solution	360
Mathematica DSolve solution	360

Internal problem ID [8784]

Book : Second order enumerated odes

Section : section 1

Problem number : 37

Date solved : Thursday, December 12, 2024 at 09:47:28 AM

CAS classification : [[_2nd_order, _missing_y]]

Solve

$$y'' + y' = \cos(x)$$

Solved as second order linear constant coeff ode

Time used: 0.139 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + \lambda e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{1}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(0)x} + c_2 e^{(-1)x}\end{aligned}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \cos(x) - A_2 \sin(x) - A_1 \sin(x) + A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + \left(-\frac{\cos(x)}{2} + \frac{\sin(x)}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$




Figure 2.123: Slope field plot
 $y'' + y' = \cos(x)$

Solved as second order linear exact ode

Time used: 0.142 (sec)

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= \cos(x) \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' + y = \int \cos(x) dx$$

We now have a first order ode to solve which is

$$y' + y = \sin(x) + c_1$$

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= 1 \\ p(x) &= \sin(x) + c_1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(\sin(x) + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(\sin(x) + c_1) \\ d(y e^x) &= ((\sin(x) + c_1) e^x) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^x &= \int (\sin(x) + c_1) e^x dx \\ &= -\frac{\cos(x) e^x}{2} + \frac{e^x \sin(x)}{2} + e^x c_1 + c_2 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$




Figure 2.124: Slope field plot
 $y'' + y' = \cos(x)$

Solved as second order missing y ode

Time used: 0.205 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x) - \cos(x) = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = r(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$r(x) = \cos(x)$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x \end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu p) = \mu r$$

$$\frac{d}{dx}(\mu p) = (\mu) (\cos(x))$$

$$\frac{d}{dx}(p e^x) = (e^x) (\cos(x))$$

$$d(p e^x) = (\cos(x) e^x) dx$$

Integrating gives

$$\begin{aligned} p e^x &= \int \cos(x) e^x dx \\ &= \frac{\cos(x) e^x}{2} + \frac{e^x \sin(x)}{2} + c_1 \end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$p(x) = \frac{\sin(x)}{2} + \frac{\cos(x)}{2} + c_1 e^{-x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{\sin(x)}{2} + \frac{\cos(x)}{2} + c_1 e^{-x}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{\sin(x)}{2} + \frac{\cos(x)}{2} + c_1 e^{-x} dx$$

$$y = -c_1 e^{-x} + \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -c_1 e^{-x} + \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_2$$




Figure 2.125: Slope field plot
 $y'' + y' = \cos(x)$

Solved as second order integrable as is ode

Time used: 0.055 (sec)

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int \cos(x) dx$$

$$y' + y = \sin(x) + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \sin(x) + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) (\sin(x) + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x) (\sin(x) + c_1) \\ d(y e^x) &= ((\sin(x) + c_1) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int (\sin(x) + c_1) e^x dx \\ &= -\frac{\cos(x) e^x}{2} + \frac{e^x \sin(x)}{2} + e^x c_1 + c_2\end{aligned}$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$




Figure 2.126: Slope field plot
 $y'' + y' = \cos(x)$

Solved as second order integrable as is ode (ABC method)

Time used: 0.056 (sec)

Writing the ode as

$$y'' + y' = \cos(x)$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int \cos(x) dx$$

$$y' + y = \sin(x) + c_1$$

Which is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = 1$$

$$p(x) = \sin(x) + c_1$$

The integrating factor μ is

$$\mu = e^{\int q dx}$$

$$= e^{\int 1 dx}$$

$$= e^x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu)(\sin(x) + c_1)$$

$$\frac{d}{dx}(y e^x) = (e^x)(\sin(x) + c_1)$$

$$d(y e^x) = ((\sin(x) + c_1) e^x) dx$$

Integrating gives

$$y e^x = \int (\sin(x) + c_1) e^x dx$$

$$= -\frac{\cos(x) e^x}{2} + \frac{e^x \sin(x)}{2} + e^x c_1 + c_2$$

Dividing throughout by the integrating factor e^x gives the final solution

$$y = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2} + c_1 + c_2 e^{-x}$$

Will add steps showing solving for IC soon.




Figure 2.127: Slope field plot
 $y'' + y' = \cos(x)$

Solved as second order ode using Kovacic algorithm

Time used: 0.077 (sec)

Writing the ode as

$$y'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.50: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \cos(x) - A_2 \sin(x) - A_1 \sin(x) + A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + \left(-\frac{\cos(x)}{2} + \frac{\sin(x)}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-x} + c_2 - \frac{\cos(x)}{2} + \frac{\sin(x)}{2}$$




Figure 2.128: Slope field plot
 $y'' + y' = \cos(x)$

Solved as second order ode adjoint method

Time used: 0.655 (sec)

In normal form the ode

$$y'' + y' = \cos(x) \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= \cos(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (\xi(x))' + (0) &= 0 \\ \xi''(x) - \xi'(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = -1, C = 0$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - \lambda e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(0)} \\ &= \frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= 0 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} \xi &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ \xi &= c_1 e^{(1)x} + c_2 e^{(0)x} \end{aligned}$$

Or

$$\xi = e^x c_1 + c_2$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(1 - \frac{e^x c_1}{e^x c_1 + c_2} \right) = \frac{c_1 \left(\frac{\cos(x)e^x}{2} + \frac{e^x \sin(x)}{2} \right) + \sin(x) c_2}{e^x c_1 + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = \frac{c_2}{e^x c_1 + c_2}$$

$$p(x) = \frac{(\cos(x) + \sin(x)) c_1 e^x + 2 \sin(x) c_2}{2 e^x c_1 + 2 c_2}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int \frac{c_2}{e^x c_1 + c_2} dx} \\ &= \frac{e^x}{e^x c_1 + c_2} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{(\cos(x) + \sin(x)) c_1 e^x + 2 \sin(x) c_2}{2 e^x c_1 + 2 c_2} \right) \\ \frac{d}{dx} \left(\frac{y e^x}{e^x c_1 + c_2} \right) &= \left(\frac{e^x}{e^x c_1 + c_2} \right) \left(\frac{(\cos(x) + \sin(x)) c_1 e^x + 2 \sin(x) c_2}{2 e^x c_1 + 2 c_2} \right) \\ d \left(\frac{y e^x}{e^x c_1 + c_2} \right) &= \left(\frac{((\cos(x) + \sin(x)) c_1 e^x + 2 \sin(x) c_2) e^x}{(2 e^x c_1 + 2 c_2) (e^x c_1 + c_2)} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y e^x}{e^x c_1 + c_2} &= \int \frac{((\cos(x) + \sin(x)) c_1 e^x + 2 \sin(x) c_2) e^x}{(2 e^x c_1 + 2 c_2) (e^x c_1 + c_2)} dx \\ &= \frac{-e^x + e^x \tan\left(\frac{x}{2}\right) - \frac{c_2}{2c_1} - \frac{c_2 \tan\left(\frac{x}{2}\right)^2}{2c_1}}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right) (e^x c_1 + c_2)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{e^x}{e^x c_1 + c_2}$ gives the final solution

$$y = \frac{c_2(2c_3c_1 - 1) e^{-x} + 2 \left(c_3c_1 - \frac{\cos(x)}{2} + \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{c_2(2c_3c_1 - 1) e^{-x} + 2 \left(c_3c_1 - \frac{\cos(x)}{2} + \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$

The constants can be merged to give

$$y = \frac{c_2(2c_1 - 1) e^{-x} + 2 \left(c_1 - \frac{\cos(x)}{2} + \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_2(2c_1 - 1) e^{-x} + 2 \left(c_1 - \frac{\cos(x)}{2} + \frac{\sin(x)}{2} - \frac{1}{2} \right) c_1}{2c_1}$$




Figure 2.129: Slope field plot
 $y'' + y' = \cos(x)$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + \frac{d}{dx}y(x) = \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 e^{-x} + C2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^x \cos(x) dx \right) + \int \cos(x) dx$$

- Compute integrals

$$y_p(x) = -\frac{\cos(x)}{2} + \frac{\sin(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 e^{-x} + C_2 - \frac{\cos(x)}{2} + \frac{\sin(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)+cos(_a), _b(_a)` *** Subl
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 21

```

dsolve(diff(y(x),x)+diff(diff(y(x),x),x) = cos(x),
        y(x),singsol=all)

```

$$y = -c_1 e^{-x} + \frac{\sin(x)}{2} - \frac{\cos(x)}{2} + c_2$$

Mathematica DSolve solution

Solving time : 0.078 (sec)

Leaf size : 28

```

DSolve[{D[y[x],{x,2}]+D[y[x],x]==Cos[x],{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{1}{2}(\sin(x) - \cos(x) - 2c_1 e^{-x}) + c_2$$

2.1.38 problem 38

Solved as second order linear constant coeff ode	361
Solved as second order can be made integrable	363
Solved as second order ode using Kovacic algorithm	365
Solved as second order ode adjoint method	368
Maple step by step solution	371
Maple trace	372
Maple dsolve solution	372
Mathematica DSolve solution	372

Internal problem ID [8785]

Book : Second order enumerated odes

Section : section 1

Problem number : 38

Date solved : Thursday, December 12, 2024 at 09:47:30 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$y'' + y = 1$$

Solved as second order linear constant coeff ode

Time used: 0.115 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.130: Slope field plot
 $y'' + y = 1$

Solved as second order can be made integrable

Time used: 0.844 (sec)

Multiplying the ode by y' gives

$$y'y'' + y'y - y' = 0$$

Integrating the above w.r.t x gives

$$\begin{aligned} \int (y'y'' + y'y - y') dx &= 0 \\ \frac{y'^2}{2} + \frac{y^2}{2} - y &= c_1 \end{aligned}$$

Which is now solved for y . Solving for the derivative gives these ODE's to solve

$$y' = \sqrt{-y^2 + 2y + 2c_1} \tag{1}$$

$$y' = -\sqrt{-y^2 + 2y + 2c_1} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Integrating gives

$$\begin{aligned} \int \frac{1}{\sqrt{-y^2 + 2c_1 + 2y}} dy &= dx \\ \arctan\left(\frac{y-1}{\sqrt{-y^2 + 2c_1 + 2y}}\right) &= x + c_2 \end{aligned}$$

Singular solutions are found by solving

$$\sqrt{-y^2 + 2c_1 + 2y} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 1 - \sqrt{1 + 2c_1}$$

$$y = 1 + \sqrt{1 + 2c_1}$$

Solving for y gives

$$y = \tan(x + c_2) \sqrt{\frac{1 + 2c_1}{\tan(x + c_2)^2 + 1}} + 1$$

Solving Eq. (2)

Integrating gives

$$\int -\frac{1}{\sqrt{-y^2 + 2c_1 + 2y}} dy = dx$$

$$-\arctan\left(\frac{y-1}{\sqrt{-y^2 + 2c_1 + 2y}}\right) = x + c_3$$

Singular solutions are found by solving

$$-\sqrt{-y^2 + 2c_1 + 2y} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 1 - \sqrt{1 + 2c_1}$$

$$y = 1 + \sqrt{1 + 2c_1}$$

Solving for y gives

$$y = -\tan(x + c_3) \sqrt{\frac{1 + 2c_1}{\tan(x + c_3)^2 + 1}} + 1$$

Will add steps showing solving for IC soon.

The solution

$$y = 1 - \sqrt{1 + 2c_1}$$

was found not to satisfy the ode or the IC. Hence it is removed. The solution

$$y = 1 + \sqrt{1 + 2c_1}$$

was found not to satisfy the ode or the IC. Hence it is removed.

Summary of solutions found

$$y = \tan(x + c_2) \sqrt{\frac{1 + 2c_1}{\tan(x + c_2)^2 + 1}} + 1$$

$$y = -\tan(x + c_3) \sqrt{\frac{1 + 2c_1}{\tan(x + c_3)^2 + 1}} + 1$$




Figure 2.131: Slope field plot
 $y'' + y = 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.109 (sec)

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.52: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.132: Slope field plot
 $y'' + y = 1$

Solved as second order ode adjoint method

Time used: 0.766 (sec)

In normal form the ode

$$y'' + y = 1 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$p(x) = 0$$

$$q(x) = 1$$

$$r(x) = 1$$

The Lagrange adjoint ode is given by

$$\xi'' - (\xi p)' + \xi q = 0$$

$$\xi'' - (0)' + (\xi(x)) = 0$$

$$\xi''(x) + \xi(x) = 0$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$\xi = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' - \frac{y(-c_1 \sin(x) + c_2 \cos(x))}{c_1 \cos(x) + c_2 \sin(x)} = \frac{c_1 \sin(x) - c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)}$$

$$p(x) = \frac{c_1 \sin(x) - c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{c_1 \sin(x) - c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{1}{c_1 \cos(x) + c_2 \sin(x)} \right) \left(\frac{c_1 \sin(x) - c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ d \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{c_1 \sin(x) - c_2 \cos(x)}{(c_1 \cos(x) + c_2 \sin(x))^2} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_1 \cos(x) + c_2 \sin(x)} &= \int \frac{c_1 \sin(x) - c_2 \cos(x)}{(c_1 \cos(x) + c_2 \sin(x))^2} dx \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x) + c_2 \sin(x)}$ gives the final solution

$$y = \cos(x) c_1 c_3 + \sin(x) c_2 c_3 + 1$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \cos(x) c_1 c_3 + \sin(x) c_2 c_3 + 1$$

The constants can be merged to give

$$y = 1 + c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.133: Slope field plot
 $y'' + y = 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + y(x) = 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) dx \right) + \sin(x) \left(\int \cos(x) dx \right)$$

- Compute integrals

$$y_p(x) = 1$$

- Substitute particular solution into general solution to ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.005 (sec)

Leaf size : 14

```

dsolve(diff(diff(y(x),x),x)+y(x) = 1,
        y(x),singsol=all)

```

$$y = \cos(x) c_1 + \sin(x) c_2 + 1$$

Mathematica DSolve solution

Solving time : 0.012 (sec)

Leaf size : 17

```

DSolve[{D[y[x],{x,2}]+y[x]==1,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow c_1 \cos(x) + c_2 \sin(x) + 1$$

2.1.39 problem 39

Solved as second order linear constant coeff ode	373
Solved as second order ode using Kovacic algorithm	375
Solved as second order ode adjoint method	378
Maple step by step solution	381
Maple trace	382
Maple dsolve solution	382
Mathematica DSolve solution	382

Internal problem ID [8786]

Book : Second order enumerated odes

Section : section 1

Problem number : 39

Date solved : Thursday, December 12, 2024 at 09:47:32 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' + y = x$$

Solved as second order linear constant coeff ode

Time used: 0.126 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2 x + A_1 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.134: Slope field plot
 $y'' + y = x$

Solved as second order ode using Kovacic algorithm

Time used: 0.134 (sec)

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\ t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.54: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2 x + A_1 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.135: Slope field plot
 $y'' + y = x$

Solved as second order ode adjoint method

Time used: 1.348 (sec)

In normal form the ode

$$y'' + y = x \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$p(x) = 0$$

$$q(x) = 1$$

$$r(x) = x$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + (\xi(x)) &= 0 \\ \xi''(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$\xi = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' - \frac{y(-c_1 \sin(x) + c_2 \cos(x))}{c_1 \cos(x) + c_2 \sin(x)} = \frac{c_1(\cos(x) + x \sin(x)) + c_2(\sin(x) - x \cos(x))}{c_1 \cos(x) + c_2 \sin(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)}$$

$$p(x) = \frac{(-c_2x + c_1) \cos(x) + \sin(x)(c_1x + c_2)}{c_1 \cos(x) + c_2 \sin(x)}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{(-c_2x + c_1) \cos(x) + \sin(x)(c_1x + c_2)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{1}{c_1 \cos(x) + c_2 \sin(x)} \right) \left(\frac{(-c_2x + c_1) \cos(x) + \sin(x)(c_1x + c_2)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ d \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{(-c_2x + c_1) \cos(x) + \sin(x)(c_1x + c_2)}{(c_1 \cos(x) + c_2 \sin(x))^2} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_1 \cos(x) + c_2 \sin(x)} &= \int \frac{(-c_2x + c_1) \cos(x) + \sin(x)(c_1x + c_2)}{(c_1 \cos(x) + c_2 \sin(x))^2} dx \\ &= \frac{-x - 2x \tan\left(\frac{x}{2}\right)^2 - x \tan\left(\frac{x}{2}\right)^4}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right) \left(\tan\left(\frac{x}{2}\right)^2 c_1 - 2 \tan\left(\frac{x}{2}\right) c_2 - c_1\right)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x) + c_2 \sin(x)}$ gives the final solution

$$y = c_2 c_3 \sin(x) + c_1 c_3 \cos(x) + x$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_2 c_3 \sin(x) + c_1 c_3 \cos(x) + x$$

The constants can be merged to give

$$y = x + c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.136: Slope field plot
 $y'' + y = x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + y(x) = x$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) x dx \right) + \sin(x) \left(\int x \cos(x) dx \right)$$

- Compute integrals

$$y_p(x) = x$$

- Substitute particular solution into general solution to ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 14

```

dsolve(diff(diff(y(x),x),x)+y(x) = x,
        y(x),singsol=all)

```

$$y = \sin(x) c_2 + \cos(x) c_1 + x$$

Mathematica DSolve solution

Solving time : 0.013 (sec)

Leaf size : 17

```

DSolve[{D[y[x],{x,2}]+y[x]==x,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow x + c_1 \cos(x) + c_2 \sin(x)$$

2.1.40 problem 40

Solved as second order linear constant coeff ode	383
Solved as second order ode using Kovacic algorithm	385
Solved as second order ode adjoint method	388
Maple step by step solution	391
Maple trace	392
Maple dsolve solution	392
Mathematica DSolve solution	392

Internal problem ID [8787]

Book : Second order enumerated odes

Section : section 1

Problem number : 40

Date solved : Thursday, December 12, 2024 at 09:47:34 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' + y = 1 + x$$

Solved as second order linear constant coeff ode

Time used: 0.126 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = 1 + x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2 x + A_1 = 1 + x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 1 + x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (1 + x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + x + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.137: Slope field plot
 $y'' + y = 1 + x$

Solved as second order ode using Kovacic algorithm

Time used: 0.138 (sec)

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\ t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.56: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2 x + A_1 = 1 + x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 1 + x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (1 + x) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + x + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.138: Slope field plot
 $y'' + y = 1 + x$

Solved as second order ode adjoint method

Time used: 1.385 (sec)

In normal form the ode

$$y'' + y = 1 + x \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 0 \\ q(x) &= 1 \\ r(x) &= 1 + x \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + (\xi(x)) &= 0 \\ \xi''(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$\xi = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' - \frac{y(-c_1 \sin(x) + c_2 \cos(x))}{c_1 \cos(x) + c_2 \sin(x)} = \frac{c_1 \sin(x) + c_1(\cos(x) + x \sin(x)) - c_2 \cos(x) + c_2(\sin(x) - x \cos(x))}{c_1 \cos(x) + c_2 \sin(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)}$$

$$p(x) = \frac{(c_1 + (-x - 1)c_2) \cos(x) + \sin(x) ((1 + x)c_1 + c_2)}{c_1 \cos(x) + c_2 \sin(x)}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{(c_1 + (-x - 1)c_2) \cos(x) + \sin(x) ((1 + x)c_1 + c_2)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{1}{c_1 \cos(x) + c_2 \sin(x)} \right) \left(\frac{(c_1 + (-x - 1)c_2) \cos(x) + \sin(x) ((1 + x)c_1 + c_2)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ d \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{(c_1 + (-x - 1)c_2) \cos(x) + \sin(x) ((1 + x)c_1 + c_2)}{(c_1 \cos(x) + c_2 \sin(x))^2} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_1 \cos(x) + c_2 \sin(x)} &= \int \frac{(c_1 + (-x - 1)c_2) \cos(x) + \sin(x) ((1 + x)c_1 + c_2)}{(c_1 \cos(x) + c_2 \sin(x))^2} dx \\ &= \frac{-x - 2 \tan\left(\frac{x}{2}\right)^2 - 2x \tan\left(\frac{x}{2}\right)^2 - x \tan\left(\frac{x}{2}\right)^4 - 1 - \tan\left(\frac{x}{2}\right)^4}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right) \left(\tan\left(\frac{x}{2}\right)^2 c_1 - 2 \tan\left(\frac{x}{2}\right) c_2 - c_1\right)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x) + c_2 \sin(x)}$ gives the final solution

$$y = c_2 c_3 \sin(x) + c_1 c_3 \cos(x) + x + 1$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_2 c_3 \sin(x) + c_1 c_3 \cos(x) + x + 1$$

The constants can be merged to give

$$y = 1 + x + c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + x + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.139: Slope field plot
 $y'' + y = 1 + x$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + y(x) = x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-i, i)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y(x) = C_1 y_1(x) + C_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C_1 \cos(x) + C_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = x + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int (x+1) \sin(x) dx \right) + \sin(x) \left(\int (x+1) \cos(x) dx \right)$$

- Compute integrals

$$y_p(x) = x + 1$$

- Substitute particular solution into general solution to ODE

$$y(x) = C_1 \cos(x) + C_2 \sin(x) + x + 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 15

```

dsolve(diff(diff(y(x),x),x)+y(x) = x+1,
        y(x),singsol=all)

```

$$y = \sin(x) c_2 + \cos(x) c_1 + x + 1$$

Mathematica DSolve solution

Solving time : 0.012 (sec)

Leaf size : 18

```

DSolve[{D[y[x],{x,2}]+y[x]==1+x,{]},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow x + c_1 \cos(x) + c_2 \sin(x) + 1$$

2.1.41 problem 41

Solved as second order linear constant coeff ode	393
Solved as second order ode using Kovacic algorithm	395
Solved as second order ode adjoint method	398
Maple step by step solution	401
Maple trace	402
Maple dsolve solution	402
Mathematica DSolve solution	402

Internal problem ID [8788]

Book : Second order enumerated odes

Section : section 1

Problem number : 41

Date solved : Thursday, December 12, 2024 at 09:47:37 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' + y = x^2 + x + 1$$

Solved as second order linear constant coeff ode

Time used: 0.128 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = x^2 + x + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3x^2 + A_2x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_3x^2 + A_2x + A_1 + 2A_3 = x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 1, A_3 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^2 + x - 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (x^2 + x - 1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^2 + x - 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.140: Slope field plot
 $y'' + y = x^2 + x + 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.138 (sec)

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.58: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_3x^2 + A_2x + A_1 + 2A_3 = x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 1, A_3 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^2 + x - 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (x^2 + x - 1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^2 + x - 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.141: Slope field plot
 $y'' + y = x^2 + x + 1$

Solved as second order ode adjoint method

Time used: 1.495 (sec)

In normal form the ode

$$y'' + y = x^2 + x + 1 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= 0 \\ q(x) &= 1 \\ r(x) &= x^2 + x + 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned}\xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + (\xi(x)) &= 0 \\ \xi''(x) + \xi(x) &= 0\end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$\xi = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' - \frac{y(-c_1 \sin(x) + c_2 \cos(x))}{c_1 \cos(x) + c_2 \sin(x)} = \frac{c_1(x^2 \sin(x) - 2 \sin(x) + 2x \cos(x)) + c_1(\cos(x) + x \sin(x)) + c_1 \sin(x)}{c_1 \cos(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} \\ p(x) &= \frac{((2x+1)c_1 - c_2(x^2 + x - 1)) \cos(x) + \sin(x)((x^2 + x - 1)c_1 + c_2(2x + 1))}{c_1 \cos(x) + c_2 \sin(x)}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)}\end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{((2x+1)c_1 - c_2(x^2 + x - 1)) \cos(x) + \sin(x)((x^2 + x - 1)c_1 + c_2(2x + 1))}{c_1 \cos(x) + c_2 \sin(x)} \right)\end{aligned}$$

$$\begin{aligned}\frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{1}{c_1 \cos(x) + c_2 \sin(x)} \right) \left(\frac{((2x+1)c_1 - c_2(x^2 + x - 1)) \cos(x) + \sin(x)((x^2 + x - 1)c_1 + c_2(2x + 1))}{c_1 \cos(x) + c_2 \sin(x)} \right)\end{aligned}$$

$$\begin{aligned}d \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{((2x+1)c_1 - c_2(x^2 + x - 1)) \cos(x) + \sin(x)((x^2 + x - 1)c_1 + c_2(2x + 1))}{(c_1 \cos(x) + c_2 \sin(x))^2} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{c_1 \cos(x) + c_2 \sin(x)} &= \int \frac{((2x+1)c_1 - c_2(x^2 + x - 1)) \cos(x) + \sin(x)((x^2 + x - 1)c_1 + c_2(2x + 1))}{(c_1 \cos(x) + c_2 \sin(x))^2} dx \\ &= \frac{-x - x^2 + 2 \tan\left(\frac{x}{2}\right)^2 - 2x \tan\left(\frac{x}{2}\right)^2 - x \tan\left(\frac{x}{2}\right)^4 - 2x^2 \tan\left(\frac{x}{2}\right)^2 - x^2 \tan\left(\frac{x}{2}\right)^4 + 1}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right) \left(\tan\left(\frac{x}{2}\right)^2 c_1 - 2 \tan\left(\frac{x}{2}\right) c_2 - c_1\right)}\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x) + c_2 \sin(x)}$ gives the final solution

$$y = c_1 c_3 \cos(x) + c_2 c_3 \sin(x) + x^2 + x - 1$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_1 c_3 \cos(x) + c_2 c_3 \sin(x) + x^2 + x - 1$$

The constants can be merged to give

$$y = x^2 + x - 1 + c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^2 + x - 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.142: Slope field plot
 $y'' + y = x^2 + x + 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + y(x) = x^2 + x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^2 + x + 1 \right]$$
- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) (x^2 + x + 1) dx \right) + \sin(x) \left(\int \cos(x) (x^2 + x + 1) dx \right)$$
- Compute integrals

$$y_p(x) = x^2 + x - 1$$
- Substitute particular solution into general solution to ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + x^2 + x - 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)
Leaf size : 18

```

dsolve(diff(diff(y(x),x),x)+y(x) = x^2+x+1,
        y(x),singsol=all)

```

$$y = \sin(x) c_2 + \cos(x) c_1 + x^2 + x - 1$$

Mathematica DSolve solution

Solving time : 0.013 (sec)
Leaf size : 21

```

DSolve[{D[y[x],{x,2}]+y[x]==1+x+x^2,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow x^2 + x + c_1 \cos(x) + c_2 \sin(x) - 1$$

2.1.42 problem 42

Solved as second order linear constant coeff ode	403
Solved as second order ode using Kovacic algorithm	405
Solved as second order ode adjoint method	408
Maple step by step solution	411
Maple trace	412
Maple dsolve solution	412
Mathematica DSolve solution	412

Internal problem ID [8789]

Book : Second order enumerated odes

Section : section 1

Problem number : 42

Date solved : Thursday, December 12, 2024 at 09:47:39 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' + y = x^3 + x^2 + x + 1$$

Solved as second order linear constant coeff ode

Time used: 0.135 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = x^3 + x^2 + x + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^3 + x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_4 x^3 + A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_4 x^3 + A_3 x^2 + A_2 x + 6x A_4 + A_1 + 2A_3 = x^3 + x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = -5, A_3 = 1, A_4 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^3 + x^2 - 5x - 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (x^3 + x^2 - 5x - 1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^3 + x^2 - 5x - 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.143: Slope field plot
 $y'' + y = x^3 + x^2 + x + 1$

Solved as second order ode using Kovacic algorithm

Time used: 0.149 (sec)

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.60: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^3 + x^2 + x + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_4 x^3 + A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_4x^3 + A_3x^2 + A_2x + 6xA_4 + A_1 + 2A_3 = x^3 + x^2 + x + 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = -5, A_3 = 1, A_4 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^3 + x^2 - 5x - 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (x^3 + x^2 - 5x - 1) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^3 + x^2 - 5x - 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.144: Slope field plot
 $y'' + y = x^3 + x^2 + x + 1$

Solved as second order ode adjoint method

Time used: 1.494 (sec)

In normal form the ode

$$y'' + y = x^3 + x^2 + x + 1 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 0 \\ q(x) &= 1 \\ r(x) &= x^3 + x^2 + x + 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned}\xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + (\xi(x)) &= 0 \\ \xi''(x) + \xi(x) &= 0\end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$\xi = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y\xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' - \frac{y(-c_1 \sin(x) + c_2 \cos(x))}{c_1 \cos(x) + c_2 \sin(x)} = \frac{c_1(x^3 \sin(x) + 3x^2 \cos(x) - 6 \cos(x) - 6x \sin(x)) + c_1(x^2 \sin(x) - 2 \sin(x))}{c_1 \cos(x) + c_2 \sin(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} \\ p(x) &= \frac{(-c_2 x^3 + (3c_1 - c_2)x^2 + (2c_1 + 5c_2)x - 5c_1 + c_2) \cos(x) + (c_1 x^3 + (c_1 + 3c_2)x^2 + (-5c_1 + 2c_2)x - c_1)}{c_1 \cos(x) + c_2 \sin(x)}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{(-c_2 x^3 + (3c_1 - c_2)x^2 + (2c_1 + 5c_2)x - 5c_1 + c_2) \cos(x) + (c_1 x^3 + (c_1 + 3c_2)x^2 + (-5c_1 + 2c_2)x - c_1)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{1}{c_1 \cos(x) + c_2 \sin(x)} \right) \left(\frac{(-c_2 x^3 + (3c_1 - c_2)x^2 + (2c_1 + 5c_2)x - 5c_1 + c_2) \cos(x) + (c_1 x^3 + (c_1 + 3c_2)x^2 + (-5c_1 + 2c_2)x - c_1)}{c_1 \cos(x) + c_2 \sin(x)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{(-c_2 x^3 + (3c_1 - c_2)x^2 + (2c_1 + 5c_2)x - 5c_1 + c_2) \cos(x) + (c_1 x^3 + (c_1 + 3c_2)x^2 + (-5c_1 + 2c_2)x - c_1)}{(c_1 \cos(x) + c_2 \sin(x))^2} \right)\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{c_1 \cos(x) + c_2 \sin(x)} &= \int \frac{(-c_2 x^3 + (3c_1 - c_2)x^2 + (2c_1 + 5c_2)x - 5c_1 + c_2) \cos(x) + (c_1 x^3 + (c_1 + 3c_2)x^2 + (-5c_1 + 2c_2)x - c_1)}{(c_1 \cos(x) + c_2 \sin(x))^2} dx \\ &= \frac{5x - x^2 - x^3 + 2 \tan\left(\frac{x}{2}\right)^2 + 10x \tan\left(\frac{x}{2}\right)^2 + 5x \tan\left(\frac{x}{2}\right)^4 - 2x^2 \tan\left(\frac{x}{2}\right)^2 - x^2 \tan\left(\frac{x}{2}\right)^2}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right) \left(\tan\left(\frac{x}{2}\right)^2 c_1 - 2 \tan\left(\frac{x}{2}\right) c_2 - c_1\right)}\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x) + c_2 \sin(x)}$ gives the final solution

$$y = c_1 c_3 \cos(x) + c_2 c_3 \sin(x) + x^3 + x^2 - 5x - 1$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_1 c_3 \cos(x) + c_2 c_3 \sin(x) + x^3 + x^2 - 5x - 1$$

The constants can be merged to give

$$y = x^3 + x^2 - 5x - 1 + c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^3 + x^2 - 5x - 1 + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.145: Slope field plot
 $y'' + y = x^3 + x^2 + x + 1$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2}y(x) + y(x) = x^3 + x^2 + x + 1$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^3 + x^2 + x + 1 \right]$$
- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) (x+1)(x^2+1) dx \right) + \sin(x) \left(\int \cos(x) (x+1)(x^2+1) dx \right)$$
- Compute integrals

$$y_p(x) = x^3 + x^2 - 5x - 1$$
- Substitute particular solution into general solution to ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + x^3 + x^2 - 5x - 1$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)
Leaf size : 23

```

dsolve(diff(diff(y(x),x),x)+y(x) = x^3+x^2+x+1,
        y(x),singsol=all)

```

$$y = \sin(x) c_2 + \cos(x) c_1 + x^3 + x^2 - 5x - 1$$

Mathematica DSolve solution

Solving time : 0.014 (sec)
Leaf size : 26

```

DSolve[{D[y[x],{x,2}]+y[x]==1+x+x^2+x^3,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow x^3 + x^2 - 5x + c_1 \cos(x) + c_2 \sin(x) - 1$$

2.1.43 problem 43

Solved as second order linear constant coeff ode	413
Solved as second order ode using Kovacic algorithm	415
Solved as second order ode adjoint method	419
Maple step by step solution	421
Maple trace	422
Maple dsolve solution	422
Mathematica DSolve solution	423

Internal problem ID [8790]

Book : Second order enumerated odes

Section : section 1

Problem number : 43

Date solved : Thursday, December 12, 2024 at 09:47:41 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' + y = \sin(x)$$

Solved as second order linear constant coeff ode

Time used: 0.141 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x)x, \sin(x)x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x)x + A_2 \sin(x)x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x) x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{\cos(x) x}{2}\right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x) x}{2} + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.146: Slope field plot
 $y'' + y = \sin(x)$

Solved as second order ode using Kovacic algorithm

Time used: 0.135 (sec)

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \end{aligned} \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.62: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x)x, \sin(x)x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x)x + A_2 \sin(x)x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{\cos(x)x}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{\cos(x)x}{2} + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.147: Slope field plot
 $y'' + y = \sin(x)$

Solved as second order ode adjoint method

Time used: 2.253 (sec)

In normal form the ode

$$y'' + y = \sin(x) \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 0 \\ q(x) &= 1 \\ r(x) &= \sin(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + (\xi(x)) &= 0 \\ \xi''(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$\xi = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' - \frac{y(-c_1 \sin(x) + c_2 \cos(x))}{c_1 \cos(x) + c_2 \sin(x)} = \frac{-\frac{\cos(x)^2 c_1}{2} + c_2 \left(-\frac{\cos(x) \sin(x)}{2} + \frac{x}{2} \right)}{c_1 \cos(x) + c_2 \sin(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} \\ p(x) &= \frac{-c_2 \cos(x) \sin(x) - \cos(x)^2 c_1 + c_2 x}{2c_2 \sin(x) + 2c_1 \cos(x)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{-c_2 \cos(x) \sin(x) - \cos(x)^2 c_1 + c_2 x}{2c_2 \sin(x) + 2c_1 \cos(x)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{1}{c_1 \cos(x) + c_2 \sin(x)} \right) \left(\frac{-c_2 \cos(x) \sin(x) - \cos(x)^2 c_1 + c_2 x}{2c_2 \sin(x) + 2c_1 \cos(x)} \right) \\ d \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{-c_2 \cos(x) \sin(x) - \cos(x)^2 c_1 + c_2 x}{(2c_2 \sin(x) + 2c_1 \cos(x)) (c_1 \cos(x) + c_2 \sin(x))} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_1 \cos(x) + c_2 \sin(x)} &= \int \frac{-c_2 \cos(x) \sin(x) - \cos(x)^2 c_1 + c_2 x}{(2c_2 \sin(x) + 2c_1 \cos(x))(c_1 \cos(x) + c_2 \sin(x))} dx \\ &= \frac{\frac{x}{2} + \frac{x \tan(\frac{x}{2})^2}{2} - \frac{x \tan(\frac{x}{2})^4}{2} - \frac{x \tan(\frac{x}{2})^6}{2}}{\left(1 + \tan\left(\frac{x}{2}\right)\right)^2 \left(c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1\right)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x) + c_2 \sin(x)}$ gives the final solution

$$y = \frac{(2c_1 c_3 - x) \cos(x)}{2} + c_2 c_3 \sin(x)$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{(2c_1 c_3 - x) \cos(x)}{2} + c_2 c_3 \sin(x)$$

The constants can be merged to give

$$y = \frac{(2c_1 - x) \cos(x)}{2} + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{(2c_1 - x) \cos(x)}{2} + c_2 \sin(x)$$




Figure 2.148: Slope field plot
 $y'' + y = \sin(x)$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2} y(x) + y(x) = \sin(x)$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial
 $r = (-I, I)$
 - 1st solution of the homogeneous ODE
 $y_1(x) = \cos(x)$
 - 2nd solution of the homogeneous ODE
 $y_2(x) = \sin(x)$
 - General solution of the ODE
 $y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$
 - Substitute in solutions of the homogeneous ODE
 $y(x) = C1 \cos(x) + C2 \sin(x) + y_p(x)$
- Find a particular solution $y_p(x)$ of the ODE
- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
 - Compute Wronskian
 $W(y_1(x), y_2(x)) = 1$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x)^2 dx \right) + \frac{\sin(x) \left(\int \sin(2x) dx \right)}{2}$$
 - Compute integrals

$$y_p(x) = \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$
 - Substitute particular solution into general solution to ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + \frac{\sin(x)}{4} - \frac{x \cos(x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)

Leaf size : 25

```

dsolve(diff(diff(y(x),x),x)+y(x) = sin(x),
        y(x),singsol=all)

```

$$y = \frac{(-x + 2c_1) \cos(x)}{2} + \frac{\sin(x) (2c_2 + 1)}{2}$$

Mathematica DSolve solution

Solving time : 0.024 (sec)

Leaf size : 22

```
DSolve[{D[y[x],{x,2}]+y[x]==Sin[x],{}}],  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \left(-\frac{x}{2} + c_1\right) \cos(x) + c_2 \sin(x)$$

2.1.44 problem 44

Solved as second order linear constant coeff ode	424
Solved as second order ode using Kovacic algorithm	426
Solved as second order ode adjoint method	430
Maple step by step solution	432
Maple trace	433
Maple dsolve solution	433
Mathematica DSolve solution	434

Internal problem ID [8791]

Book : Second order enumerated odes

Section : section 1

Problem number : 44

Date solved : Thursday, December 12, 2024 at 09:47:45 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' + y = \cos(x)$$

Solved as second order linear constant coeff ode

Time used: 0.138 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +i \\ \lambda_2 &= -i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i \\ \lambda_2 &= -i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x)x, \sin(x)x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x)x + A_2 \sin(x)x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\sin(x)x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(\frac{\sin(x)x}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{\sin(x)x}{2} + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.149: Slope field plot
 $y'' + y = \cos(x)$

Solved as second order ode using Kovacic algorithm

Time used: 0.135 (sec)

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.64: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x)x, \sin(x)x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x)x + A_2 \sin(x)x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\sin(x)x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(\frac{\sin(x)x}{2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{\sin(x)x}{2} + c_1 \cos(x) + c_2 \sin(x)$$




Figure 2.150: Slope field plot
 $y'' + y = \cos(x)$

Solved as second order ode adjoint method

Time used: 2.262 (sec)

In normal form the ode

$$y'' + y = \cos(x) \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 0 \\ q(x) &= 1 \\ r(x) &= \cos(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + (\xi(x)) &= 0 \\ \xi''(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$A\xi''(x) + B\xi'(x) + C\xi(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $\xi = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$\xi = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$\xi = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$\xi = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' - \frac{y(-c_1 \sin(x) + c_2 \cos(x))}{c_1 \cos(x) + c_2 \sin(x)} = \frac{c_1 \left(\frac{\cos(x) \sin(x)}{2} + \frac{x}{2} \right) - \frac{\cos(x)^2 c_2}{2}}{c_1 \cos(x) + c_2 \sin(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} \\ p(x) &= \frac{c_1 \cos(x) \sin(x) - \cos(x)^2 c_2 + c_1 x}{2c_2 \sin(x) + 2c_1 \cos(x)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{-c_1 \sin(x) + c_2 \cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{1}{c_1 \cos(x) + c_2 \sin(x)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{c_1 \cos(x) \sin(x) - \cos(x)^2 c_2 + c_1 x}{2c_2 \sin(x) + 2c_1 \cos(x)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{1}{c_1 \cos(x) + c_2 \sin(x)} \right) \left(\frac{c_1 \cos(x) \sin(x) - \cos(x)^2 c_2 + c_1 x}{2c_2 \sin(x) + 2c_1 \cos(x)} \right) \\ d \left(\frac{y}{c_1 \cos(x) + c_2 \sin(x)} \right) &= \left(\frac{c_1 \cos(x) \sin(x) - \cos(x)^2 c_2 + c_1 x}{(2c_2 \sin(x) + 2c_1 \cos(x)) (c_1 \cos(x) + c_2 \sin(x))} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_1 \cos(x) + c_2 \sin(x)} &= \int \frac{c_1 \cos(x) \sin(x) - \cos(x)^2 c_2 + c_1 x}{(2c_2 \sin(x) + 2c_1 \cos(x))(c_1 \cos(x) + c_2 \sin(x))} dx \\ &= \frac{-x \tan\left(\frac{x}{2}\right) - 2x \tan\left(\frac{x}{2}\right)^3 - x \tan\left(\frac{x}{2}\right)^5 - \frac{1}{2} - \frac{\tan\left(\frac{x}{2}\right)^2}{2} + \frac{\tan\left(\frac{x}{2}\right)^4}{2} + \frac{\tan\left(\frac{x}{2}\right)^6}{2}}{\left(1 + \tan\left(\frac{x}{2}\right)^2\right)^2 \left(c_1 \tan\left(\frac{x}{2}\right)^2 - 2 \tan\left(\frac{x}{2}\right) c_2 - c_1\right)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x) + c_2 \sin(x)}$ gives the final solution

$$y = -\frac{1}{2} + (2c_1 c_3 + 1) \cos\left(\frac{x}{2}\right)^2 + \sin\left(\frac{x}{2}\right) (2c_2 c_3 + x) \cos\left(\frac{x}{2}\right) - c_1 c_3$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = -\frac{1}{2} + (2c_1 c_3 + 1) \cos\left(\frac{x}{2}\right)^2 + \sin\left(\frac{x}{2}\right) (2c_2 c_3 + x) \cos\left(\frac{x}{2}\right) - c_1 c_3$$

The constants can be merged to give

$$y = -\frac{1}{2} + (2c_1 + 1) \cos\left(\frac{x}{2}\right)^2 + \sin\left(\frac{x}{2}\right) (2c_2 + x) \cos\left(\frac{x}{2}\right) - c_1$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = -\frac{1}{2} + (2c_1 + 1) \cos\left(\frac{x}{2}\right)^2 + \sin\left(\frac{x}{2}\right) (2c_2 + x) \cos\left(\frac{x}{2}\right) - c_1$$




Figure 2.151: Slope field plot $y'' + y = \cos(x)$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2} y(x) + y(x) = \cos(x)$$

- Highest derivative means the order of the ODE is 2
- $\frac{d^2}{dx^2} y(x)$
- Characteristic polynomial of homogeneous ODE $r^2 + 1 = 0$
- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y(x) = C1y_1(x) + C2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(x) \left(\int \sin(2x) dx \right)}{2} + \sin(x) \left(\int \cos(x)^2 dx \right)$$

- Compute integrals

$$y_p(x) = \frac{\cos(x)}{4} + \frac{\sin(x)x}{2}$$

- Substitute particular solution into general solution to ODE

$$y(x) = C1 \cos(x) + C2 \sin(x) + \frac{\cos(x)}{4} + \frac{\sin(x)x}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.005 (sec)

Leaf size : 18

```

dsolve(diff(diff(y(x),x),x)+y(x) = cos(x),
        y(x),singsol=all)

```

$$y = \frac{(2c_2 + x) \sin(x)}{2} + \cos(x) c_1$$

Mathematica DSolve solution

Solving time : 0.021 (sec)

Leaf size : 28

```
DSolve[{D[y[x],{x,2}]+y[x]==Cos[x],{}}],  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{2}(x \sin(x) + \cos(x) + 2c_1 \cos(x) + 2c_2 \sin(x))$$

2.1.45 problem 45

Solved as second order missing x ode	435
Maple step by step solution	449
Maple trace	449
Maple dsolve solution	449
Mathematica DSolve solution	450

Internal problem ID [8792]

Book : Second order enumerated odes

Section : section 1

Problem number : 45

Date solved : Thursday, December 12, 2024 at 09:47:48 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$yy''^2 + y' = 0$$

Solved as second order missing x ode

Time used: 11.262 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y)^2 \left(\frac{d}{dy} p(y) \right)^2 + p(y) = 0$$

Which is now solved as first order ode for $p(y)$.

Solving for the derivative gives these ODE's to solve

$$p' = -\frac{1}{\sqrt{-py}} \tag{1}$$

$$p' = \frac{1}{\sqrt{-py}} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(y, p) dy + N(y, p) dp = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dp &= \left(-\frac{1}{\sqrt{-py}} \right) dy \\ \left(\frac{1}{\sqrt{-py}} \right) dy + dp &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(y, p) &= \frac{1}{\sqrt{-py}} \\ N(y, p) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial p} = \frac{\partial N}{\partial y}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial p} &= \frac{\partial}{\partial p} \left(\frac{1}{\sqrt{-py}} \right) \\ &= -\frac{1}{2\sqrt{-py} p} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial y} &= \frac{\partial}{\partial y} (1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial p} \neq \frac{\partial N}{\partial y}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial p} - \frac{\partial N}{\partial y} \right) \\ &= 1 \left(\left(\frac{y}{2(-py)^{3/2}} \right) - (0) \right) \\ &= -\frac{1}{2\sqrt{-py} p} \end{aligned}$$

Since A depends on p , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial y} - \frac{\partial M}{\partial p} \right) \\ &= \sqrt{-py} \left((0) - \left(\frac{y}{2(-py)^{3/2}} \right) \right) \\ &= \frac{1}{2p} \end{aligned}$$

Since B does not depend on y , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dp} \\ &= e^{\int \frac{1}{2p} \, dp} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\frac{\ln(p)}{2}} \\ &= \sqrt{p} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \sqrt{p} \left(\frac{1}{\sqrt{-py}} \right) \\ &= \frac{\sqrt{p}}{\sqrt{-py}} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \sqrt{p}(1) \\ &= \sqrt{p} \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dp}{dy} &= 0 \\ \left(\frac{\sqrt{p}}{\sqrt{-py}} \right) + (\sqrt{p}) \frac{dp}{dy} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(y, p)$

$$\frac{\partial \phi}{\partial y} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial p} = \bar{N} \tag{2}$$

Integrating (2) w.r.t. p gives

$$\int \frac{\partial \phi}{\partial p} \, dp = \int \bar{N} \, dp$$

$$\int \frac{\partial \phi}{\partial p} \, dp = \int \sqrt{p} \, dp$$

$$\phi = \frac{2p^{3/2}}{3} + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both y and p . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (1) says that $\frac{\partial \phi}{\partial y} = \frac{\sqrt{p}}{\sqrt{-py}}$. Therefore equation (4) becomes

$$\frac{\sqrt{p}}{\sqrt{-py}} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{\sqrt{p}}{\sqrt{-py}}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{\sqrt{p}}{\sqrt{-py}} \right) dy$$

$$f(y) = -\frac{2\sqrt{-py}}{\sqrt{p}} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{2p^{3/2}}{3} - \frac{2\sqrt{-py}}{\sqrt{p}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into the constant c_1 gives the solution as

$$c_1 = \frac{2p^{3/2}}{3} - \frac{2\sqrt{-py}}{\sqrt{p}}$$

Solving Eq. (2)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (B)$$

Comparing (A,B) shows that

$$\frac{\partial \phi}{\partial x} = M$$

$$\frac{\partial \phi}{\partial y} = N$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(y, p) dy + N(y, p) dp = 0 \quad (1A)$$

Therefore

$$\begin{aligned} dp &= \left(\frac{1}{\sqrt{-py}} \right) dy \\ \left(-\frac{1}{\sqrt{-py}} \right) dy + dp &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(y, p) &= -\frac{1}{\sqrt{-py}} \\ N(y, p) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial p} = \frac{\partial N}{\partial y}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial p} &= \frac{\partial}{\partial p} \left(-\frac{1}{\sqrt{-py}} \right) \\ &= \frac{1}{2\sqrt{-py} p} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial y} &= \frac{\partial}{\partial y} (1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial p} \neq \frac{\partial N}{\partial y}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial p} - \frac{\partial N}{\partial y} \right) \\ &= 1 \left(\left(-\frac{y}{2(-py)^{3/2}} \right) - (0) \right) \\ &= \frac{1}{2\sqrt{-py} p} \end{aligned}$$

Since A depends on p , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial y} - \frac{\partial M}{\partial p} \right) \\ &= -\sqrt{-py} \left((0) - \left(-\frac{y}{2(-py)^{3/2}} \right) \right) \\ &= \frac{1}{2p} \end{aligned}$$

Since B does not depend on y , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned}\mu &= e^{\int B \, dp} \\ &= e^{\int \frac{1}{2p} \, dp}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\frac{\ln(p)}{2}} \\ &= \sqrt{p}\end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \sqrt{p} \left(-\frac{1}{\sqrt{-py}} \right) \\ &= -\frac{\sqrt{p}}{\sqrt{-py}}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \sqrt{p}(1) \\ &= \sqrt{p}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dp}{dy} &= 0 \\ \left(-\frac{\sqrt{p}}{\sqrt{-py}} \right) + (\sqrt{p}) \frac{dp}{dy} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(y, p)$

$$\frac{\partial \phi}{\partial y} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial p} = \bar{N} \tag{2}$$

Integrating (2) w.r.t. p gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial p} \, dp &= \int \bar{N} \, dp \\ \int \frac{\partial \phi}{\partial p} \, dp &= \int \sqrt{p} \, dp \\ \phi &= \frac{2p^{3/2}}{3} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both y and p . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (1) says that $\frac{\partial \phi}{\partial y} = -\frac{\sqrt{p}}{\sqrt{-py}}$. Therefore equation (4) becomes

$$-\frac{\sqrt{p}}{\sqrt{-py}} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{\sqrt{p}}{\sqrt{-py}}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(-\frac{\sqrt{p}}{\sqrt{-py}} \right) dy$$

$$f(y) = \frac{2\sqrt{-py}}{\sqrt{p}} + c_2$$

Where c_2 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{2p^{3/2}}{3} + \frac{2\sqrt{-py}}{\sqrt{p}} + c_2$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_2 and c_2 constants into the constant c_2 gives the solution as

$$c_2 = \frac{2p^{3/2}}{3} + \frac{2\sqrt{-py}}{\sqrt{p}}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{2y^{3/2}}{3} - \frac{2\sqrt{-yy'}}{\sqrt{y'}} = c_1$$

Solving for the derivative gives these ODE's to solve

$$y' = \frac{(12c_1 + 24\sqrt{-y})^{2/3}}{4} \quad (1)$$

$$y' = \left(-\frac{(12c_1 + 24\sqrt{-y})^{1/3}}{4} + \frac{i\sqrt{3}(12c_1 + 24\sqrt{-y})^{1/3}}{4} \right)^2 \quad (2)$$

$$y' = \left(-\frac{(12c_1 + 24\sqrt{-y})^{1/3}}{4} - \frac{i\sqrt{3}(12c_1 + 24\sqrt{-y})^{1/3}}{4} \right)^2 \quad (3)$$

$$y' = \frac{(12c_1 - 24\sqrt{-y})^{2/3}}{4} \quad (4)$$

$$y' = \left(-\frac{(12c_1 - 24\sqrt{-y})^{1/3}}{4} - \frac{i\sqrt{3}(12c_1 - 24\sqrt{-y})^{1/3}}{4} \right)^2 \quad (5)$$

$$y' = \left(-\frac{(12c_1 - 24\sqrt{-y})^{1/3}}{4} + \frac{i\sqrt{3}(12c_1 - 24\sqrt{-y})^{1/3}}{4} \right)^2 \quad (6)$$

Now each of the above is solved separately.

Solving Eq. (1)

Integrating gives

$$\int \frac{4}{(12c_1 + 24\sqrt{-y})^{2/3}} dy = dx$$

$$-\frac{(-3c_1 + 2\sqrt{-y})(12c_1 + 24\sqrt{-y})^{1/3}}{8} = x + c_3$$

Singular solutions are found by solving

$$\frac{(12c_1 + 24\sqrt{-y})^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_1^2}{4}$$

Solving Eq. (2)

Integrating gives

$$\int \frac{16}{(12c_1 + 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2} dy = dx$$

$$-\frac{(i\sqrt{3} - 1) (12c_1 + 24\sqrt{-y})^{1/3} (-3c_1 + 2\sqrt{-y})}{16} = x + c_4$$

Singular solutions are found by solving

$$\frac{(12c_1 + 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_1^2}{4}$$

Solving Eq. (3)

Integrating gives

$$\int \frac{16}{(12c_1 + 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2} dy = dx$$

$$-\frac{3(12c_1 + 24\sqrt{-y})^{1/3} (1 + i\sqrt{3}) \left(c_1 - \frac{2\sqrt{-y}}{3}\right)}{16} = x + c_5$$

Singular solutions are found by solving

$$\frac{(12c_1 + 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_1^2}{4}$$

Solving Eq. (4)

Integrating gives

$$\int \frac{4}{(12c_1 - 24\sqrt{-y})^{2/3}} dy = dx$$

$$\frac{(3c_1 + 2\sqrt{-y}) (12c_1 - 24\sqrt{-y})^{1/3}}{8} = x + c_6$$

Singular solutions are found by solving

$$\frac{(12c_1 - 24\sqrt{-y})^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_1^2}{4}$$

Solving Eq. (5)

Integrating gives

$$\int \frac{16}{(12c_1 - 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2} dy = dx$$

$$\frac{3\left(c_1 + \frac{2\sqrt{-y}}{3}\right) (12c_1 - 24\sqrt{-y})^{1/3} (1 + i\sqrt{3})}{16} = x + c_7$$

Singular solutions are found by solving

$$\frac{(12c_1 - 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_1^2}{4}$$

Solving Eq. (6)

Integrating gives

$$\int \frac{16}{(12c_1 - 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2} dy = dx$$

$$\frac{(i\sqrt{3} - 1) (12c_1 - 24\sqrt{-y})^{1/3} (3c_1 + 2\sqrt{-y})}{16} = x + c_8$$

Singular solutions are found by solving

$$\frac{(12c_1 - 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_1^2}{4}$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{2y^{3/2}}{3} + \frac{2\sqrt{-y}y'}{\sqrt{y'}} = c_2$$

Solving for the derivative gives these ODE's to solve

$$y' = \frac{(12c_2 + 24\sqrt{-y})^{2/3}}{4} \tag{1}$$

$$y' = \left(-\frac{(12c_2 + 24\sqrt{-y})^{1/3}}{4} + \frac{i\sqrt{3} (12c_2 + 24\sqrt{-y})^{1/3}}{4} \right)^2 \tag{2}$$

$$y' = \left(-\frac{(12c_2 + 24\sqrt{-y})^{1/3}}{4} - \frac{i\sqrt{3} (12c_2 + 24\sqrt{-y})^{1/3}}{4} \right)^2 \tag{3}$$

$$y' = \frac{(12c_2 - 24\sqrt{-y})^{2/3}}{4} \quad (4)$$

$$y' = \left(-\frac{(12c_2 - 24\sqrt{-y})^{1/3}}{4} - \frac{i\sqrt{3}(12c_2 - 24\sqrt{-y})^{1/3}}{4} \right)^2 \quad (5)$$

$$y' = \left(-\frac{(12c_2 - 24\sqrt{-y})^{1/3}}{4} + \frac{i\sqrt{3}(12c_2 - 24\sqrt{-y})^{1/3}}{4} \right)^2 \quad (6)$$

Now each of the above is solved separately.

Solving Eq. (1)

Integrating gives

$$\int \frac{4}{(12c_2 + 24\sqrt{-y})^{2/3}} dy = dx$$

$$\frac{(-3c_2 + 2\sqrt{-y})(12c_2 + 24\sqrt{-y})^{1/3}}{8} = x + c_9$$

Singular solutions are found by solving

$$\frac{(12c_2 + 24\sqrt{-y})^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_2^2}{4}$$

Solving Eq. (2)

Integrating gives

$$\int \frac{16}{(12c_2 + 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2} dy = dx$$

$$\frac{(i\sqrt{3} - 1)(12c_2 + 24\sqrt{-y})^{1/3} (-3c_2 + 2\sqrt{-y})}{16} = x + C_{10}$$

Singular solutions are found by solving

$$\frac{(12c_2 + 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_2^2}{4}$$

Solving Eq. (3)

Integrating gives

$$\int \frac{16}{(12c_2 + 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2} dy = dx$$

$$\frac{3(12c_2 + 24\sqrt{-y})^{1/3} (1 + i\sqrt{3}) \left(c_2 - \frac{2\sqrt{-y}}{3} \right)}{16} = x + C_{11}$$

Singular solutions are found by solving

$$\frac{(12c_2 + 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_2^2}{4}$$

Solving Eq. (4)

Integrating gives

$$\int \frac{4}{(12c_2 - 24\sqrt{-y})^{2/3}} dy = dx$$

$$\frac{(3c_2 + 2\sqrt{-y})(12c_2 - 24\sqrt{-y})^{1/3}}{8} = x + C_{12}$$

Singular solutions are found by solving

$$\frac{(12c_2 - 24\sqrt{-y})^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_2^2}{4}$$

Solving Eq. (5)

Integrating gives

$$\int \frac{16}{(12c_2 - 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2} dy = dx$$

$$-\frac{3\left(c_2 + \frac{2\sqrt{-y}}{3}\right)(12c_2 - 24\sqrt{-y})^{1/3} (1 + i\sqrt{3})}{16} = x + C_{13}$$

Singular solutions are found by solving

$$\frac{(12c_2 - 24\sqrt{-y})^{2/3} (1 + i\sqrt{3})^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_2^2}{4}$$

Solving Eq. (6)

Integrating gives

$$\int \frac{16}{(12c_2 - 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2} dy = dx$$

$$\frac{(i\sqrt{3} - 1)(12c_2 - 24\sqrt{-y})^{1/3} (3c_2 + 2\sqrt{-y})}{16} = x + C_{14}$$

Singular solutions are found by solving

$$\frac{(12c_2 - 24\sqrt{-y})^{2/3} (i\sqrt{3} - 1)^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = -\frac{c_2^2}{4}$$

Will add steps showing solving for IC soon.

Solving for y from the above solution(s) gives (after possible removing of solutions that do not verify)

$$y = -\frac{c_1^2}{4}$$

$$y = -\frac{c_2^2}{4}$$

$$y = -\frac{c_1 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_3 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_3 + 6x)^2 c_3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_3 + 6x) c_3^2}{3}$$

$$y = -\frac{c_1 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_6 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_6 + 6x)^2 c_6}{3} + \frac{2x \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_6 + 6x) c_6^2}{3}$$

$$y = -\frac{c_2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6_C12 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6_C12 + 6x)^2 _C12}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6_C12 + 6x) _C12^2}{3}$$

$$y = -\frac{c_2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6c_9 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6c_9 + 6x)^2 c_9}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6c_9 + 6x) c_9^2}{3}$$

$$y = -\frac{i_C10\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C10 - 3i\sqrt{3}x - 6c_2_Z - 3_C10 - 3x)^2}{3} - \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C10 - 3i\sqrt{3}x - 6c_2_Z - 3_C10 - 3x)}{3}$$

$$y = -\frac{i_C14\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C14 - 3i\sqrt{3}x - 6c_2_Z - 3_C14 - 3x)^2}{3} - \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C14 - 3i\sqrt{3}x - 6c_2_Z - 3_C14 - 3x)}{3}$$

$$y = -\frac{ic_4\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_4 - 3i\sqrt{3}x - 6c_1_Z - 3c_4 - 3x)^2}{3} - \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_4 - 3i\sqrt{3}x - 6c_1_Z - 3c_4 - 3x)}{3}$$

$$y = -\frac{ic_8\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_8 - 3i\sqrt{3}x - 6c_1_Z - 3c_8 - 3x)^2}{3} - \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_8 - 3i\sqrt{3}x - 6c_1_Z - 3c_8 - 3x)}{3}$$

$$y = \frac{i_C11\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C11 + 3i\sqrt{3}x - 6c_2_Z - 3_C11 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C11 + 3i\sqrt{3}x - 6c_2_Z - 3_C11 - 3x)}{3}$$

$$y = \frac{i_C13\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C13 + 3i\sqrt{3}x - 6c_2_Z - 3_C13 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C13 + 3i\sqrt{3}x - 6c_2_Z - 3_C13 - 3x)}{3}$$

$$y = \frac{ic_5\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c_5 + 3i\sqrt{3}x - 6c_1_Z - 3c_5 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c_5 + 3i\sqrt{3}x - 6c_1_Z - 3c_5 - 3x)}{3}$$

$$y = \frac{ic_7\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c_7 + 3i\sqrt{3}x - 6c_1_Z - 3c_7 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c_7 + 3i\sqrt{3}x - 6c_1_Z - 3c_7 - 3x)}{3}$$

Summary of solutions found

$$y = -\frac{c_1^2}{4}$$

$$y = -\frac{c_2^2}{4}$$

$$y = -\frac{c_1 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_3 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_3 + 6x)^2 c_3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_3 + 6x) c_3^2}{3} - \frac{c_1^2}{4}$$

$$y = -\frac{c_1 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_6 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_6 + 6x)^2 c_6}{3} + \frac{2x \operatorname{RootOf}(_Z^4 - 6c_1_Z + 6c_6 + 6x) c_6^2}{3} - \frac{c_1^2}{4}$$

$$y = -\frac{c_2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6_C12 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6_C12 + 6x)^2 _C12}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6_C12 + 6x) _C12^2}{3} - \frac{c_2^2}{4}$$

$$y = -\frac{c_2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6c_9 + 6x)^3}{3} + \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6c_9 + 6x)^2 c_9}{3}$$

$$+ \frac{2 \operatorname{RootOf}(_Z^4 - 6c_2_Z + 6c_9 + 6x)^2 x}{3} - \frac{c_2^2}{4}$$

$$y = -\frac{i_C10\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C10 - 3i\sqrt{3}x - 6c_2_Z - 3_C10 - 3x)^2}{3}$$

$$- \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C10 - 3i\sqrt{3}x - 6c_2_Z - 3_C10 - 3x)^2}{3}$$

$$- \frac{c_2 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C10 - 3i\sqrt{3}x - 6c_2_Z - 3_C10 - 3x)^3}{3}$$

$$- \frac{_C10 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C10 - 3i\sqrt{3}x - 6c_2_Z - 3_C10 - 3x)^2}{3}$$

$$- \frac{x \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C10 - 3i\sqrt{3}x - 6c_2_Z - 3_C10 - 3x)^2}{3} - \frac{c_2^2}{4}$$

$$y = -\frac{i_C14\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C14 - 3i\sqrt{3}x - 6c_2_Z - 3_C14 - 3x)^2}{3}$$

$$- \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C14 - 3i\sqrt{3}x - 6c_2_Z - 3_C14 - 3x)^2}{3}$$

$$- \frac{c_2 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C14 - 3i\sqrt{3}x - 6c_2_Z - 3_C14 - 3x)^3}{3}$$

$$- \frac{_C14 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C14 - 3i\sqrt{3}x - 6c_2_Z - 3_C14 - 3x)^2}{3}$$

$$- \frac{x \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}_C14 - 3i\sqrt{3}x - 6c_2_Z - 3_C14 - 3x)^2}{3} - \frac{c_2^2}{4}$$

$$y = -\frac{ic_4\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_4 - 3i\sqrt{3}x - 6c_1_Z - 3c_4 - 3x)^2}{3}$$

$$- \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_4 - 3i\sqrt{3}x - 6c_1_Z - 3c_4 - 3x)^2}{3}$$

$$- \frac{c_1 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_4 - 3i\sqrt{3}x - 6c_1_Z - 3c_4 - 3x)^3}{3}$$

$$- \frac{c_4 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_4 - 3i\sqrt{3}x - 6c_1_Z - 3c_4 - 3x)^2}{3}$$

$$- \frac{x \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_4 - 3i\sqrt{3}x - 6c_1_Z - 3c_4 - 3x)^2}{3} - \frac{c_1^2}{4}$$

$$y = -\frac{ic_8\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_8 - 3i\sqrt{3}x - 6c_1_Z - 3c_8 - 3x)^2}{3}$$

$$- \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_8 - 3i\sqrt{3}x - 6c_1_Z - 3c_8 - 3x)^2}{3}$$

$$- \frac{c_1 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_8 - 3i\sqrt{3}x - 6c_1_Z - 3c_8 - 3x)^3}{3}$$

$$- \frac{c_8 \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_8 - 3i\sqrt{3}x - 6c_1_Z - 3c_8 - 3x)^2}{3}$$

$$- \frac{x \operatorname{RootOf}(_Z^4 - 3i\sqrt{3}c_8 - 3i\sqrt{3}x - 6c_1_Z - 3c_8 - 3x)^2}{3} - \frac{c_1^2}{4}$$

$$y = \frac{i_{C11}\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C11 + 3i\sqrt{3}x - 6c2_Z - 3_C11 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C11 + 3i\sqrt{3}x - 6c2_Z - 3_C11 - 3x)^2}{3} - \frac{c2 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C11 + 3i\sqrt{3}x - 6c2_Z - 3_C11 - 3x)^3}{3} - \frac{_C11 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C11 + 3i\sqrt{3}x - 6c2_Z - 3_C11 - 3x)^2}{3} - \frac{x \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C11 + 3i\sqrt{3}x - 6c2_Z - 3_C11 - 3x)^2}{3} - \frac{c2^2}{4}$$

$$y = \frac{i_{C13}\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C13 + 3i\sqrt{3}x - 6c2_Z - 3_C13 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C13 + 3i\sqrt{3}x - 6c2_Z - 3_C13 - 3x)^2}{3} - \frac{c2 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C13 + 3i\sqrt{3}x - 6c2_Z - 3_C13 - 3x)^3}{3} - \frac{_C13 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C13 + 3i\sqrt{3}x - 6c2_Z - 3_C13 - 3x)^2}{3} - \frac{x \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}_C13 + 3i\sqrt{3}x - 6c2_Z - 3_C13 - 3x)^2}{3} - \frac{c2^2}{4}$$

$$y = \frac{ic5\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c5 + 3i\sqrt{3}x - 6c1_Z - 3c5 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c5 + 3i\sqrt{3}x - 6c1_Z - 3c5 - 3x)^2}{3} - \frac{c1 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c5 + 3i\sqrt{3}x - 6c1_Z - 3c5 - 3x)^3}{3} - \frac{c5 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c5 + 3i\sqrt{3}x - 6c1_Z - 3c5 - 3x)^2}{3} - \frac{x \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c5 + 3i\sqrt{3}x - 6c1_Z - 3c5 - 3x)^2}{3} - \frac{c1^2}{4}$$

$$y = \frac{ic7\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c7 + 3i\sqrt{3}x - 6c1_Z - 3c7 - 3x)^2}{3} + \frac{ix\sqrt{3} \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c7 + 3i\sqrt{3}x - 6c1_Z - 3c7 - 3x)^2}{3} - \frac{c1 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c7 + 3i\sqrt{3}x - 6c1_Z - 3c7 - 3x)^3}{3} - \frac{c7 \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c7 + 3i\sqrt{3}x - 6c1_Z - 3c7 - 3x)^2}{3} - \frac{x \operatorname{RootOf}(_Z^4 + 3i\sqrt{3}c7 + 3i\sqrt{3}x - 6c1_Z - 3c7 - 3x)^2}{3} - \frac{c1^2}{4}$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
  *** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each res
  *** Sublevel 3 ***
Methods for second order ODEs:
  --- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-(-_b(_a)*_a)^(1/2)/
  symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 1/3*_b]

```

Maple dsolve solution

Solving time : 0.150 (sec)

Leaf size : 263

```

dsolve(y(x)*diff(diff(y(x),x),x)^2+diff(y(x),x) = 0,
  y(x),singsol=all)

```

$$\begin{aligned}
 & y = c_1 \\
 & y = 0 \\
 & - \left(\int^y \frac{-a}{(_a^{3/2} (c_1 - 3\sqrt{-a}))^{2/3}} d_a \right) - x - c_2 = 0 \\
 & - \left(\int^y \frac{-a}{(_a^{3/2} (c_1 + 3\sqrt{-a}))^{2/3}} d_a \right) - x - c_2 = 0 \\
 & \frac{-4 \left(\int^y \frac{-a}{(_a^{3/2} (c_1 - 3\sqrt{-a}))^{2/3}} d_a \right) + 2i(-x - c_2) \sqrt{3} + 2x + 2c_2}{(-i\sqrt{3} - 1)^2} = 0 \\
 & \frac{-4 \left(\int^y \frac{-a}{(_a^{3/2} (c_1 - 3\sqrt{-a}))^{2/3}} d_a \right) + 2i(x + c_2) \sqrt{3} + 2x + 2c_2}{(1 - i\sqrt{3})^2} = 0 \\
 & \frac{-4 \left(\int^y \frac{-a}{(_a^{3/2} (c_1 + 3\sqrt{-a}))^{2/3}} d_a \right) + 2i(-x - c_2) \sqrt{3} + 2x + 2c_2}{(-i\sqrt{3} - 1)^2} = 0 \\
 & \frac{-4 \left(\int^y \frac{-a}{(_a^{3/2} (c_1 + 3\sqrt{-a}))^{2/3}} d_a \right) + 2i(x + c_2) \sqrt{3} + 2x + 2c_2}{(1 - i\sqrt{3})^2} = 0
 \end{aligned}$$

Mathematica DSolve solution

Solving time : 60.97 (sec)

Leaf size : 23861

```
DSolve[{y[x]*D[y[x],{x,2}]^2+D[y[x],x]==0,{x}},  
y[x],x,IncludeSingularSolutions->True]
```

Too large to display

2.1.46 problem 46

Solved as second order missing x ode	451
Maple step by step solution	457
Maple trace	457
Maple dsolve solution	458
Mathematica DSolve solution	458

Internal problem ID [8793]

Book : Second order enumerated odes

Section : section 1

Problem number : 46

Date solved : Thursday, December 12, 2024 at 09:48:00 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$yy''^2 + y'^3 = 0$$

Solved as second order missing x ode

Time used: 2.196 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y)^2 \left(\frac{d}{dy} p(y) \right)^2 + p(y)^3 = 0$$

Which is now solved as first order ode for $p(y)$.

Solving for p' gives

$$p' = \frac{\sqrt{-py}}{y} \tag{1}$$

$$p' = -\frac{\sqrt{-py}}{y} \tag{2}$$

In canonical form, the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= \frac{\sqrt{-py}}{y} \end{aligned} \tag{1}$$

An ode of the form $p' = \frac{M(y,p)}{N(y,p)}$ is called homogeneous if the functions $M(y, p)$ and $N(y, p)$ are both homogeneous functions and of the same order. Recall that a function $f(y, p)$ is homogeneous of order n if

$$f(t^n y, t^n p) = t^n f(y, p)$$

In this case, it can be seen that both $M = \sqrt{-py}$ and $N = y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{p}{y}$, or $p = uy$. Hence

$$\frac{dp}{dy} = \frac{du}{dy}y + u$$

Applying the transformation $p = uy$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dy}y + u &= \sqrt{-u} \\ \frac{du}{dy} &= \frac{\sqrt{-u(y)} - u(y)}{y}\end{aligned}$$

Or

$$u'(y) - \frac{\sqrt{-u(y)} - u(y)}{y} = 0$$

Or

$$u'(y)y - \sqrt{-u(y)} + u(y) = 0$$

Which is now solved as separable in $u(y)$.

The ode $u'(y) = \frac{\sqrt{-u(y)} - u(y)}{y}$ is separable as it can be written as

$$\begin{aligned}u'(y) &= \frac{\sqrt{-u(y)} - u(y)}{y} \\ &= f(y)g(u)\end{aligned}$$

Where

$$\begin{aligned}f(y) &= \frac{1}{y} \\ g(u) &= \sqrt{-u} - u\end{aligned}$$

Integrating gives

$$\begin{aligned}\int \frac{1}{g(u)} du &= \int f(y) dy \\ \int \frac{1}{\sqrt{-u} - u} du &= \int \frac{1}{y} dy \\ \ln \left(\frac{1}{(\sqrt{-u(y)} + 1)^2} \right) &= \ln(y) + c_1\end{aligned}$$

We now need to find the singular solutions, these are found by finding for what values $g(u)$ is zero, since we had to divide by this above. Solving $g(u) = 0$ or $\sqrt{-u} - u = 0$ for $u(y)$ gives

$$u(y) = 0$$

Now we go over each such singular solution and check if it verifies the ode itself and any initial conditions given. If it does not then the singular solution will not be used.

Therefore the solutions found are

$$\begin{aligned}\ln \left(\frac{1}{(\sqrt{-u(y)} + 1)^2} \right) &= \ln(y) + c_1 \\ u(y) &= 0\end{aligned}$$

Converting $\ln \left(\frac{1}{(\sqrt{-u(y)+1})^2} \right) = \ln(y) + c_1$ back to p gives

$$\ln \left(\frac{1}{\left(\sqrt{-\frac{p}{y}} + 1 \right)^2} \right) = \ln(y) + c_1$$

Converting $u(y) = 0$ back to p gives

$$p = 0$$

In canonical form, the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= -\frac{\sqrt{-py}}{y} \end{aligned} \tag{1}$$

An ode of the form $p' = \frac{M(y,p)}{N(y,p)}$ is called homogeneous if the functions $M(y, p)$ and $N(y, p)$ are both homogeneous functions and of the same order. Recall that a function $f(y, p)$ is homogeneous of order n if

$$f(t^n y, t^n p) = t^n f(y, p)$$

In this case, it can be seen that both $M = -\sqrt{-py}$ and $N = y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{p}{y}$, or $p = uy$. Hence

$$\frac{dp}{dy} = \frac{du}{dy}y + u$$

Applying the transformation $p = uy$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dy}y + u &= -\sqrt{-u} \\ \frac{du}{dy} &= \frac{-\sqrt{-u(y)} - u(y)}{y} \end{aligned}$$

Or

$$u'(y) - \frac{-\sqrt{-u(y)} - u(y)}{y} = 0$$

Or

$$u'(y)y + \sqrt{-u(y)} + u(y) = 0$$

Which is now solved as separable in $u(y)$.

The ode $u'(y) = -\frac{\sqrt{-u(y)}+u(y)}{y}$ is separable as it can be written as

$$\begin{aligned} u'(y) &= -\frac{\sqrt{-u(y)} + u(y)}{y} \\ &= f(y)g(u) \end{aligned}$$

Where

$$\begin{aligned} f(y) &= \frac{1}{y} \\ g(u) &= -\sqrt{-u} - u \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(u)} du &= \int f(y) dy \\ \int \frac{1}{-\sqrt{-u} - u} du &= \int \frac{1}{y} dy \\ -2 \ln \left(\sqrt{-u(y)} - 1 \right) &= \ln(y) + c_2 \end{aligned}$$

We now need to find the singular solutions, these are found by finding for what values $g(u)$ is zero, since we had to divide by this above. Solving $g(u) = 0$ or $-\sqrt{-u} - u = 0$ for $u(y)$ gives

$$\begin{aligned}u(y) &= -1 \\u(y) &= 0\end{aligned}$$

Now we go over each such singular solution and check if it verifies the ode itself and any initial conditions given. If it does not then the singular solution will not be used.

Therefore the solutions found are

$$\begin{aligned}-2 \ln \left(\sqrt{-u(y)} - 1 \right) &= \ln(y) + c_2 \\u(y) &= -1 \\u(y) &= 0\end{aligned}$$

Converting $-2 \ln \left(\sqrt{-u(y)} - 1 \right) = \ln(y) + c_2$ back to p gives

$$-2 \ln \left(\sqrt{-\frac{p}{y}} - 1 \right) = \ln(y) + c_2$$

Converting $u(y) = -1$ back to p gives

$$p = -y$$

Converting $u(y) = 0$ back to p gives

$$p = 0$$

Solving for p gives

$$\begin{aligned}p &= \left(-\frac{2y e^{c_1} (\sqrt{y e^{c_1}} - 1)}{\sqrt{y e^{c_1}}} + y e^{c_1} - 1 \right) e^{-c_1} \\p &= \left(-\frac{2y e^{c_1} (\sqrt{y e^{c_1}} + 1)}{\sqrt{y e^{c_1}}} + y e^{c_1} - 1 \right) e^{-c_1}\end{aligned}$$

Solving for p gives

$$\begin{aligned}p &= -\left(\frac{2y e^{c_2} (\sqrt{y e^{c_2}} - 1)}{\sqrt{y e^{c_2}}} - y e^{c_2} + 1 \right) e^{-c_2} \\p &= -\left(\frac{2y e^{c_2} (\sqrt{y e^{c_2}} + 1)}{\sqrt{y e^{c_2}}} - y e^{c_2} + 1 \right) e^{-c_2}\end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 0$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned}\int dy &= \int 0 dx + c_3 \\y &= c_3\end{aligned}$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \left(-\frac{2y e^{c_1} (\sqrt{y e^{c_1}} - 1)}{\sqrt{y e^{c_1}}} + y e^{c_1} - 1 \right) e^{-c_1}$$

Integrating gives

$$\int -\frac{\sqrt{y e^{c_1}} e^{c_1}}{y e^{c_1} \sqrt{y e^{c_1}} - 2y e^{c_1} + \sqrt{y e^{c_1}}} dy = dx$$

$$\frac{2 + (-2\sqrt{y e^{c_1}} + 2) \ln(\sqrt{y e^{c_1}} - 1)}{\sqrt{y e^{c_1}} - 1} = x + c_4$$

Singular solutions are found by solving

$$-\frac{(y e^{c_1} \sqrt{y e^{c_1}} - 2y e^{c_1} + \sqrt{y e^{c_1}}) e^{-c_1}}{\sqrt{y e^{c_1}}} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{-c_1}$$

Solving for y gives

$$y = \frac{\left(\text{LambertW}\left(e^{\frac{c_4}{2} + \frac{x}{2}}\right)^2 + 2 \text{LambertW}\left(e^{\frac{c_4}{2} + \frac{x}{2}}\right) + 1 \right) e^{-c_1}}{\text{LambertW}\left(e^{\frac{c_4}{2} + \frac{x}{2}}\right)^2}$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \left(-\frac{2y e^{c_1} (1 + \sqrt{y e^{c_1}})}{\sqrt{y e^{c_1}}} + y e^{c_1} - 1 \right) e^{-c_1}$$

Integrating gives

$$\int -\frac{\sqrt{y e^{c_1}} e^{c_1}}{y e^{c_1} \sqrt{y e^{c_1}} + 2y e^{c_1} + \sqrt{y e^{c_1}}} dy = dx$$

$$\frac{-2 + (-2\sqrt{y e^{c_1}} - 2) \ln(\sqrt{y e^{c_1}} + 1)}{\sqrt{y e^{c_1}} + 1} = x + c_5$$

Solving for y gives

$$y = \frac{\left(\text{LambertW}\left(-e^{\frac{c_5}{2} + \frac{x}{2}}\right)^2 + 2 \text{LambertW}\left(-e^{\frac{c_5}{2} + \frac{x}{2}}\right) + 1 \right) e^{-c_1}}{\text{LambertW}\left(-e^{\frac{c_5}{2} + \frac{x}{2}}\right)^2}$$

For solution (4) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -y$$

Integrating gives

$$\int -\frac{1}{y} dy = dx$$

$$-\ln(y) = x + c_6$$

Singular solutions are found by solving

$$-y = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 0$$

Solving for y gives

$$y = e^{-x-c_6}$$

For solution (5) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\left(\frac{2y e^{c_2}(\sqrt{y e^{c_2}} - 1)}{\sqrt{y e^{c_2}}} - y e^{c_2} + 1\right) e^{-c_2}$$

Integrating gives

$$\int -\frac{\sqrt{y e^{c_2}} e^{c_2}}{y e^{c_2} \sqrt{y e^{c_2}} - 2y e^{c_2} + \sqrt{y e^{c_2}}} dy = dx$$

$$\frac{2 + (-2\sqrt{y e^{c_2}} + 2) \ln(\sqrt{y e^{c_2}} - 1)}{\sqrt{y e^{c_2}} - 1} = x + c_7$$

Singular solutions are found by solving

$$-\frac{(y e^{c_2} \sqrt{y e^{c_2}} - 2y e^{c_2} + \sqrt{y e^{c_2}}) e^{-c_2}}{\sqrt{y e^{c_2}}} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{-c_2}$$

Solving for y gives

$$y = \frac{\left(\text{LambertW}\left(e^{\frac{c_7}{2} + \frac{x}{2}}\right)^2 + 2\text{LambertW}\left(e^{\frac{c_7}{2} + \frac{x}{2}}\right) + 1\right) e^{-c_2}}{\text{LambertW}\left(e^{\frac{c_7}{2} + \frac{x}{2}}\right)^2}$$

For solution (6) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\left(\frac{2y e^{c_2}(1 + \sqrt{y e^{c_2}})}{\sqrt{y e^{c_2}}} - y e^{c_2} + 1\right) e^{-c_2}$$

Integrating gives

$$\int -\frac{\sqrt{y e^{c_2}} e^{c_2}}{y e^{c_2} \sqrt{y e^{c_2}} + 2y e^{c_2} + \sqrt{y e^{c_2}}} dy = dx$$

$$\frac{-2 + (-2\sqrt{y e^{c_2}} - 2) \ln(\sqrt{y e^{c_2}} + 1)}{\sqrt{y e^{c_2}} + 1} = x + c_8$$

Solving for y gives

$$y = \frac{\left(\text{LambertW}\left(-e^{\frac{c_8}{2} + \frac{x}{2}}\right)^2 + 2\text{LambertW}\left(-e^{\frac{c_8}{2} + \frac{x}{2}}\right) + 1\right) e^{-c_2}}{\text{LambertW}\left(-e^{\frac{c_8}{2} + \frac{x}{2}}\right)^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 0$$

$$y = c_3$$

$$y = e^{-c_1}$$

$$y = e^{-c_2}$$

$$y = \frac{\left(\text{LambertW}\left(-e^{\frac{c_5}{2} + \frac{x}{2}}\right)^2 + 2 \text{LambertW}\left(-e^{\frac{c_5}{2} + \frac{x}{2}}\right) + 1 \right) e^{-c_1}}{\text{LambertW}\left(-e^{\frac{c_5}{2} + \frac{x}{2}}\right)^2}$$

$$y = \frac{\left(\text{LambertW}\left(-e^{\frac{c_8}{2} + \frac{x}{2}}\right)^2 + 2 \text{LambertW}\left(-e^{\frac{c_8}{2} + \frac{x}{2}}\right) + 1 \right) e^{-c_2}}{\text{LambertW}\left(-e^{\frac{c_8}{2} + \frac{x}{2}}\right)^2}$$

$$y = \frac{\left(\text{LambertW}\left(e^{\frac{c_4}{2} + \frac{x}{2}}\right)^2 + 2 \text{LambertW}\left(e^{\frac{c_4}{2} + \frac{x}{2}}\right) + 1 \right) e^{-c_1}}{\text{LambertW}\left(e^{\frac{c_4}{2} + \frac{x}{2}}\right)^2}$$

$$y = \frac{\left(\text{LambertW}\left(e^{\frac{c_7}{2} + \frac{x}{2}}\right)^2 + 2 \text{LambertW}\left(e^{\frac{c_7}{2} + \frac{x}{2}}\right) + 1 \right) e^{-c_2}}{\text{LambertW}\left(e^{\frac{c_7}{2} + \frac{x}{2}}\right)^2}$$

$$y = e^{-x-c_6}$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each res
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
Try integration with the canonical coordinates of the symmetry [0, y]
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(-_b(_a))^(3/2)-_b(_a)^2,
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[1, 0]

```

Maple dsolve solution

Solving time : 0.331 (sec)

Leaf size : 166

```
dsolve(y(x)*diff(diff(y(x),x),x)^2+diff(y(x),x)^3 = 0,
      y(x),singsol=all)
```

$$y = c_1$$

$$y = 0$$

$$y = \frac{c_2(\text{LambertW}(c_1 e^{\frac{x}{2}-1}) + 1)^2}{\text{LambertW}(c_1 e^{\frac{x}{2}-1})^2}$$

$$y = \frac{c_2(\text{LambertW}(-c_1 e^{\frac{x}{2}-1}) + 1)^2}{\text{LambertW}(-c_1 e^{\frac{x}{2}-1})^2}$$

$$y$$

$$= e^{-\left(\int e^{2\text{RootOf}(e^{-Z}\ln((e^{-Z}+1)^2)+c_1 e^{-Z}-2e^{-Z}-Z+x e^{-Z}+\ln((e^{-Z}+1)^2)+c_1-2-Z+x-2)} dx\right)-2\left(\int e^{\text{RootOf}(e^{-Z}\ln((e^{-Z}+1)^2)+c_1 e^{-Z}-2e^{-Z}-Z+x e^{-Z}+\ln((e^{-Z}+1)^2)+c_1-2-Z+x-2)} dx\right)}$$

Mathematica DSolve solution

Solving time : 3.059 (sec)

Leaf size : 361

```
DSolve[{y[x]*D[y[x],{x,2}]^2+D[y[x],x]^3==0,{}},
      y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \text{InverseFunction}\left[-4\left(\frac{1}{2}\log\left(2\sqrt{\#1}-ic_1\right)-\frac{ic_1}{2(2\sqrt{\#1}-ic_1)}\right)\&\right][x+c_2]$$

$$y(x) \rightarrow \text{InverseFunction}\left[-4\left(\frac{ic_1}{2(2\sqrt{\#1}+ic_1)}+\frac{1}{2}\log\left(2\sqrt{\#1}+ic_1\right)\right)\&\right][x+c_2]$$

$$y(x) \rightarrow \text{InverseFunction}\left[-4\left(\frac{1}{2}\log\left(2\sqrt{\#1}-i(-c_1)\right)-\frac{i(-c_1)}{2(2\sqrt{\#1}-i(-c_1))}\right)\&\right][x+c_2]$$

$$y(x) \rightarrow \text{InverseFunction}\left[-4\left(\frac{i(-c_1)}{2(2\sqrt{\#1}+i(-1)c_1)}+\frac{1}{2}\log\left(2\sqrt{\#1}+i(-1)c_1\right)\right)\&\right][x+c_2]$$

$$y(x) \rightarrow \text{InverseFunction}\left[-4\left(\frac{1}{2}\log\left(2\sqrt{\#1}-ic_1\right)-\frac{ic_1}{2(2\sqrt{\#1}-ic_1)}\right)\&\right][x+c_2]$$

$$y(x) \rightarrow \text{InverseFunction}\left[-4\left(\frac{ic_1}{2(2\sqrt{\#1}+ic_1)}+\frac{1}{2}\log\left(2\sqrt{\#1}+ic_1\right)\right)\&\right][x+c_2]$$

2.1.47 problem 47

Solved as second order missing x ode	459
Maple step by step solution	461
Maple trace	461
Maple dsolve solution	462
Mathematica DSolve solution	462

Internal problem ID [8794]

Book : Second order enumerated odes

Section : section 1

Problem number : 47

Date solved : Thursday, December 12, 2024 at 09:48:02 AM

CAS classification :

[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]

Solve

$$y^2 y''^2 + y' = 0$$

Solved as second order missing x ode

Time used: 5.322 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$y^2 p(y)^2 \left(\frac{d}{dy} p(y) \right)^2 + p(y) = 0$$

Which is now solved as first order ode for $p(y)$.

Solving for the derivative gives these ODE's to solve

$$p' = -\frac{1}{\sqrt{-p}y} \tag{1}$$

$$p' = \frac{1}{\sqrt{-p}y} \tag{2}$$

Now each of the above is solved separately.

Solving Eq. (1)

The ode $p' = -\frac{1}{\sqrt{-p}y}$ is separable as it can be written as

$$\begin{aligned} p' &= -\frac{1}{\sqrt{-p}y} \\ &= f(y)g(p) \end{aligned}$$

Where

$$f(y) = -\frac{1}{y}$$

$$g(p) = \frac{1}{\sqrt{-p}}$$

Integrating gives

$$\int \frac{1}{g(p)} dp = \int f(y) dy$$

$$\int \sqrt{-p} dp = \int -\frac{1}{y} dy$$

$$-\frac{2(-p)^{3/2}}{3} = \ln\left(\frac{1}{y}\right) + c_1$$

Solving for p gives

$$p = -\frac{\left(-12 \ln\left(\frac{1}{y}\right) - 12c_1\right)^{2/3}}{4}$$

Solving Eq. (2)

The ode $p' = \frac{1}{\sqrt{-p}y}$ is separable as it can be written as

$$p' = \frac{1}{\sqrt{-p}y}$$

$$= f(y)g(p)$$

Where

$$f(y) = \frac{1}{y}$$

$$g(p) = \frac{1}{\sqrt{-p}}$$

Integrating gives

$$\int \frac{1}{g(p)} dp = \int f(y) dy$$

$$\int \sqrt{-p} dp = \int \frac{1}{y} dy$$

$$-\frac{2(-p)^{3/2}}{3} = \ln(y) + c_2$$

Solving for p gives

$$p = -\frac{(-12 \ln(y) - 12c_2)^{2/3}}{4}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{(-12 \ln(y) - 12c_2)^{2/3}}{4}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y -\frac{4}{(-12 \ln(\tau) - 12c_2)^{2/3}} d\tau = x + c_3$$

Singular solutions are found by solving

$$-\frac{(-12 \ln(y) - 12c_2)^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{-c_2}$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{\left(-12 \ln\left(\frac{1}{y}\right) - 12c_1\right)^{2/3}}{4}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y -\frac{4}{\left(-12 \ln\left(\frac{1}{\tau}\right) - 12c_1\right)^{2/3}} d\tau = x + c_4$$

Singular solutions are found by solving

$$-\frac{\left(-12 \ln\left(\frac{1}{y}\right) - 12c_1\right)^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{c_1}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\int^y -\frac{4}{\left(-12 \ln(\tau) - 12c_2\right)^{2/3}} d\tau = x + c_3$$

$$\int^y -\frac{4}{\left(-12 \ln\left(\frac{1}{\tau}\right) - 12c_1\right)^{2/3}} d\tau = x + c_4$$

$$y = e^{-c_2}$$

$$y = e^{c_1}$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each res
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-(-_b(_a))^(1/2)/_a =
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 0]

```

Maple dsolve solution

Solving time : 0.142 (sec)

Leaf size : 233

```
dsolve(diff(diff(y(x),x),x)^2*y(x)^2+diff(y(x),x) = 0,
        y(x),singsol=all)
```

$$\begin{aligned}
 & y = c_1 \\
 & y = 0 \\
 & -4 \left(\int^y \frac{1}{(-12 \ln(_a) + 8c_1)^{2/3}} d_a \right) - x - c_2 = 0 \\
 & -4 \left(\int^y \frac{1}{(12 \ln(_a) - 8c_1)^{2/3}} d_a \right) - x - c_2 = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(-12 \ln(_a) + 8c_1)^{2/3}} d_a \right) + 2i(-x - c_2) \sqrt{3} + 2x + 2c_2}{(-i\sqrt{3} - 1)^2} = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(-12 \ln(_a) + 8c_1)^{2/3}} d_a \right) + 2i(x + c_2) \sqrt{3} + 2x + 2c_2}{(1 - i\sqrt{3})^2} = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(12 \ln(_a) - 8c_1)^{2/3}} d_a \right) + 2i(-x - c_2) \sqrt{3} + 2x + 2c_2}{(-i\sqrt{3} - 1)^2} = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(12 \ln(_a) - 8c_1)^{2/3}} d_a \right) + 2i(x + c_2) \sqrt{3} + 2x + 2c_2}{(1 - i\sqrt{3})^2} = 0
 \end{aligned}$$

Mathematica DSolve solution

Solving time : 3.841 (sec)

Leaf size : 449

```
DSolve[{y[x]^2*D[y[x],{x,2}]^2+D[y[x],x]==0,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$\begin{aligned}
 & y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{-ic_1} (-\log(\#1) - ic_1)^{2/3} \Gamma\left(\frac{1}{3}, -ic_1 - \log(\#1)\right)}{(c_1 - i \log(\#1))^{2/3}} \& \right] [x + c_2] \\
 & y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{ic_1} (-\log(\#1) + ic_1)^{2/3} \Gamma\left(\frac{1}{3}, ic_1 - \log(\#1)\right)}{(i \log(\#1) + c_1)^{2/3}} \& \right] [x + c_2] \\
 & y(x) \\
 & \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{-i(-c_1)} (-\log(\#1) - i(-1)c_1)^{2/3} \Gamma\left(\frac{1}{3}, -i(-1)c_1 - \log(\#1)\right)}{(-i \log(\#1) - c_1)^{2/3}} \& \right] [x + c_2] \\
 & y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{-ic_1} (-\log(\#1) - ic_1)^{2/3} \Gamma\left(\frac{1}{3}, -ic_1 - \log(\#1)\right)}{(c_1 - i \log(\#1))^{2/3}} \& \right] [x + c_2] \\
 & y(x) \\
 & \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{i(-c_1)} (-\log(\#1) + i(-c_1))^{2/3} \Gamma\left(\frac{1}{3}, i(-c_1) - \log(\#1)\right)}{(i \log(\#1) - c_1)^{2/3}} \& \right] [x + c_2] \\
 & y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{ic_1} (-\log(\#1) + ic_1)^{2/3} \Gamma\left(\frac{1}{3}, ic_1 - \log(\#1)\right)}{(i \log(\#1) + c_1)^{2/3}} \& \right] [x + c_2]
 \end{aligned}$$

2.1.48 problem 48

Solved as second order missing x ode	463
Maple step by step solution	485
Maple trace	485
Maple dsolve solution	486
Mathematica DSolve solution	487

Internal problem ID [8795]

Book : Second order enumerated odes

Section : section 1

Problem number : 48

Date solved : Thursday, December 12, 2024 at 09:48:08 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$yy''^4 + y'^2 = 0$$

Solved as second order missing x ode

Time used: 56.729 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y)^4 \left(\frac{d}{dy} p(y) \right)^4 + p(y)^2 = 0$$

Which is now solved as first order ode for $p(y)$.

Solving for the derivative gives these ODE's to solve

$$p' = \frac{(-p^2 y^3)^{1/4}}{py} \tag{1}$$

$$p' = \frac{i(-p^2 y^3)^{1/4}}{py} \tag{2}$$

$$p' = -\frac{(-p^2 y^3)^{1/4}}{py} \tag{3}$$

$$p' = -\frac{i(-p^2 y^3)^{1/4}}{py} \tag{4}$$

Now each of the above is solved separately.

Solving Eq. (1)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(y, p) dy + N(y, p) dp = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dp &= \left(\frac{(-p^2 y^3)^{1/4}}{py} \right) dy \\ \left(-\frac{(-p^2 y^3)^{1/4}}{py} \right) dy + dp &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(y, p) &= -\frac{(-p^2 y^3)^{1/4}}{py} \\ N(y, p) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial p} = \frac{\partial N}{\partial y}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial p} &= \frac{\partial}{\partial p} \left(-\frac{(-p^2 y^3)^{1/4}}{py} \right) \\ &= -\frac{y^2}{2(-p^2 y^3)^{3/4}} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial y} &= \frac{\partial}{\partial y}(1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial p} \neq \frac{\partial N}{\partial y}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial p} - \frac{\partial N}{\partial y} \right) \\ &= 1 \left(\left(\frac{(-p^2 y^3)^{1/4}}{p^2 y} + \frac{y^2}{2(-p^2 y^3)^{3/4}} \right) - (0) \right) \\ &= -\frac{y^2}{2(-p^2 y^3)^{3/4}} \end{aligned}$$

Since A depends on p , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial y} - \frac{\partial M}{\partial p} \right) \\ &= -\frac{py}{(-p^2 y^3)^{1/4}} \left((0) - \left(\frac{(-p^2 y^3)^{1/4}}{p^2 y} + \frac{y^2}{2(-p^2 y^3)^{3/4}} \right) \right) \\ &= \frac{1}{2p} \end{aligned}$$

Since B does not depend on y , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dp} \\ &= e^{\int \frac{1}{2p} \, dp} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\frac{\ln(p)}{2}} \\ &= \sqrt{p} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \sqrt{p} \left(-\frac{(-p^2 y^3)^{1/4}}{py} \right) \\ &= -\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \sqrt{p}(1) \\ &= \sqrt{p} \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dp}{dy} &= 0 \\ \left(-\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y} \right) + (\sqrt{p}) \frac{dp}{dy} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(y, p)$

$$\frac{\partial \phi}{\partial y} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial p} = \bar{N} \tag{2}$$

Integrating (2) w.r.t. p gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial p} dp &= \int \bar{N} dp \\ \int \frac{\partial \phi}{\partial p} dp &= \int \sqrt{p} dp \\ \phi &= \frac{2p^{3/2}}{3} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both y and p . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (1) says that $\frac{\partial \phi}{\partial y} = -\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y}$. Therefore equation (4) becomes

$$-\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(-\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y} \right) dy \\ f(y) &= -\frac{4(-p^2 y^3)^{1/4}}{3\sqrt{p}} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{2p^{3/2}}{3} - \frac{4(-p^2 y^3)^{1/4}}{3\sqrt{p}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into the constant c_1 gives the solution as

$$c_1 = \frac{2p^{3/2}}{3} - \frac{4(-p^2 y^3)^{1/4}}{3\sqrt{p}}$$

Solving Eq. (2)

Writing the ode as

$$\begin{aligned}p' &= \frac{i(-p^2 y^3)^{1/4}}{py} \\ p' &= \omega(y, p)\end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_y + \omega(\eta_p - \xi_y) - \omega^2 \xi_p - \omega_y \xi - \omega_p \eta = 0 \quad (\text{A})$$

To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = pa_3 + ya_2 + a_1 \quad (\text{1E})$$

$$\eta = pb_3 + yb_2 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{i(-p^2y^3)^{1/4}(b_3 - a_2)}{py} + \frac{\sqrt{-p^2y^3}a_3}{p^2y^2} - \left(-\frac{i(-p^2y^3)^{1/4}}{py^2} - \frac{3ipy}{4(-p^2y^3)^{3/4}} \right) (pa_3 + ya_2 + a_1) - \left(-\frac{i(-p^2y^3)^{1/4}}{p^2y} - \frac{iy^2}{2(-p^2y^3)^{3/4}} \right) (pb_3 + yb_2 + b_1) = 0 \quad (\text{5E})$$

Putting the above in normal form gives

$$\frac{ip^4y^3a_3 - 3ip^3y^4a_2 + 6ip^3y^4b_3 + 2ip^2y^5b_2 + ip^3y^3a_1 + 2ip^2y^4b_1 - 4b_2p^2y^2(-p^2y^3)^{3/4} - 4(-p^2y^3)^{5/4}a_3}{4p^2y^2(-p^2y^3)^{3/4}} = 0$$

Setting the numerator to zero gives

$$-ip^4y^3a_3 + 3ip^3y^4a_2 - 6ip^3y^4b_3 - 2ip^2y^5b_2 - ip^3y^3a_1 - 2ip^2y^4b_1 + 4b_2p^2y^2(-p^2y^3)^{3/4} + 4(-p^2y^3)^{5/4}a_3 = 0 \quad (\text{6E})$$

Since the PDE has radicals, simplifying gives

$$-p^2y^2 \left(ip^2ya_3 - 3ipy^2a_2 + 6ipy^2b_3 + 2iy^3b_2 + ipya_1 + 2iy^2b_1 - 4(-p^2y^3)^{3/4}b_2 + 4(-p^2y^3)^{1/4}ya_3 \right) = 0$$

Looking at the above PDE shows the following are all the terms with $\{p, y\}$ in them.

$$\{p, y, (-p^2y^3)^{1/4}, (-p^2y^3)^{3/4}\}$$

The following substitution is now made to be able to collect on all terms with $\{p, y\}$ in them

$$\{p = v_1, y = v_2, (-p^2y^3)^{1/4} = v_3, (-p^2y^3)^{3/4} = v_4\}$$

The above PDE (6E) now becomes

$$-v_1^2v_2^2(-3iv_1v_2^2a_2 + iv_1^2v_2a_3 + 2iv_2^3b_2 + 6iv_1v_2^2b_3 + iv_1v_2a_1 + 2iv_2^2b_1 + 4v_3v_2a_3 - 4v_4b_2) = 0 \quad (\text{7E})$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$-iv_2^3 a_3 v_1^4 + (3ia_2 - 6ib_3) v_1^3 v_2^4 - ia_1 v_1^3 v_2^3 - 2ib_2 v_1^2 v_2^5 - 2ib_1 v_1^2 v_2^4 - 4a_3 v_3 v_1^2 v_2^3 + 4v_4 b_2 v_1^2 v_2^2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -2ib_1 &= 0 \\ -2ib_2 &= 0 \\ -ia_1 &= 0 \\ -ia_3 &= 0 \\ -4a_3 &= 0 \\ 4b_2 &= 0 \\ 3ia_2 - 6ib_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= a_2 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= \frac{a_2}{2} \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= y \\ \eta &= \frac{p}{2} \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(y, p) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dy}{\xi} = \frac{dp}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial y} + \eta \frac{\partial}{\partial p}\right) S(y, p) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned} \frac{dp}{dy} &= \frac{\eta}{\xi} \\ &= \frac{\frac{p}{2}}{y} \\ &= \frac{p}{2y} \end{aligned}$$

This is easily solved to give

$$p = c_1 \sqrt{y}$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = \frac{p}{\sqrt{y}}$$

And S is found from

$$\begin{aligned} dS &= \frac{dy}{\xi} \\ &= \frac{dy}{y} \end{aligned}$$

Integrating gives

$$\begin{aligned} S &= \int \frac{dy}{T} \\ &= \ln(y) \end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_y + \omega(y, p)S_p}{R_y + \omega(y, p)R_p} \quad (2)$$

Where in the above R_y, R_p, S_y, S_p are all partial derivatives and $\omega(y, p)$ is the right hand side of the original ode given by

$$\omega(y, p) = \frac{i(-p^2 y^3)^{1/4}}{py}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_y &= -\frac{p}{2y^{3/2}} \\ R_p &= \frac{1}{\sqrt{y}} \\ S_y &= \frac{1}{y} \\ S_p &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{2\sqrt{y}p}{2i(-p^2 y^3)^{1/4} - p^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for y, p in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{2R}{(-1+i)\sqrt{2}\sqrt{R} - R^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S .

Since the ode has the form $\frac{d}{dR}S(R) = f(R)$, then we only need to integrate $f(R)$.

$$\begin{aligned} \int dS &= \int \frac{2R}{i\sqrt{R}\sqrt{2} - \sqrt{R}\sqrt{2} - R^2} dR \\ S(R) &= -\frac{(4i\sqrt{2} + 4\sqrt{2})\sqrt{2} \left(\ln \left(\frac{R^3 + 2R^{3/2}\sqrt{2} + 4}{R^3 - 2R^{3/2}\sqrt{2} + 4} \right) + 2 \arctan \left(\frac{R^{3/2}\sqrt{2}}{2} + 1 \right) + 2 \arctan \left(\frac{R^{3/2}\sqrt{2}}{2} - 1 \right) \right)}{48} \end{aligned} \quad (i)$$

$$S(R) = \left(-\frac{1}{6} - \frac{i}{6} \right) \ln \left(\frac{R^3 + 2R^{3/2}\sqrt{2} + 4}{R^3 - 2R^{3/2}\sqrt{2} + 4} \right) + \left(\frac{1}{6} - \frac{i}{6} \right) \ln \left(\frac{R^3 - 2R^{3/2}\sqrt{2} + 4}{R^3 + 2R^{3/2}\sqrt{2} + 4} \right) + \frac{2i \arctan \left(\frac{R^3}{4} \right)}{3} -$$

To complete the solution, we just need to transform the above back to y, p coordinates. This results in

$$\ln(y) = \left(-\frac{1}{6} - \frac{i}{6}\right) \ln\left(\frac{\frac{p^3}{y^{3/2}} + 2\left(\frac{p}{\sqrt{y}}\right)^{3/2} \sqrt{2+4}}{\frac{p^3}{y^{3/2}} - 2\left(\frac{p}{\sqrt{y}}\right)^{3/2} \sqrt{2+4}}\right) + \left(\frac{1}{6} - \frac{i}{6}\right) \ln\left(\frac{\frac{p^3}{y^{3/2}} - 2\left(\frac{p}{\sqrt{y}}\right)^{3/2} \sqrt{2+4}}{\frac{p^3}{y^{3/2}} + 2\left(\frac{p}{\sqrt{y}}\right)^{3/2} \sqrt{2+4}}\right) + \frac{2i \arctan}{3}$$

Solving Eq. (3)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(y, p) dy + N(y, p) dp = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dp &= \left(-\frac{(-p^2 y^3)^{1/4}}{py}\right) dy \\ \left(\frac{(-p^2 y^3)^{1/4}}{py}\right) dy + dp &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(y, p) &= \frac{(-p^2 y^3)^{1/4}}{py} \\ N(y, p) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial p} = \frac{\partial N}{\partial y}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial p} &= \frac{\partial}{\partial p} \left(\frac{(-p^2 y^3)^{1/4}}{py} \right) \\ &= \frac{y^2}{2(-p^2 y^3)^{3/4}}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial y} &= \frac{\partial}{\partial y} (1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial p} \neq \frac{\partial N}{\partial y}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial p} - \frac{\partial N}{\partial y} \right) \\ &= 1 \left(\left(-\frac{(-p^2 y^3)^{1/4}}{p^2 y} - \frac{y^2}{2(-p^2 y^3)^{3/4}} \right) - (0) \right) \\ &= \frac{y^2}{2(-p^2 y^3)^{3/4}}\end{aligned}$$

Since A depends on p , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial y} - \frac{\partial M}{\partial p} \right) \\ &= \frac{py}{(-p^2 y^3)^{1/4}} \left((0) - \left(-\frac{(-p^2 y^3)^{1/4}}{p^2 y} - \frac{y^2}{2(-p^2 y^3)^{3/4}} \right) \right) \\ &= \frac{1}{2p}\end{aligned}$$

Since B does not depend on y , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned}\mu &= e^{\int B dp} \\ &= e^{\int \frac{1}{2p} dp}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\frac{\ln(p)}{2}} \\ &= \sqrt{p}\end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \sqrt{p} \left(\frac{(-p^2 y^3)^{1/4}}{py} \right) \\ &= \frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \sqrt{p}(1) \\ &= \sqrt{p}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dp}{dy} &= 0 \\ \left(\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y} \right) + (\sqrt{p}) \frac{dp}{dy} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(y, p)$

$$\frac{\partial \phi}{\partial y} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial p} = \bar{N} \quad (2)$$

Integrating (2) w.r.t. p gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial p} dp &= \int \bar{N} dp \\ \int \frac{\partial \phi}{\partial p} dp &= \int \sqrt{p} dp \\ \phi &= \frac{2p^{3/2}}{3} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both y and p . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (1) says that $\frac{\partial \phi}{\partial y} = \frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y}$. Therefore equation (4) becomes

$$\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{(-p^2 y^3)^{1/4}}{\sqrt{p} y} \right) dy \\ f(y) &= \frac{4(-p^2 y^3)^{1/4}}{3\sqrt{p}} + c_4\end{aligned}$$

Where c_4 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{2p^{3/2}}{3} + \frac{4(-p^2 y^3)^{1/4}}{3\sqrt{p}} + c_4$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_4 and c_2 constants into the constant c_4 gives the solution as

$$c_4 = \frac{2p^{3/2}}{3} + \frac{4(-p^2y^3)^{1/4}}{3\sqrt{p}}$$

Solving Eq. (4)

Writing the ode as

$$p' = -\frac{i(-p^2y^3)^{1/4}}{py}$$

$$p' = \omega(y, p)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_y + \omega(\eta_p - \xi_y) - \omega^2\xi_p - \omega_y\xi - \omega_p\eta = 0 \quad (\text{A})$$

To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as ansatz gives

$$\xi = pa_3 + ya_2 + a_1 \quad (\text{1E})$$

$$\eta = pb_3 + yb_2 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 - \frac{i(-p^2y^3)^{1/4}(b_3 - a_2)}{py} + \frac{\sqrt{-p^2y^3}a_3}{p^2y^2} - \left(\frac{i(-p^2y^3)^{1/4}}{py^2} + \frac{3ipy}{4(-p^2y^3)^{3/4}} \right) (pa_3 + ya_2 + a_1) - \left(\frac{i(-p^2y^3)^{1/4}}{p^2y} + \frac{iy^2}{2(-p^2y^3)^{3/4}} \right) (pb_3 + yb_2 + b_1) = 0 \quad (\text{5E})$$

Putting the above in normal form gives

$$\frac{ip^4y^3a_3 - 3ip^3y^4a_2 + 6ip^3y^4b_3 + 2ip^2y^5b_2 + ip^3y^3a_1 + 2ip^2y^4b_1 + 4b_2p^2y^2(-p^2y^3)^{3/4} + 4(-p^2y^3)^{5/4}a_3}{4p^2y^2(-p^2y^3)^{3/4}} = 0$$

Setting the numerator to zero gives

$$ip^4y^3a_3 - 3ip^3y^4a_2 + 6ip^3y^4b_3 + 2ip^2y^5b_2 + ip^3y^3a_1 + 2ip^2y^4b_1 + 4b_2p^2y^2(-p^2y^3)^{3/4} + 4(-p^2y^3)^{5/4}a_3 = 0 \quad (\text{6E})$$

Since the PDE has radicals, simplifying gives

$$p^2y^2 \left(ip^2ya_3 - 3ipy^2a_2 + 6ipy^2b_3 + 2iy^3b_2 + ipya_1 + 2iy^2b_1 + 4(-p^2y^3)^{3/4}b_2 - 4(-p^2y^3)^{1/4}ya_3 \right) = 0$$

Looking at the above PDE shows the following are all the terms with $\{p, y\}$ in them.

$$\{p, y, (-p^2y^3)^{1/4}, (-p^2y^3)^{3/4}\}$$

The following substitution is now made to be able to collect on all terms with $\{p, y\}$ in them

$$\left\{ p = v_1, y = v_2, (-p^2 y^3)^{1/4} = v_3, (-p^2 y^3)^{3/4} = v_4 \right\}$$

The above PDE (6E) now becomes

$$v_1^2 v_2^2 (-3iv_1 v_2^2 a_2 + a_3 v_1^2 v_2 i + 2b_2 v_2^3 i + 6b_3 v_1 v_2^2 i + a_1 v_1 v_2 i + 2b_1 v_2^2 i - 4v_3 v_2 a_3 + 4v_4 b_2) = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$iv_2^3 a_3 v_1^4 + (-3ia_2 + 6ib_3) v_1^3 v_2^4 + ia_1 v_1^3 v_2^3 + 2ib_2 v_1^2 v_2^5 + 2ib_1 v_1^2 v_2^4 - 4a_3 v_3 v_1^2 v_2^3 + 4v_4 b_2 v_1^2 v_2^2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} ia_1 &= 0 \\ ia_3 &= 0 \\ 2ib_1 &= 0 \\ 2ib_2 &= 0 \\ -4a_3 &= 0 \\ 4b_2 &= 0 \\ -3ia_2 + 6ib_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= a_2 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= \frac{a_2}{2} \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= y \\ \eta &= \frac{p}{2} \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(y, p) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dy}{\xi} = \frac{dp}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial y} + \eta \frac{\partial}{\partial p}\right) S(y, p) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned}\frac{dp}{dy} &= \frac{\eta}{\xi} \\ &= \frac{p}{2y} \\ &= \frac{p}{2y}\end{aligned}$$

This is easily solved to give

$$p = c_1 \sqrt{y}$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = \frac{p}{\sqrt{y}}$$

And S is found from

$$\begin{aligned}dS &= \frac{dy}{\xi} \\ &= \frac{dy}{y}\end{aligned}$$

Integrating gives

$$\begin{aligned}S &= \int \frac{dy}{y} \\ &= \ln(y)\end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_y + \omega(y, p)S_p}{R_y + \omega(y, p)R_p} \quad (2)$$

Where in the above R_y, R_p, S_y, S_p are all partial derivatives and $\omega(y, p)$ is the right hand side of the original ode given by

$$\omega(y, p) = -\frac{i(-p^2 y^3)^{1/4}}{py}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_y &= -\frac{p}{2y^{3/2}} \\ R_p &= \frac{1}{\sqrt{y}} \\ S_y &= \frac{1}{y} \\ S_p &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{2\sqrt{y}p}{2i(-p^2 y^3)^{1/4} + p^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for y, p in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{2R}{(-1+i)\sqrt{2}\sqrt{R} + R^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S .

Since the ode has the form $\frac{d}{dR}S(R) = f(R)$, then we only need to integrate $f(R)$.

$$\int dS = \int -\frac{2R}{i\sqrt{R}\sqrt{2} - \sqrt{R}\sqrt{2} + R^2} dR$$

$$S(R) = -\frac{(-4i\sqrt{2} - 4\sqrt{2})\sqrt{2} \left(\ln \left(\frac{R^3 + 2R^{3/2}\sqrt{2} + 4}{R^3 - 2R^{3/2}\sqrt{2} + 4} \right) + 2 \arctan \left(\frac{R^{3/2}\sqrt{2}}{2} + 1 \right) + 2 \arctan \left(\frac{R^{3/2}\sqrt{2}}{2} - 1 \right) \right)}{48} - \dots$$

$$S(R) = \left(\frac{1}{6} + \frac{i}{6} \right) \ln \left(\frac{R^3 + 2R^{3/2}\sqrt{2} + 4}{R^3 - 2R^{3/2}\sqrt{2} + 4} \right) + \left(-\frac{1}{6} + \frac{i}{6} \right) \ln \left(\frac{R^3 - 2R^{3/2}\sqrt{2} + 4}{R^3 + 2R^{3/2}\sqrt{2} + 4} \right) + \frac{2i \arctan \left(\frac{R^3}{4} \right)}{3} + \dots$$

To complete the solution, we just need to transform the above back to y, p coordinates.

This results in

$$\ln(y) = \left(\frac{1}{6} + \frac{i}{6} \right) \ln \left(\frac{\frac{p^3}{y^{3/2}} + 2 \left(\frac{p}{\sqrt{y}} \right)^{3/2} \sqrt{2} + 4}{\frac{p^3}{y^{3/2}} - 2 \left(\frac{p}{\sqrt{y}} \right)^{3/2} \sqrt{2} + 4} \right) + \left(-\frac{1}{6} + \frac{i}{6} \right) \ln \left(\frac{\frac{p^3}{y^{3/2}} - 2 \left(\frac{p}{\sqrt{y}} \right)^{3/2} \sqrt{2} + 4}{\frac{p^3}{y^{3/2}} + 2 \left(\frac{p}{\sqrt{y}} \right)^{3/2} \sqrt{2} + 4} \right) + \frac{2i \arctan \left(\frac{p^3}{4} \right)}{3}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{2y'^{3/2}}{3} - \frac{4(-y'^2 y^3)^{1/4}}{3\sqrt{y'}} = c_1$$

Solving for the derivative gives these ODE's to solve

$$y' = \frac{\left(16(-y^3)^{1/4} + 12c_1 \right)^{2/3}}{4} \quad (1)$$

$$y' = \left(-\frac{\left(16(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} + \frac{i\sqrt{3} \left(16(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} \right)^2 \quad (2)$$

$$y' = \left(-\frac{\left(16(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} - \frac{i\sqrt{3} \left(16(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} \right)^2 \quad (3)$$

$$y' = \frac{\left(16i(-y^3)^{1/4} + 12c_1 \right)^{2/3}}{4} \quad (4)$$

$$y' = \left(-\frac{\left(16i(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} + \frac{i\sqrt{3} \left(16i(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} \right)^2 \quad (5)$$

$$y' = \left(-\frac{\left(16i(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} - \frac{i\sqrt{3} \left(16i(-y^3)^{1/4} + 12c_1 \right)^{1/3}}{4} \right)^2 \quad (6)$$

$$y' = \frac{\left(-16(-y^3)^{1/4} + 12c_1 \right)^{2/3}}{4} \quad (7)$$

$$y' = \left(-\frac{(-16(-y^3)^{1/4} + 12c_1)^{1/3}}{4} - \frac{i\sqrt{3}(-16(-y^3)^{1/4} + 12c_1)^{1/3}}{4} \right)^2 \quad (8)$$

$$y' = \left(-\frac{(-16(-y^3)^{1/4} + 12c_1)^{1/3}}{4} + \frac{i\sqrt{3}(-16(-y^3)^{1/4} + 12c_1)^{1/3}}{4} \right)^2 \quad (9)$$

$$y' = \frac{(-16i(-y^3)^{1/4} + 12c_1)^{2/3}}{4} \quad (10)$$

$$y' = \left(-\frac{(-16i(-y^3)^{1/4} + 12c_1)^{1/3}}{4} - \frac{i\sqrt{3}(-16i(-y^3)^{1/4} + 12c_1)^{1/3}}{4} \right)^2 \quad (11)$$

$$y' = \left(-\frac{(-16i(-y^3)^{1/4} + 12c_1)^{1/3}}{4} + \frac{i\sqrt{3}(-16i(-y^3)^{1/4} + 12c_1)^{1/3}}{4} \right)^2 \quad (12)$$

Now each of the above is solved separately.

Solving Eq. (1)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{(16(-\tau^3)^{1/4} + 12c_1)^{2/3}} d\tau = x + c_7$$

Solving Eq. (2)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{(16(-\tau^3)^{1/4} + 12c_1)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + c_8$$

Solving Eq. (3)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{(16(-\tau^3)^{1/4} + 12c_1)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + c_9$$

Solving Eq. (4)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{(16i(-\tau^3)^{1/4} + 12c_1)^{2/3}} d\tau = x + _C10$$

Solving Eq. (5)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{(16i(-\tau^3)^{1/4} + 12c_1)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C11$$

Solving Eq. (6)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _ C12$$

Solving Eq. (7)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{\left(-16(-\tau^3)^{1/4} + 12c_1\right)^{2/3}} d\tau = x + _ C13$$

Singular solutions are found by solving

$$\frac{\left(-16(-y^3)^{1/4} + 12c_1\right)^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{3(-6c_1)^{1/3} c_1}{8}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

Solving Eq. (8)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _ C14$$

Singular solutions are found by solving

$$\frac{\left(-16(-y^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{3(-6c_1)^{1/3} c_1}{8}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

Solving Eq. (9)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _ C15$$

Singular solutions are found by solving

$$\frac{\left(-16(-y^3)^{1/4} + 12c_1\right)^{2/3} (i\sqrt{3} - 1)^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{3(-6c_1)^{1/3} c_1}{8}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

Solving Eq. (10)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{\left(-16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3}} d\tau = x + _C16$$

Solving Eq. (11)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C17$$

Solving Eq. (12)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C18$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{2y'^{3/2}}{3} + \frac{4(-y'^2 y^3)^{1/4}}{3\sqrt{y'}} = c_4$$

Solving for the derivative gives these ODE's to solve

$$y' = \frac{\left(16(-y^3)^{1/4} + 12c_4\right)^{2/3}}{4} \tag{1}$$

$$y' = \left(-\frac{\left(16(-y^3)^{1/4} + 12c_4\right)^{1/3}}{4} + \frac{i\sqrt{3} \left(16(-y^3)^{1/4} + 12c_4\right)^{1/3}}{4} \right)^2 \tag{2}$$

$$y' = \left(-\frac{\left(16(-y^3)^{1/4} + 12c_4\right)^{1/3}}{4} - \frac{i\sqrt{3} \left(16(-y^3)^{1/4} + 12c_4\right)^{1/3}}{4} \right)^2 \tag{3}$$

$$y' = \frac{(16i(-y^3)^{1/4} + 12c_4)^{2/3}}{4} \quad (4)$$

$$y' = \left(-\frac{(16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} + \frac{i\sqrt{3}(16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} \right)^2 \quad (5)$$

$$y' = \left(-\frac{(16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} - \frac{i\sqrt{3}(16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} \right)^2 \quad (6)$$

$$y' = \frac{(-16(-y^3)^{1/4} + 12c_4)^{2/3}}{4} \quad (7)$$

$$y' = \left(-\frac{(-16(-y^3)^{1/4} + 12c_4)^{1/3}}{4} - \frac{i\sqrt{3}(-16(-y^3)^{1/4} + 12c_4)^{1/3}}{4} \right)^2 \quad (8)$$

$$y' = \left(-\frac{(-16(-y^3)^{1/4} + 12c_4)^{1/3}}{4} + \frac{i\sqrt{3}(-16(-y^3)^{1/4} + 12c_4)^{1/3}}{4} \right)^2 \quad (9)$$

$$y' = \frac{(-16i(-y^3)^{1/4} + 12c_4)^{2/3}}{4} \quad (10)$$

$$y' = \left(-\frac{(-16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} - \frac{i\sqrt{3}(-16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} \right)^2 \quad (11)$$

$$y' = \left(-\frac{(-16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} + \frac{i\sqrt{3}(-16i(-y^3)^{1/4} + 12c_4)^{1/3}}{4} \right)^2 \quad (12)$$

Now each of the above is solved separately.

Solving Eq. (1)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{(16(-\tau^3)^{1/4} + 12c_4)^{2/3}} d\tau = x + _C19$$

Solving Eq. (2)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{(16(-\tau^3)^{1/4} + 12c_4)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C20$$

Solving Eq. (3)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{(16(-\tau^3)^{1/4} + 12c_4)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C21$$

Solving Eq. (4)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{\left(16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3}} d\tau = x + _C22$$

Solving Eq. (5)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C23$$

Solving Eq. (6)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C24$$

Solving Eq. (7)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{\left(-16(-\tau^3)^{1/4} + 12c_4\right)^{2/3}} d\tau = x + _C25$$

Singular solutions are found by solving

$$\frac{\left(-16(-y^3)^{1/4} + 12c_4\right)^{2/3}}{4} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{3(-6c_4)^{1/3} c_4}{8}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

Solving Eq. (8)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C26$$

Singular solutions are found by solving

$$\frac{\left(-16(-y^3)^{1/4} + 12c_4\right)^{2/3} (1 + i\sqrt{3})^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{3(-6c_4)^{1/3} c_4}{8}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

Solving Eq. (9)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C27$$

Singular solutions are found by solving

$$\frac{\left(-16(-y^3)^{1/4} + 12c_4\right)^{2/3} (i\sqrt{3} - 1)^2}{16} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = \frac{3(-6c_4)^{1/3} c_4}{8}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

Solving Eq. (10)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{4}{\left(-16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3}} d\tau = x + _C28$$

Solving Eq. (11)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C29$$

Solving Eq. (12)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C30$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\ln(y) = \left(-\frac{1}{6} - \frac{i}{6}\right) \ln\left(\frac{\frac{y'^3}{y^{3/2}} + 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}{\frac{y'^3}{y^{3/2}} - 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}\right) + \left(\frac{1}{6} - \frac{i}{6}\right) \ln\left(\frac{\frac{y'^3}{y^{3/2}} - 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}{\frac{y'^3}{y^{3/2}} + 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}\right) + \frac{2i \arctan}{3}$$

Unable to solve. Terminating.

For solution (4) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\ln(y) = \left(\frac{1}{6} + \frac{i}{6}\right) \ln\left(\frac{\frac{y'^3}{y^{3/2}} + 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}{\frac{y'^3}{y^{3/2}} - 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}\right) + \left(-\frac{1}{6} + \frac{i}{6}\right) \ln\left(\frac{\frac{y'^3}{y^{3/2}} - 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}{\frac{y'^3}{y^{3/2}} + 2\left(\frac{y'}{\sqrt{y}}\right)^{3/2} \sqrt{2} + 4}\right) + \frac{2i \arctan}{\dots}$$

Unable to solve. Terminating.

Will add steps showing solving for IC soon.

Summary of solutions found

$$\int^y \frac{4}{(-16(-\tau^3)^{1/4} + 12c_1)^{2/3}} d\tau = x + _ C13$$

$$\int^y \frac{4}{(16(-\tau^3)^{1/4} + 12c_1)^{2/3}} d\tau = x + c_7$$

$$\int^y \frac{4}{(-16(-\tau^3)^{1/4} + 12c_4)^{2/3}} d\tau = x + _ C25$$

$$\int^y \frac{4}{(16(-\tau^3)^{1/4} + 12c_4)^{2/3}} d\tau = x + _ C19$$

$$\int^y \frac{4}{(-16i(-\tau^3)^{1/4} + 12c_4)^{2/3}} d\tau = x + _ C28$$

$$\int^y \frac{4}{(-16i(-\tau^3)^{1/4} + 12c_1)^{2/3}} d\tau = x + _ C16$$

$$\int^y \frac{4}{(16i(-\tau^3)^{1/4} + 12c_1)^{2/3}} d\tau = x + _ C10$$

$$\int^y \frac{4}{(16i(-\tau^3)^{1/4} + 12c_4)^{2/3}} d\tau = x + _ C22$$

$$\int^y \frac{16}{(16(-\tau^3)^{1/4} + 12c_4)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _ C20$$

$$\int^y \frac{16}{(-16(-\tau^3)^{1/4} + 12c_4)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _ C26$$

$$\int^y \frac{16}{(16(-\tau^3)^{1/4} + 12c_4)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _ C21$$

$$\int^y \frac{16}{(-16(-\tau^3)^{1/4} + 12c_1)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _ C15$$

$$\int^y \frac{16}{(16(-\tau^3)^{1/4} + 12c_1)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + c_8$$

$$\int^y \frac{16}{\left(-16(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C14$$

$$\int^y \frac{16}{\left(-16(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C27$$

$$\int^y \frac{16}{\left(16(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + c_9$$

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C17$$

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C30$$

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C29$$

$$\int^y \frac{16}{\left(-16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C18$$

$$\int^y \frac{16}{\left(16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C24$$

$$\int^y \frac{16}{\left(16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (1 + i\sqrt{3})^2} d\tau = x + _C12$$

$$\int^y \frac{16}{\left(16i(-\tau^3)^{1/4} + 12c_1\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C11$$

$$\int^y \frac{16}{\left(16i(-\tau^3)^{1/4} + 12c_4\right)^{2/3} (i\sqrt{3} - 1)^2} d\tau = x + _C23$$

$$y = \frac{3(-6c_1)^{1/3} c_1}{8}$$

$$y = \frac{3(-6c_4)^{1/3} c_4}{8}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

$$y = \frac{3\left(-\frac{(-6c_1)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_1)^{1/3}}{4}\right) c_1}{4}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} - \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

$$y = \frac{3\left(-\frac{(-6c_4)^{1/3}}{4} + \frac{i\sqrt{3}(-6c_4)^{1/3}}{4}\right) c_4}{4}$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
  *** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of  $d^2y/dx^2$ : 4 solutions were found. Trying to solve each res
  *** Sublevel 3 ***
Methods for second order ODEs:
  --- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form  $\mu(x,y)$ 
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`,  $(\text{diff}(\_b(\_a), \_a)) \* \_b(\_a) - (-\_a^3 \* \_b(\_a)^2)^{(1/2)}$ 
  symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[ $\_a, 1/2 \* \_b$ ]

```

Maple dsolve solution

Solving time : 22.898 (sec)

Leaf size : 2829

```
dsolve(y(x)*diff(diff(y(x),x),x)^4+diff(y(x),x)^2 = 0,
      y(x),singsol=all)
```

$y = c_1$
 $y = 0$

$$\int^y \frac{-a^2}{\sqrt{-a^3 (2a - (c_1 a)^{1/4}) (-2a^3 + (c_1 a)^{1/4} a^2)^{1/3}}} da - x - c_2 = 0$$

$$\int^y \frac{-a^2}{\sqrt{-a^3 (i(c_1 a)^{1/4} - 2a) ((i(c_1 a)^{1/4} - 2a) a^2)^{1/3}}} da - x - c_2 = 0$$

$$\int^y \frac{-a^2}{\sqrt{-a^3 (2a + (c_1 a)^{1/4}) (-2a^3 - (c_1 a)^{1/4} a^2)^{1/3}}} da - x - c_2 = 0$$

$$\int^y \frac{-a^2}{\sqrt{-a^3 (i(c_1 a)^{1/4} + 2a) (- (i(c_1 a)^{1/4} + 2a) a^2)^{1/3}}} da - x - c_2 = 0$$

$$\sqrt{2} \left(\int^y \frac{-a^2}{\sqrt{(-2a + (c_1 a)^{1/4}) (1 + i\sqrt{3}) a^3 (-2a^3 + (c_1 a)^{1/4} a^2)^{1/3}}} da \right) - x - c_2 = 0$$

$$\sqrt{2} \left(\int^y \frac{-a^2}{\sqrt{(i - \sqrt{3}) a^3 ((i(c_1 a)^{1/4} - 2a) a^2)^{1/3} ((c_1 a)^{1/4} + 2i a)}} da \right) - x - c_2 = 0$$

$$\sqrt{2} \left(\int^y \frac{-a^2}{\sqrt{-2a^3 (1 + i\sqrt{3}) (-a + \frac{(c_1 a)^{1/4}}{2}) (-2a^3 - (c_1 a)^{1/4} a^2)^{1/3}}} da \right) - x - c_2 = 0$$

$$\sqrt{2} \left(\int^y \frac{-a^2}{\sqrt{-a^3 (i(c_1 a)^{1/4} + 2a) (- (i(c_1 a)^{1/4} + 2a) a^2)^{1/3} (1 + i\sqrt{3})}} da \right) - x - c_2 = 0$$

$$- \left(\int^y \frac{-a^2}{\sqrt{-a^3 (2a - (c_1 a)^{1/4}) (-2a^3 + (c_1 a)^{1/4} a^2)^{1/3}}} da \right) - x - c_2 = 0$$

$$- \left(\int^y \frac{-a^2}{\sqrt{-a^3 (i(c_1 a)^{1/4} - 2a) ((i(c_1 a)^{1/4} - 2a) a^2)^{1/3}}} da \right) - x - c_2 = 0$$

$$- \left(\int^y \frac{-a^2}{\sqrt{-a^3 (2a + (c_1 a)^{1/4}) (-2a^3 - (c_1 a)^{1/4} a^2)^{1/3}}} da \right) - x - c_2 = 0$$

$$- \left(\int^y \frac{-a^2}{\sqrt{-a^3 (i(c_1 a)^{1/4} + 2a) (- (i(c_1 a)^{1/4} + 2a) a^2)^{1/3}}} da \right) - x - c_2 = 0$$

$$-\sqrt{2} \left(\int^y \frac{-a^2}{\sqrt{(1 - i\sqrt{3}) a^3 (-2a + (c_1 a)^{1/4}) (-2a^3 + (c_1 a)^{1/4} a^2)^{1/3}}} da \right) - x - c_2 = 0$$

$$\sqrt{2} \left(\int^y \frac{-a^2}{\sqrt{\dots}} da \right) - x - c_2 = 0$$

Mathematica DSolve solution

Solving time : 6.004 (sec)

Leaf size : 1237

```
DSolve[{y[x]*D[y[x],{x,2}]^4+D[y[x],x]^2==0,{}},  
y[x],x,IncludeSingularSolutions->True]
```

Too large to display

2.1.49 problem 49

Solved as first order quadrature ode	488
Solved as first order homogeneous class D2 ode	489
Solved as first order ode of type differential	490
Maple step by step solution	491
Maple trace	491
Maple dsolve solution	492
Mathematica DSolve solution	492

Internal problem ID [8796]

Book : Second order enumerated odes

Section : section 1

Problem number : 49

Date solved : Thursday, December 12, 2024 at 09:49:05 AM

CAS classification :

[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]

Solve

$$y^3 y''^2 + yy' = 0$$

Factoring the ode gives these factors

$$y = 0 \tag{1}$$

$$y' = 0 \tag{2}$$

Now each of the above equations is solved in turn.

Solving equation (1)

Solving for y from

$$y = 0$$

Solving gives $y = 0$

Solving equation (2)

Solved as first order quadrature ode

Time used: 0.014 (sec)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_1$$

$$y = c_1$$




Figure 2.152: Slope field plot
 $y' = 0$

Summary of solutions found

$$y = c_1$$

Solved as first order homogeneous class D2 ode

Time used: 0.155 (sec)

Applying change of variables $y = u(x)x$, then the ode becomes

$$u'(x)x + u(x) = 0$$

Which is now solved The ode $u'(x) = -\frac{u(x)}{x}$ is separable as it can be written as

$$\begin{aligned} u'(x) &= -\frac{u(x)}{x} \\ &= f(x)g(u) \end{aligned}$$

Where

$$\begin{aligned} f(x) &= -\frac{1}{x} \\ g(u) &= u \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(u)} du &= \int f(x) dx \\ \int \frac{1}{u} du &= \int -\frac{1}{x} dx \\ \ln(u(x)) &= \ln\left(\frac{1}{x}\right) + c_1 \end{aligned}$$

We now need to find the singular solutions, these are found by finding for what values $g(u)$ is zero, since we had to divide by this above. Solving $g(u) = 0$ or $u = 0$ for $u(x)$ gives

$$u(x) = 0$$

Now we go over each such singular solution and check if it verifies the ode itself and any initial conditions given. If it does not then the singular solution will not be used.

Therefore the solutions found are

$$\begin{aligned} \ln(u(x)) &= \ln\left(\frac{1}{x}\right) + c_1 \\ u(x) &= 0 \end{aligned}$$

Solving for $u(x)$ gives

$$u(x) = \frac{e^{c_1}}{x}$$

Converting $u(x) = 0$ back to y gives

$$y = 0$$

Converting $u(x) = \frac{e^{c_1}}{x}$ back to y gives

$$y = e^{c_1}$$




Figure 2.153: Slope field plot
 $y' = 0$

Summary of solutions found

$$y = 0$$

$$y = e^{c_1}$$

Solved as first order ode of type differential

Time used: 0.011 (sec)

Writing the ode as

$$y' = 0 \tag{1}$$

Which becomes

$$(1) dy = (0) dx \tag{2}$$

But the RHS is complete differential because

$$(0) dx = d(0)$$

Hence (2) becomes

$$(1) dy = d(0)$$

Integrating gives

$$y = c_1$$




Figure 2.154: Slope field plot
 $y' = 0$

Summary of solutions found

$$y = c_1$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each res
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful
-----
* Tackling next ODE.
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`

```

Maple dsolve solution

Solving time : 0.142 (sec)

Leaf size : 233

```
dsolve(y(x)^3*diff(diff(y(x),x),x)^2+y(x)*diff(y(x),x) = 0,
y(x),singsol=all)
```

$$\begin{aligned}
 & y = c_1 \\
 & y = 0 \\
 & -4 \left(\int^y \frac{1}{(-12 \ln(_a) + 8c_1)^{2/3}} d_a \right) - x - c_2 = 0 \\
 & -4 \left(\int^y \frac{1}{(12 \ln(_a) - 8c_1)^{2/3}} d_a \right) - x - c_2 = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(-12 \ln(_a) + 8c_1)^{2/3}} d_a \right) + 2i(-x - c_2) \sqrt{3} + 2x + 2c_2}{(-i\sqrt{3} - 1)^2} = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(-12 \ln(_a) + 8c_1)^{2/3}} d_a \right) + 2i(x + c_2) \sqrt{3} + 2x + 2c_2}{(1 - i\sqrt{3})^2} = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(12 \ln(_a) - 8c_1)^{2/3}} d_a \right) + 2i(-x - c_2) \sqrt{3} + 2x + 2c_2}{(-i\sqrt{3} - 1)^2} = 0 \\
 & \frac{-16 \left(\int^y \frac{1}{(12 \ln(_a) - 8c_1)^{2/3}} d_a \right) + 2i(x + c_2) \sqrt{3} + 2x + 2c_2}{(1 - i\sqrt{3})^2} = 0
 \end{aligned}$$

Mathematica DSolve solution

Solving time : 3.77 (sec)

Leaf size : 459

```
DSolve[{y[x]^3*D[y[x],{x,2}]^2+y[x]*D[y[x],x]==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

 $y(x) \rightarrow 0$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{-ic_1} (-\log(\#1) - ic_1)^{2/3} \Gamma\left(\frac{1}{3}, -ic_1 - \log(\#1)\right)}{(c_1 - i \log(\#1))^{2/3}} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{ic_1} (-\log(\#1) + ic_1)^{2/3} \Gamma\left(\frac{1}{3}, ic_1 - \log(\#1)\right)}{(i \log(\#1) + c_1)^{2/3}} \& \right] [x + c_2]$$

 $y(x) \rightarrow 0$ $y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{-i(-c_1)} (-\log(\#1) - i(-1)c_1)^{2/3} \Gamma\left(\frac{1}{3}, -i(-1)c_1 - \log(\#1)\right)}{(-i \log(\#1) - c_1)^{2/3}} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{-ic_1} (-\log(\#1) - ic_1)^{2/3} \Gamma\left(\frac{1}{3}, -ic_1 - \log(\#1)\right)}{(c_1 - i \log(\#1))^{2/3}} \& \right] [x + c_2]$$

 $y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{i(-c_1)} (-\log(\#1) + i(-c_1))^{2/3} \Gamma\left(\frac{1}{3}, i(-c_1) - \log(\#1)\right)}{(i \log(\#1) - c_1)^{2/3}} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\left(\frac{2}{3}\right)^{2/3} e^{ic_1} (-\log(\#1) + ic_1)^{2/3} \Gamma\left(\frac{1}{3}, ic_1 - \log(\#1)\right)}{(i \log(\#1) + c_1)^{2/3}} \& \right] [x + c_2]$$

2.1.50 problem 50

Solved as second order missing x ode	493
Maple step by step solution	495
Maple trace	496
Maple dsolve solution	496
Mathematica DSolve solution	496

Internal problem ID [8797]

Book : Second order enumerated odes

Section : section 1

Problem number : 50

Date solved : Thursday, December 12, 2024 at 09:49:06 AM

CAS classification :

[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible

Solve

$$yy'' + y'^3 = 0$$

Solved as second order missing x ode

Time used: 0.480 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) + p(y)^3 = 0$$

Which is now solved as first order ode for $p(y)$.

The ode $p' = -\frac{p^2}{y}$ is separable as it can be written as

$$\begin{aligned} p' &= -\frac{p^2}{y} \\ &= f(y)g(p) \end{aligned}$$

Where

$$\begin{aligned} f(y) &= -\frac{1}{y} \\ g(p) &= p^2 \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(p)} dp &= \int f(y) dy \\ \int \frac{1}{p^2} dp &= \int -\frac{1}{y} dy \\ -\frac{1}{p} &= \ln\left(\frac{1}{y}\right) + c_1 \end{aligned}$$

We now need to find the singular solutions, these are found by finding for what values $g(p)$ is zero, since we had to divide by this above. Solving $g(p) = 0$ or $p^2 = 0$ for p gives

$$p = 0$$

Now we go over each such singular solution and check if it verifies the ode itself and any initial conditions given. If it does not then the singular solution will not be used.

Therefore the solutions found are

$$-\frac{1}{p} = \ln\left(\frac{1}{y}\right) + c_1$$

$$p = 0$$

Solving for p gives

$$p = -\frac{1}{\ln\left(\frac{1}{y}\right) + c_1}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 0$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_2$$

$$y = c_2$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{1}{\ln\left(\frac{1}{y}\right) + c_1}$$

Integrating gives

$$\int \left(-\ln\left(\frac{1}{y}\right) - c_1\right) dy = dx$$

$$-y\left(\ln\left(\frac{1}{y}\right) + c_1 + 1\right) = x + c_3$$

Solving for y gives

$$y = \frac{x + c_3}{\text{LambertW}\left((x + c_3)e^{-c_1-1}\right)}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2$$

$$y = \frac{x + c_3}{\text{LambertW}\left((x + c_3)e^{-c_1-1}\right)}$$

Maple step by step solution

Let's solve

$$\left(\frac{d^2}{dx^2}y(x)\right)y(x) + \left(\frac{d}{dx}y(x)\right)^3 = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Define new dependent variable u

$$u(x) = \frac{d}{dx}y(x)$$

- Compute $\frac{d^2}{dx^2}y(x)$

$$\frac{d}{dx}u(x) = \frac{d^2}{dx^2}y(x)$$

- Use chain rule on the lhs

$$\left(\frac{d}{dx}y(x)\right)\left(\frac{d}{dy}u(y)\right) = \frac{d^2}{dx^2}y(x)$$

- Substitute in the definition of u

$$u(y)\left(\frac{d}{dy}u(y)\right) = \frac{d^2}{dx^2}y(x)$$

- Make substitutions $\frac{d}{dx}y(x) = u(y)$, $\frac{d^2}{dx^2}y(x) = u(y)\left(\frac{d}{dy}u(y)\right)$ to reduce order of ODE

$$u(y)\left(\frac{d}{dy}u(y)\right)y + u(y)^3 = 0$$

- Solve for the highest derivative

$$\frac{d}{dy}u(y) = -\frac{u(y)^2}{y}$$

- Separate variables

$$\frac{\frac{d}{dy}u(y)}{u(y)^2} = -\frac{1}{y}$$

- Integrate both sides with respect to y

$$\int \frac{\frac{d}{dy}u(y)}{u(y)^2} dy = \int -\frac{1}{y} dy + C1$$

- Evaluate integral

$$-\frac{1}{u(y)} = -\ln(y) + C1$$

- Solve for $u(y)$

$$u(y) = \frac{1}{\ln(y) - C1}$$

- Solve 1st ODE for $u(y)$

$$u(y) = \frac{1}{\ln(y) - C1}$$

- Revert to original variables with substitution $u(y) = \frac{d}{dx}y(x)$, $y = y(x)$

$$\frac{d}{dx}y(x) = \frac{1}{\ln(y(x)) - C1}$$

- Solve for the highest derivative

$$\frac{d}{dx}y(x) = \frac{1}{\ln(y(x)) - C1}$$

- Separate variables

$$\left(\frac{d}{dx}y(x)\right)(\ln(y(x)) - C1) = 1$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}y(x)\right)(\ln(y(x)) - C1) dx = \int 1 dx + C2$$

- Evaluate integral

$$-C1y(x) + y(x)\ln(y(x)) - y(x) = x + C2$$

- Solve for $y(x)$

$$y(x) = e^{LambertW((x+C2)e^{-C1-1})+C1+1}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`

```

Maple dsolve solution

Solving time : 0.037 (sec)

Leaf size : 27

```

dsolve(y(x)*diff(diff(y(x),x),x)+diff(y(x),x)^3 = 0,
        y(x),singsol=all)

```

$$\begin{aligned}
 y &= 0 \\
 y &= c_1 \\
 y &= \frac{x + c_2}{\text{LambertW}((x + c_2)e^{c_1-1})}
 \end{aligned}$$

Mathematica DSolve solution

Solving time : 60.147 (sec)

Leaf size : 26

```

DSolve[{y[x]*D[y[x],{x,2}]+D[y[x],x]^3==0,{]},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{x + c_2}{W(e^{-1-c_1}(x + c_2))}$$

2.1.51 problem 51

Solved as first order quadrature ode	497
Solved as first order homogeneous class D2 ode	498
Solved as first order ode of type differential	499
Maple step by step solution	500
Maple trace	500
Maple dsolve solution	501
Mathematica DSolve solution	502

Internal problem ID [8798]

Book : Second order enumerated odes

Section : section 1

Problem number : 51

Date solved : Thursday, December 12, 2024 at 09:49:07 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$yy''^3 + y^3y' = 0$$

Factoring the ode gives these factors

$$y = 0 \tag{1}$$

$$y'y^2 = 0 \tag{2}$$

Now each of the above equations is solved in turn.

Solving equation (1)

Solving for y from

$$y = 0$$

Solving gives $y = 0$

Solving equation (2)

Solved as first order quadrature ode

Time used: 0.017 (sec)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_1$$

$$y = c_1$$




Figure 2.155: Slope field plot
 $y'y^2 = 0$

Summary of solutions found

$$y = c_1$$

Solved as first order homogeneous class D2 ode

Time used: 0.160 (sec)

Applying change of variables $y = u(x)x$, then the ode becomes

$$(u'(x)x + u(x))u(x)^2x^2 = 0$$

Which is now solved The ode $u'(x) = -\frac{u(x)}{x}$ is separable as it can be written as

$$\begin{aligned} u'(x) &= -\frac{u(x)}{x} \\ &= f(x)g(u) \end{aligned}$$

Where

$$\begin{aligned} f(x) &= -\frac{1}{x} \\ g(u) &= u \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(u)} du &= \int f(x) dx \\ \int \frac{1}{u} du &= \int -\frac{1}{x} dx \\ \ln(u(x)) &= \ln\left(\frac{1}{x}\right) + c_1 \end{aligned}$$

We now need to find the singular solutions, these are found by finding for what values $g(u)$ is zero, since we had to divide by this above. Solving $g(u) = 0$ or $u = 0$ for $u(x)$ gives

$$u(x) = 0$$

Now we go over each such singular solution and check if it verifies the ode itself and any initial conditions given. If it does not then the singular solution will not be used.

Therefore the solutions found are

$$\begin{aligned} \ln(u(x)) &= \ln\left(\frac{1}{x}\right) + c_1 \\ u(x) &= 0 \end{aligned}$$

Solving for $u(x)$ gives

$$u(x) = \frac{e^{c_1}}{x}$$

Converting $u(x) = 0$ back to y gives

$$y = 0$$

Converting $u(x) = \frac{e^{c_1}}{x}$ back to y gives

$$y = e^{c_1}$$




Figure 2.156: Slope field plot
 $y'y^2 = 0$

Summary of solutions found

$$y = 0$$

$$y = e^{c_1}$$

Solved as first order ode of type differential

Time used: 0.011 (sec)

Writing the ode as

$$y' = 0 \tag{1}$$

Which becomes

$$(1) dy = (0) dx \tag{2}$$

But the RHS is complete differential because

$$(0) dx = d(0)$$

Hence (2) becomes

$$(1) dy = d(0)$$

Integrating gives

$$y = c_1$$




Figure 2.157: Slope field plot
 $y'y^2 = 0$

Summary of solutions found

$$y = c_1$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 3 solutions were found. Trying to solve each result
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
Try integration with the canonical coordinates of the symmetry [0, y]
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)^2+(-_b(_a))^(1/3), _b
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[1, 0]

```

Maple dsolve solution

Solving time : 0.283 (sec)

Leaf size : 126

```
dsolve(y(x)*diff(diff(y(x),x),x)^3+y(x)^3*diff(y(x),x) = 0,
y(x),singsol=all)
```

$$y = 0$$

$$y = c_1$$

$$y = e^{\int \text{RootOf}\left(x - \left(f^{-Z} - \frac{1}{-f^2 - (-f)^{1/3}} d_f\right) + c_1\right) dx + c_2}$$

$$y = e^{\int \text{RootOf}\left(x + 2 \left(f^{-Z} \frac{1}{i\sqrt{3}(-f)^{1/3} + 2f^2 + (-f)^{1/3}} d_f\right) + c_1\right) dx + c_2}$$

$$y = e^{\int \text{RootOf}\left(x - 2 \left(f^{-Z} \frac{1}{i\sqrt{3}(-f)^{1/3} - 2f^2 - (-f)^{1/3}} d_f\right) + c_1\right) dx + c_2}$$

Mathematica DSolve solution

Solving time : 4.228 (sec)

Leaf size : 800

```
DSolve[{y[x]*D[y[x],{x,2}]^3+y[x]^3*D[y[x],x]==0,{}},
      y[x],x,IncludeSingularSolutions->True]
```

$y(x) \rightarrow 0$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 - \frac{3\#1^{5/3}}{5c_1} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, \frac{3\#1^{5/3}}{5c_1} \right)}{\left(-\#1^{5/3} + \frac{5c_1}{3} \right)^{3/5}} \& \right] [x + c_2]$$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 + \frac{3\sqrt[3]{-1}\#1^{5/3}}{5c_1} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, -\frac{3\sqrt[3]{-1}\#1^{5/3}}{5c_1} \right)}{\left(\sqrt[3]{-1}\#1^{5/3} + \frac{5c_1}{3} \right)^{3/5}} \& \right] [x + c_2]$$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 - \frac{3(-1)^{2/3}\#1^{5/3}}{5c_1} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, \frac{3(-1)^{2/3}\#1^{5/3}}{5c_1} \right)}{\left(-(-1)^{2/3}\#1^{5/3} + \frac{5c_1}{3} \right)^{3/5}} \& \right] [x + c_2]$$

$y(x) \rightarrow 0$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 - \frac{3\#1^{5/3}}{5(-c_1)} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, \frac{3\#1^{5/3}}{5(-c_1)} \right)}{\left(-\#1^{5/3} + \frac{5(-c_1)}{3} \right)^{3/5}} \& \right] [x + c_2]$$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 + \frac{3\sqrt[3]{-1}\#1^{5/3}}{5(-c_1)} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, -\frac{3\sqrt[3]{-1}\#1^{5/3}}{5(-c_1)} \right)}{\left(\sqrt[3]{-1}\#1^{5/3} + \frac{5}{3}(-1)c_1 \right)^{3/5}} \& \right] [x + c_2]$$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 - \frac{3(-1)^{2/3}\#1^{5/3}}{5(-c_1)} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, \frac{3(-1)^{2/3}\#1^{5/3}}{5(-c_1)} \right)}{\left(-(-1)^{2/3}\#1^{5/3} + \frac{5(-c_1)}{3} \right)^{3/5}} \& \right] [x + c_2]$$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 - \frac{3\#1^{5/3}}{5c_1} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, \frac{3\#1^{5/3}}{5c_1} \right)}{\left(-\#1^{5/3} + \frac{5c_1}{3} \right)^{3/5}} \& \right] [x + c_2]$$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 + \frac{3\sqrt[3]{-1}\#1^{5/3}}{5c_1} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, -\frac{3\sqrt[3]{-1}\#1^{5/3}}{5c_1} \right)}{\left(\sqrt[3]{-1}\#1^{5/3} + \frac{5c_1}{3} \right)^{3/5}} \& \right] [x + c_2]$$

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\frac{\#1 \left(1 - \frac{3(-1)^{2/3}\#1^{5/3}}{5c_1} \right)^{3/5} \text{Hypergeometric2F1} \left(\frac{3}{5}, \frac{3}{5}, \frac{8}{5}, \frac{3(-1)^{2/3}\#1^{5/3}}{5c_1} \right)}{\left(-(-1)^{2/3}\#1^{5/3} + \frac{5c_1}{3} \right)^{3/5}} \& \right] [x + c_2]$$

2.1.52 problem 52

Maple step by step solution 504
 Maple trace 504
 Maple dsolve solution 505
 Mathematica DSolve solution 506

Internal problem ID [8799]

Book : Second order enumerated odes

Section : section 1

Problem number : 52

Date solved : Thursday, December 12, 2024 at 09:49:08 AM

CAS classification : [[_2nd_order, _missing_x]]

Solve

$$yy''^3 + y^3y'^5 = 0$$

Factoring the ode gives these factors

$$y = 0 \tag{1}$$

$$y'^5 y^2 = 0 \tag{2}$$

Now each of the above equations is solved in turn.

Solving equation (1)

Solving for y from

$$y = 0$$

Solving gives $y = 0$

Solving equation (2)

Solving for the derivative gives these ODE's to solve

$$y' = 0 \tag{1}$$

$$y' = 0 \tag{2}$$

$$y' = 0 \tag{3}$$

$$y' = 0 \tag{4}$$

$$y' = 0 \tag{5}$$

Now each of the above is solved separately.

Solving Eq. (1)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_1$$

$$y = c_1$$

Solving Eq. (2)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_2$$

$$y = c_2$$

Solving Eq. (3)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_3$$

$$y = c_3$$

Solving Eq. (4)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_4$$

$$y = c_4$$

Solving Eq. (5)

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int 0 dx + c_5$$

$$y = c_5$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 3 solutions were found. Trying to solve each result
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-(-_a^2*_b(_a)^2)^(1/3)
    symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 5*_b]

```


Maple dsolve solution

Solving time : 0.311 (sec)

Leaf size : 208

```
dsolve(y(x)*diff(diff(y(x),x),x)^3+y(x)^3*diff(y(x),x)^5 = 0,
      y(x),singsol=all)
```

$$y = 0$$

$$y = c_1$$

$$\int^y \frac{1}{\text{RootOf}\left(5 \left(\int_{-g}^{-Z} \frac{1}{-a(-a^2-f)^{1/3}-5f} d-f\right) - \ln(-a^5 + 125) + 5c_1\right)} d_a - x - c_2 = 0$$

$$\int^y \frac{1}{\text{RootOf}\left(-i \ln(-a^5 + 125) + \sqrt{3} \ln(-a^5 + 125) + 20 \left(\int_{-g}^{-Z} \frac{1}{2i_a(-a^2-f)^{1/3} + 5i_f + 5\sqrt{3}_f} d-f\right) - 20\right)} d_a - x - c_2 = 0$$

$$\int^y \frac{1}{\text{RootOf}\left(\sqrt{3} \ln(-a^5 + 125) + i \ln(-a^5 + 125) + 20 \left(\int_{-g}^{-Z} \frac{1}{-2i_a(-a^2-f)^{1/3} + 5\sqrt{3}_f - 5i_f} d-f\right) - 20\right)} d_a - x - c_2 = 0$$

Mathematica DSolve solution

Solving time : 26.332 (sec)

Leaf size : 449

```
DSolve[{y[x]*D[y[x],{x,2}]^3+y[x]^3*D[y[x],x]^5==0,{}},
  y[x],x,IncludeSingularSolutions->True]
```

$y(x) \rightarrow 0$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, \frac{3 \#1^{5/3}}{5c_1} \right)}{c_1^3} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, -\frac{3i(-i+\sqrt{3}) \#1^{5/3}}{10c_1} \right)}{c_1^3} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, \frac{3i(i+\sqrt{3}) \#1^{5/3}}{10c_1} \right)}{c_1^3} \& \right] [x + c_2]$$

$y(x) \rightarrow 0$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, \frac{3 \#1^{5/3}}{5(-c_1)} \right)}{(-c_1)^3} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, -\frac{3i(-i+\sqrt{3}) \#1^{5/3}}{10(-c_1)} \right)}{(-c_1)^3} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, \frac{3i(i+\sqrt{3}) \#1^{5/3}}{10(-c_1)} \right)}{(-c_1)^3} \& \right] [x + c_2]$$

$y(x) \rightarrow 0$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, \frac{3 \#1^{5/3}}{5c_1} \right)}{c_1^3} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, -\frac{3i(-i+\sqrt{3}) \#1^{5/3}}{10c_1} \right)}{c_1^3} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{27 \#1 \text{ Hypergeometric2F1} \left(\frac{3}{5}, 3, \frac{8}{5}, \frac{3i(i+\sqrt{3}) \#1^{5/3}}{10c_1} \right)}{c_1^3} \& \right] [x + c_2]$$

2.2 section 2

2.2.1	problem 1	508
2.2.2	problem 2	511
2.2.3	problem 3	514
2.2.4	problem 4	515
2.2.5	problem 5	516
2.2.6	problem 6	522
2.2.7	problem 8	528
2.2.8	problem 9	535
2.2.9	problem 10	536
2.2.10	problem 11	539
2.2.11	problem 12	544
2.2.12	problem 13	547
2.2.13	problem 14	550
2.2.14	problem 15	551
2.2.15	problem 16	564
2.2.16	problem 17	577
2.2.17	problem 18	590
2.2.18	problem 19	606
2.2.19	problem 20	620
2.2.20	problem 21	648
2.2.21	problem 22	651
2.2.22	problem 23	653
2.2.23	problem 24	656
2.2.24	problem 25	679
2.2.25	problem 25	696
2.2.26	problem 26	705
2.2.27	problem 27	715
2.2.28	problem 28	728
2.2.29	problem 29	738
2.2.30	problem 30	745
2.2.31	problem 31	758
2.2.32	problem 32	771
2.2.33	problem 33	783
2.2.34	problem 34	797
2.2.35	problem 35	808
2.2.36	problem 36	820
2.2.37	problem 37	822
2.2.38	problem 38	833
2.2.39	problem 39	844
2.2.40	problem 40	850
2.2.41	problem 41	855
2.2.42	problem 42	858
2.2.43	problem 43	861
2.2.44	problem 44	872
2.2.45	problem 45	881
2.2.46	problem 46	887
2.2.47	problem 47	904
2.2.48	problem 48	906
2.2.49	problem 49	918

2.2.1 problem 1

Solved as second nonlinear ode solved by Mainardi Liouville method	508
Maple step by step solution	509
Maple trace	509
Maple dsolve solution	510
Mathematica DSolve solution	510

Internal problem ID [8800]

Book : Second order enumerated odes

Section : section 2

Problem number : 1

Date solved : Thursday, December 12, 2024 at 09:49:08 AM

CAS classification : [_Liouville, [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + xy' + yy'^2 = 0$$

Solved as second nonlinear ode solved by Mainardi Liouville method

Time used: 0.210 (sec)

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \tag{1A}$$

Where in this problem

$$f(x) = x$$

$$g(y) = y$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \tag{2A}$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \tag{3A}$$

And the last term in Eq (2A) can be written as

$$\begin{aligned} g \frac{dy}{dx} &= \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx} \\ &= \frac{d}{dx} \int g dy \end{aligned} \tag{4A}$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \tag{5A}$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \tag{6A}$$

Where c_2 is a new arbitrary constant. But since $g = y$ and $f = x$, then

$$\begin{aligned}\int -g dy &= \int -y dy \\ &= -\frac{y^2}{2} \\ \int -f dx &= \int -x dx \\ &= -\frac{x^2}{2}\end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = c_2 e^{-\frac{y^2}{2}} e^{-\frac{x^2}{2}}$$

Which is now solved as first order separable ode. The ode $y' = c_2 e^{-\frac{y^2}{2}} e^{-\frac{x^2}{2}}$ is separable as it can be written as

$$\begin{aligned}y' &= c_2 e^{-\frac{y^2}{2}} e^{-\frac{x^2}{2}} \\ &= f(x)g(y)\end{aligned}$$

Where

$$\begin{aligned}f(x) &= e^{-\frac{x^2}{2}} c_2 \\ g(y) &= e^{-\frac{y^2}{2}}\end{aligned}$$

Integrating gives

$$\begin{aligned}\int \frac{1}{g(y)} dy &= \int f(x) dx \\ \int e^{\frac{y^2}{2}} dy &= \int e^{-\frac{x^2}{2}} c_2 dx \\ -\frac{i\sqrt{\pi}\sqrt{2}\operatorname{erf}\left(\frac{i\sqrt{2}y}{2}\right)}{2} &= \frac{c_2\sqrt{\pi}\sqrt{2}\operatorname{erf}\left(\frac{\sqrt{2}x}{2}\right)}{2} + c_3\end{aligned}$$

Will add steps showing solving for IC soon.

Solving for y from the above solution(s) gives (after possible removing of solutions that do not verify)

$$y = -i \operatorname{RootOf}\left(i\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{2}x}{2}\right)c_2 + i\sqrt{2}c_3 - \operatorname{erf}(_Z)\sqrt{\pi}\right)\sqrt{2}$$

Summary of solutions found

$$y = -i \operatorname{RootOf}\left(i\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{2}x}{2}\right)c_2 + i\sqrt{2}c_3 - \operatorname{erf}(_Z)\sqrt{\pi}\right)\sqrt{2}$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.014 (sec)

Leaf size : 37

```
dsolve(diff(diff(y(x),x),x)+diff(y(x),x)*x+y(x)*diff(y(x),x)^2 = 0,
y(x),singsol=all)
```

$$y = -i \operatorname{RootOf} \left(i\sqrt{\pi} \operatorname{erf} \left(\frac{\sqrt{2}x}{2} \right) c_1 + i\sqrt{2} c_2 - \operatorname{erf}(_Z) \sqrt{\pi} \right) \sqrt{2}$$

Mathematica DSolve solution

Solving time : 0.169 (sec)

Leaf size : 44

```
DSolve[{D[y[x],{x,2}]+x*D[y[x],x]+y[x]*(D[y[x],x])^2==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -i\sqrt{2}\operatorname{erf}^{-1} \left(i \left(\sqrt{\frac{2}{\pi}} c_2 - c_1 \operatorname{erf} \left(\frac{x}{\sqrt{2}} \right) \right) \right)$$

2.2.2 problem 2

Solved as second nonlinear ode solved by Mainardi Liouville method	511
Maple step by step solution	512
Maple trace	512
Maple dsolve solution	512
Mathematica DSolve solution	513

Internal problem ID [8801]

Book : Second order enumerated odes

Section : section 2

Problem number : 2

Date solved : Thursday, December 12, 2024 at 09:49:09 AM

CAS classification : [_Liouville, [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + \sin(x)y' + yy'^2 = 0$$

Solved as second nonlinear ode solved by Mainardi Liouville method

Time used: 0.240 (sec)

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \quad (1A)$$

Where in this problem

$$f(x) = \sin(x)$$

$$g(y) = y$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \quad (2A)$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \quad (3A)$$

And the last term in Eq (2A) can be written as

$$\begin{aligned} g \frac{dy}{dx} &= \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx} \\ &= \frac{d}{dx} \int g dy \end{aligned} \quad (4A)$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \quad (5A)$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \quad (6A)$$

Where c_2 is a new arbitrary constant. But since $g = y$ and $f = \sin(x)$, then

$$\begin{aligned}\int -g dy &= \int -y dy \\ &= -\frac{y^2}{2} \\ \int -f dx &= \int -\sin(x) dx \\ &= \cos(x)\end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = c_2 e^{-\frac{y^2}{2}} e^{\cos(x)}$$

Which is now solved as first order separable ode. The ode $y' = c_2 e^{-\frac{y^2}{2}} e^{\cos(x)}$ is separable as it can be written as

$$\begin{aligned}y' &= c_2 e^{-\frac{y^2}{2}} e^{\cos(x)} \\ &= f(x)g(y)\end{aligned}$$

Where

$$\begin{aligned}f(x) &= e^{\cos(x)} c_2 \\ g(y) &= e^{-\frac{y^2}{2}}\end{aligned}$$

Integrating gives

$$\begin{aligned}\int \frac{1}{g(y)} dy &= \int f(x) dx \\ \int e^{\frac{y^2}{2}} dy &= \int e^{\cos(x)} c_2 dx \\ -\frac{i\sqrt{\pi}\sqrt{2}\operatorname{erf}\left(\frac{i\sqrt{2}y}{2}\right)}{2} &= \int e^{\cos(x)} c_2 dx + 2c_3\end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$-\frac{i\sqrt{\pi}\sqrt{2}\operatorname{erf}\left(\frac{i\sqrt{2}y}{2}\right)}{2} = \int e^{\cos(x)} c_2 dx + 2c_3$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.011 (sec)

Leaf size : 35

```
dsolve(diff(diff(y(x),x),x)+sin(x)*diff(y(x),x)+y(x)*diff(y(x),x)^2 = 0,
y(x),singsol=all)
```

$$y = -i \operatorname{RootOf}\left(i\sqrt{2}c_1\left(\int e^{\cos(x)} dx\right) + i\sqrt{2}c_2 - \operatorname{erf}(_Z)\sqrt{\pi}\right)\sqrt{2}$$

Mathematica DSolve solution

Solving time : 0.433 (sec)

Leaf size : 47

```
DSolve[{D[y[x], {x, 2}] + Sin[x]*D[y[x], x] + y[x]*(D[y[x], x])^2 == 0, {}},  
y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -i\sqrt{2}\operatorname{erf}^{-1}\left(i\sqrt{\frac{2}{\pi}}\left(\int_1^x -e^{\cos(K[1])}c_1 dK[1] + c_2\right)\right)$$

2.2.3 problem 3

Maple step by step solution	514
Maple trace	514
Maple dsolve solution	514
Mathematica DSolve solution	514

Internal problem ID [8802]

Book : Second order enumerated odes

Section : section 2

Problem number : 3

Date solved : Wednesday, December 18, 2024 at 02:16:36 AM

CAS classification :

[_Liouville, [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + (1 - x)y' + y^2y'^2 = 0$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.015 (sec)

Leaf size : 61

```
dsolve(diff(diff(y(x),x),x)+(1-x)*diff(y(x),x)+y(x)^2*diff(y(x),x)^2 = 0,
y(x),singsol=all)
```

$$c_1 \operatorname{erf}\left(\frac{i\sqrt{2}(x-1)}{2}\right) - c_2 + \frac{2 \cdot 3^{5/6} y \pi}{9 \Gamma\left(\frac{2}{3}\right) (-y^3)^{1/3}} - \frac{y \Gamma\left(\frac{1}{3}, -\frac{y^3}{3}\right) 3^{1/3}}{3 (-y^3)^{1/3}} = 0$$

Mathematica DSolve solution

Solving time : 0.512 (sec)

Leaf size : 67

```
DSolve[{D[y[x],{x,2}]+(1-x)*D[y[x],x]+y[x]^2*(D[y[x],x])^2==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \operatorname{InverseFunction}\left[-\frac{\#1 \Gamma\left(\frac{1}{3}, -\frac{\#1^3}{3}\right)}{3^{2/3} \sqrt[3]{-\#1^3}} \&\right] \left[c_2 - \sqrt{\frac{\pi}{2e}} c_1 \operatorname{erfi}\left(\frac{x-1}{\sqrt{2}}\right) \right]$$

2.2.4 problem 4

Maple step by step solution	515
Maple trace	515
Maple dsolve solution	515
Mathematica DSolve solution	515

Internal problem ID [8803]

Book : Second order enumerated odes

Section : section 2

Problem number : 4

Date solved : Wednesday, December 18, 2024 at 02:16:36 AM

CAS classification :

[_Liouville, [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + (\sin(x) + 2x)y' + \cos(y)yy'^2 = 0$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.009 (sec)

Leaf size : 34

```
dsolve(diff(diff(y(x),x),x)+(sin(x)+2*x)*diff(y(x),x)+cos(y(x))*y(x)*diff(y(x),x)^2 =
y(x),singsol=all)
```

$$\int^y e^{\cos(a)+a\sin(a)} da - c_1 \left(\int e^{-x^2+\cos(x)} dx \right) - c_2 = 0$$

Mathematica DSolve solution

Solving time : 1.456 (sec)

Leaf size : 53

```
DSolve[{D[y[x],{x,2}]+(Sin[x]+2*x)*D[y[x],x]+Cos[y[x]]*y[x]*(D[y[x],x])^2==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} e^{\cos(K[1])+K[1]\sin(K[1])} dK[1] \& \right] \left[\int_1^x -e^{\cos(K[2])-K[2]^2} c_1 dK[2] + c_2 \right]$$

2.2.5 problem 5

Solved as second order missing x ode	516
Solved as second order can be made integrable	518
Maple step by step solution	520
Maple trace	520
Maple dsolve solution	521
Mathematica DSolve solution	521

Internal problem ID [8804]

Book : Second order enumerated odes

Section : section 2

Problem number : 5

Date solved : Thursday, December 12, 2024 at 09:49:11 AM

CAS classification :

[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]

Solve

$$y''y' + y^2 = 0$$

Solved as second order missing x ode

Time used: 1.107 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y)^2 \left(\frac{d}{dy} p(y) \right) + y^2 = 0$$

Which is now solved as first order ode for $p(y)$.

The ode $p' = -\frac{y^2}{p^2}$ is separable as it can be written as

$$\begin{aligned} p' &= -\frac{y^2}{p^2} \\ &= f(y)g(p) \end{aligned}$$

Where

$$\begin{aligned} f(y) &= -y^2 \\ g(p) &= \frac{1}{p^2} \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(p)} dp &= \int f(y) dy \\ \int p^2 dp &= \int -y^2 dy \\ \frac{p^3}{3} &= -\frac{y^3}{3} + c_1 \end{aligned}$$

Solving for p gives

$$p = (-y^3 + 3c_1)^{1/3}$$

$$p = -\frac{(-y^3 + 3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2}$$

$$p = -\frac{(-y^3 + 3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = (-y^3 + 3c_1)^{1/3}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{1}{(-\tau^3 + 3c_1)^{1/3}} d\tau = x + c_2$$

Singular solutions are found by solving

$$(-y^3 + 3c_1)^{1/3} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 3^{1/3}c_1^{1/3}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2}$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{(-y^3 + 3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{1}{-\frac{(-\tau^3 + 3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-\tau^3 + 3c_1)^{1/3}}{2}} d\tau = x + c_3$$

Singular solutions are found by solving

$$-\frac{(-y^3 + 3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 3^{1/3}c_1^{1/3}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2}$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{(-y^3 + 3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{1}{-\frac{(-\tau^3+3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-\tau^3+3c_1)^{1/3}}{2}} d\tau = x + c_4$$

Singular solutions are found by solving

$$-\frac{(-y^3 + 3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$\begin{aligned} y &= 3^{1/3}c_1^{1/3} \\ y &= -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2} \\ y &= -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2} \end{aligned}$$

Will add steps showing solving for IC soon.

The solution

$$y = 3^{1/3}c_1^{1/3}$$

was found not to satisfy the ode or the IC. Hence it is removed. The solution

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2}$$

was found not to satisfy the ode or the IC. Hence it is removed. The solution

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2}$$

was found not to satisfy the ode or the IC. Hence it is removed.

Summary of solutions found

$$\int^y \frac{1}{-\frac{(-\tau^3+3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-\tau^3+3c_1)^{1/3}}{2}} d\tau = x + c_3$$

$$\int^y \frac{1}{-\frac{(-\tau^3+3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-\tau^3+3c_1)^{1/3}}{2}} d\tau = x + c_4$$

$$\int^y \frac{1}{(-\tau^3 + 3c_1)^{1/3}} d\tau = x + c_2$$

Solved as second order can be made integrable

Time used: 0.569 (sec)

Multiplying the ode by y' gives

$$y^2y' + y'^2y'' = 0$$

Integrating the above w.r.t x gives

$$\begin{aligned} \int (y^2y' + y'^2y'') dx &= 0 \\ \frac{y^3}{3} + \frac{y'^3}{3} &= c_1 \end{aligned}$$

Which is now solved for y . Solving for the derivative gives these ODE's to solve

$$y' = (-y^3 + 3c_1)^{1/3} \quad (1)$$

$$y' = -\frac{(-y^3 + 3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2} \quad (2)$$

$$y' = -\frac{(-y^3 + 3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2} \quad (3)$$

Now each of the above is solved separately.

Solving Eq. (1)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{1}{(-\tau^3 + 3c_1)^{1/3}} d\tau = x + c_2$$

Singular solutions are found by solving

$$(-y^3 + 3c_1)^{1/3} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 3^{1/3}c_1^{1/3}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2}$$

Solving Eq. (2)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{1}{-\frac{(-\tau^3+3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-\tau^3+3c_1)^{1/3}}{2}} d\tau = x + c_3$$

Singular solutions are found by solving

$$-\frac{(-y^3 + 3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 3^{1/3}c_1^{1/3}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2}$$

Solving Eq. (3)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{1}{-\frac{(-\tau^3+3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-\tau^3+3c_1)^{1/3}}{2}} d\tau = x + c_4$$

Singular solutions are found by solving

$$-\frac{(-y^3 + 3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-y^3 + 3c_1)^{1/3}}{2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 3^{1/3}c_1^{1/3}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2}$$

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2}$$

Will add steps showing solving for IC soon.

The solution

$$y = 3^{1/3}c_1^{1/3}$$

was found not to satisfy the ode or the IC. Hence it is removed. The solution

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} - \frac{i3^{5/6}c_1^{1/3}}{2}$$

was found not to satisfy the ode or the IC. Hence it is removed. The solution

$$y = -\frac{3^{1/3}c_1^{1/3}}{2} + \frac{i3^{5/6}c_1^{1/3}}{2}$$

was found not to satisfy the ode or the IC. Hence it is removed.

Summary of solutions found

$$\int^y \frac{1}{-\frac{(-\tau^3+3c_1)^{1/3}}{2} + \frac{i\sqrt{3}(-\tau^3+3c_1)^{1/3}}{2}} d\tau = x + c_4$$

$$\int^y \frac{1}{-\frac{(-\tau^3+3c_1)^{1/3}}{2} - \frac{i\sqrt{3}(-\tau^3+3c_1)^{1/3}}{2}} d\tau = x + c_3$$

$$\int^y \frac{1}{(-\tau^3 + 3c_1)^{1/3}} d\tau = x + c_2$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
Try integration with the canonical coordinates of the symmetry [0, y]
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(_b(_a)^3+1)/_b(_a), _b(_a), expli
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[1, 0]

```


Maple dsolve solution

Solving time : 0.087 (sec)

Leaf size : 61

```
dsolve(diff(y(x),x)*diff(diff(y(x),x),x)+y(x)^2 = 0,
        y(x),singsol=all)
```

$$y = 0$$

$$y = e^{\frac{\sqrt{3} \left(\int \tan \left(\text{RootOf} \left(-\sqrt{3} \ln \left(\cos \left(_Z \right)^2 \right) - 2\sqrt{3} \ln \left(\tan \left(_Z \right) + \sqrt{3} \right) + 6\sqrt{3} c_1 + 6\sqrt{3} x + 6_Z \right) dx \right) + c_2 + \frac{x}{2}}{2}}$$

Mathematica DSolve solution

Solving time : 1.855 (sec)

Leaf size : 180

```
DSolve[{D[y[x],{x,2}]*D[y[x],x]+y[x]^2==0,{x]},
        y[x],x,IncludeSingularSolutions->True]
```

 $y(x)$

→

$$\sqrt[3]{1 + \text{InverseFunction} \left[\frac{1}{6} \log(\#1^2 - \#1 + 1) + \frac{\arctan\left(\frac{2\#1-1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3} \log(\#1 + 1) \& \right] [-x + c_1]^3 \sqrt[3]{\text{Inv}}$$

2.2.6 problem 6

Solved as second order missing x ode	522
Solved as second order can be made integrable	524
Maple step by step solution	526
Maple trace	526
Maple dsolve solution	526
Mathematica DSolve solution	527

Internal problem ID [8805]

Book : Second order enumerated odes

Section : section 2

Problem number : 6

Date solved : Thursday, December 12, 2024 at 09:49:13 AM

CAS classification :

[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]

Solve

$$y''y' + y^n = 0$$

Solved as second order missing x ode

Time used: 1.533 (sec)

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dp}{dy} \frac{dy}{dx} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y)^2 \left(\frac{d}{dy} p(y) \right) + y^n = 0$$

Which is now solved as first order ode for $p(y)$.

The ode $p' = -\frac{y^n}{p^2}$ is separable as it can be written as

$$\begin{aligned} p' &= -\frac{y^n}{p^2} \\ &= f(y)g(p) \end{aligned}$$

Where

$$\begin{aligned} f(y) &= -y^n \\ g(p) &= \frac{1}{p^2} \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(p)} dp &= \int f(y) dy \\ \int p^2 dp &= \int -y^n dy \\ \frac{p^3}{3} &= -\frac{y^{n+1}}{n+1} + c_1 \end{aligned}$$

Solving for p gives

$$p = \frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{n+1}$$

$$p = -\frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2(n+1)} - \frac{i\sqrt{3}((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2(n+1)}$$

$$p = -\frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2(n+1)} + \frac{i\sqrt{3}((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2n+2}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{n+1}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{n+1}{((3c_1n - 3\tau^{n+1} + 3c_1)(n+1)^2)^{1/3}} d\tau = x + c_2$$

Singular solutions are found by solving

$$\frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{n+1} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{\frac{\ln(c_1n+c_1)}{n+1}}$$

For solution (2) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2(n+1)} - \frac{i\sqrt{3}((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2(n+1)}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y -\frac{2(n+1)}{((3c_1n - 3\tau^{n+1} + 3c_1)(n+1)^2)^{1/3} (1+i\sqrt{3})} d\tau = x + c_3$$

Singular solutions are found by solving

$$-\frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3} (1+i\sqrt{3})}{2(n+1)} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{\frac{\ln(c_1n+c_1)}{n+1}}$$

For solution (3) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2(n+1)} + \frac{i\sqrt{3}((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3}}{2n+2}$$

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{2n+2}{((3c_1n - 3\tau^{n+1} + 3c_1)(n+1)^2)^{1/3} (i\sqrt{3}-1)} d\tau = x + c_4$$

Singular solutions are found by solving

$$\frac{((3c_1n - 3y^{n+1} + 3c_1)(n+1)^2)^{1/3} (i\sqrt{3}-1)}{2n+2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{\frac{\ln(c_1n+c_1)}{n+1}}$$

Will add steps showing solving for IC soon.

The solution

$$y = e^{\frac{\ln(c_1n+c_1)}{n+1}}$$

was found not to satisfy the ode or the IC. Hence it is removed.

Summary of solutions found

$$\int^y \frac{n+1}{((3c_1n - 3\tau^{n+1} + 3c_1)(n+1)^2)^{1/3}} d\tau = x + c_2$$

$$\int^y \frac{2n+2}{((3c_1n - 3\tau^{n+1} + 3c_1)(n+1)^2)^{1/3} (i\sqrt{3}-1)} d\tau = x + c_4$$

$$\int^y \frac{2(n+1)}{((3c_1n - 3\tau^{n+1} + 3c_1)(n+1)^2)^{1/3} (1+i\sqrt{3})} d\tau = x + c_3$$

Solved as second order can be made integrable

Time used: 1.503 (sec)

Multiplying the ode by y' gives

$$y'^2 y'' + y^{n-1} y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y'^2 y'' + y^{n-1} y' y) dx = 0$$

$$\frac{y'^3}{3} + \frac{y^2 e^{(n-1)\ln(y)}}{n+1} = c_1$$

Which is now solved for y . Solving for the derivative gives these ODE's to solve

$$y' = \frac{((-3y^2 e^{(n-1)\ln(y)} + 3c_1n + 3c_1)(n+1)^2)^{1/3}}{n+1} \quad (1)$$

$$y' = -\frac{((-3y^2 e^{(n-1)\ln(y)} + 3c_1n + 3c_1)(n+1)^2)^{1/3}}{2(n+1)} \quad (2)$$

$$-\frac{i\sqrt{3}((-3y^2 e^{(n-1)\ln(y)} + 3c_1n + 3c_1)(n+1)^2)^{1/3}}{2(n+1)}$$

$$y' = -\frac{((-3y^2 e^{(n-1)\ln(y)} + 3c_1n + 3c_1)(n+1)^2)^{1/3}}{2(n+1)} \quad (3)$$

$$+\frac{i\sqrt{3}((-3y^2 e^{(n-1)\ln(y)} + 3c_1n + 3c_1)(n+1)^2)^{1/3}}{2n+2}$$

Now each of the above is solved separately.

Solving Eq. (1)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{n+1}{((-3\tau^2 e^{(n-1)\ln(\tau)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3}} d\tau = x + c_2$$

Singular solutions are found by solving

$$\frac{((-3y^2 e^{(n-1)\ln(y)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3}}{n+1} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{\frac{\ln(c_1(n+1))}{n+1}}$$

Solving Eq. (2)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y -\frac{2(n+1)}{((-3\tau^2 e^{(n-1)\ln(\tau)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3} (1+i\sqrt{3})} d\tau = x + c_3$$

Singular solutions are found by solving

$$-\frac{((-3y^2 e^{(n-1)\ln(y)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3} (1+i\sqrt{3})}{2(n+1)} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{\frac{\ln(c_1(n+1))}{n+1}}$$

Solving Eq. (3)

Unable to integrate (or integral too complicated), and since no initial conditions are given, then the result can be written as

$$\int^y \frac{2n+2}{((-3\tau^2 e^{(n-1)\ln(\tau)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3} (i\sqrt{3}-1)} d\tau = x + c_4$$

Singular solutions are found by solving

$$\frac{((-3y^2 e^{(n-1)\ln(y)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3} (i\sqrt{3}-1)}{2n+2} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = e^{\frac{\ln(c_1(n+1))}{n+1}}$$

Will add steps showing solving for IC soon.

The solution

$$y = e^{\frac{\ln(c_1(n+1))}{n+1}}$$

was found not to satisfy the ode or the IC. Hence it is removed.

Summary of solutions found

$$\int^y \frac{n+1}{((-3\tau^2 e^{(n-1)\ln(\tau)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3}} d\tau = x + c_2$$

$$\int^y \frac{2n+2}{((-3\tau^2 e^{(n-1)\ln(\tau)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3} (i\sqrt{3}-1)} d\tau = x + c_4$$

$$\int^y \frac{2(n+1)}{((-3\tau^2 e^{(n-1)\ln(\tau)} + 3c_1 n + 3c_1)(n+1)^2)^{1/3} (1+i\sqrt{3})} d\tau = x + c_3$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+_a^n/_b(_a) = 0, _b(_a), HIN
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[-3/(n-2)*_a, -_b*(n+1)/

```

Maple dsolve solution

Solving time : 0.044 (sec)

Leaf size : 169

```

dsolve(diff(y(x),x)*diff(diff(y(x),x),x)+y(x)^n = 0,
y(x),singsol=all)

```

$$\frac{(-2n-2) \left(\int^y \frac{1}{(-3a^{n+1}-c_1)(n+1)^2)^{1/3}} da \right) - (x+c_2)(1+i\sqrt{3})}{1+i\sqrt{3}} = 0$$

$$\frac{2i(n+1) \left(\int^y \frac{1}{(-3a^{n+1}-c_1)(n+1)^2)^{1/3}} da \right) + (x+c_2)(\sqrt{3}+i)}{\sqrt{3}+i} = 0$$

$$\left(\int^y \frac{1}{(-3a^{n+1}-c_1)(n+1)^2)^{1/3}} da \right) n + \int^y \frac{1}{(-3a^{n+1}-c_1)(n+1)^2)^{1/3}} da - c_2 - x = 0$$

Mathematica DSolve solution

Solving time : 3.397 (sec)

Leaf size : 910

```
DSolve[{D[y[x], {x, 2}] * D[y[x], x] + y[x]^n == 0, {}},
        y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\#1 \sqrt[3]{n+1} \sqrt[3]{1 - \frac{\#1^{n+1}}{c_1(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)c_1} \right)}{\sqrt[3]{-3\#1^{n+1} + 3c_1(n+1)}} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{(-1)^{2/3} \#1 \sqrt[3]{n+1} \sqrt[3]{1 - \frac{\#1^{n+1}}{c_1(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)c_1} \right)}{\sqrt[3]{-3\#1^{n+1} + 3c_1(n+1)}} \& \right] + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\sqrt[3]{-\frac{1}{3}\#1\sqrt[3]{n+1}} \sqrt[3]{1 - \frac{\#1^{n+1}}{c_1(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)c_1} \right)}{\sqrt[3]{-\#1^{n+1} + c_1(n+1)}} \& \right] + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\#1 \sqrt[3]{n+1} \sqrt[3]{1 - \frac{\#1^{n+1}}{(-c_1)(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)(-c_1)} \right)}{\sqrt[3]{-3\#1^{n+1} + 3(-c_1)(n+1)}} \& \right] + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{(-1)^{2/3} \#1 \sqrt[3]{n+1} \sqrt[3]{1 - \frac{\#1^{n+1}}{(-c_1)(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)(-c_1)} \right)}{\sqrt[3]{-3\#1^{n+1} + 3(-c_1)(n+1)}} \& \right] + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\sqrt[3]{-\frac{1}{3}\#1\sqrt[3]{n+1}} \sqrt[3]{1 - \frac{\#1^{n+1}}{(-c_1)(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)(-c_1)} \right)}{\sqrt[3]{-\#1^{n+1} + (-c_1)(n+1)}} \& \right] + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{\#1 \sqrt[3]{n+1} \sqrt[3]{1 - \frac{\#1^{n+1}}{c_1(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)c_1} \right)}{\sqrt[3]{-3\#1^{n+1} + 3c_1(n+1)}} \& \right] [x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction} \left[\frac{(-1)^{2/3} \#1 \sqrt[3]{n+1} \sqrt[3]{1 - \frac{\#1^{n+1}}{c_1(n+1)}} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{1}{n+1}, 1 + \frac{1}{n+1}, \frac{\#1^{n+1}}{(n+1)c_1} \right)}{\sqrt[3]{-3\#1^{n+1} + 3c_1(n+1)}} \& \right] + c_2]$$

2.2.7 problem 8

Solved as first order homogeneous class C ode	528
Solved using Lie symmetry for first order ode	529
Maple step by step solution	533
Maple trace	533
Maple dsolve solution	533
Mathematica DSolve solution	534

Internal problem ID [8806]

Book : Second order enumerated odes

Section : section 2

Problem number : 8

Date solved : Tuesday, December 17, 2024 at 01:01:59 PM

CAS classification : [[_homogeneous, 'class C'], _dAlembert]

Solve

$$y' = (x + y)^4$$

Solved as first order homogeneous class C ode

Time used: 0.497 (sec)

Let

$$z = x + y \tag{1}$$

Then

$$z'(x) = 1 + y'$$

Therefore

$$y' = z'(x) - 1$$

Hence the given ode can now be written as

$$z'(x) - 1 = z^4$$

This is separable first order ode. Integrating

$$\int dx = \int \frac{1}{z^4 + 1} dz$$

$$x + c_1 = \frac{\sqrt{2} \left(\ln \left(\frac{z^2 + z\sqrt{2} + 1}{z^2 - z\sqrt{2} + 1} \right) + 2 \arctan(z\sqrt{2} + 1) + 2 \arctan(z\sqrt{2} - 1) \right)}{8}$$

Replacing z back by its value from (1) then the above gives the solution as




Figure 2.158: Slope field plot
 $y' = (x + y)^4$

Summary of solutions found

$$\frac{\sqrt{2} \left(\ln \left(\frac{(x+y)^2 + (x+y)\sqrt{2}+1}{(x+y)^2 - (x+y)\sqrt{2}+1} \right) + 2 \arctan \left((x+y)\sqrt{2}+1 \right) + 2 \arctan \left((x+y)\sqrt{2}-1 \right) \right)}{8} = x + c_1$$

Solved using Lie symmetry for first order ode

Time used: 1.251 (sec)

Writing the ode as

$$y' = (x + y)^4$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + (x + y)^4 (b_3 - a_2) - (x + y)^8 a_3 - 4(x + y)^3 (xa_2 + ya_3 + a_1) - 4(x + y)^3 (xb_2 + yb_3 + b_1) = 0 \quad (5\text{E})$$

Putting the above in normal form gives

$$\begin{aligned} & -x^8 a_3 - 8x^7 y a_3 - 28x^6 y^2 a_3 - 56x^5 y^3 a_3 - 70x^4 y^4 a_3 - 56x^3 y^5 a_3 - 28x^2 y^6 a_3 - 8x y^7 a_3 \\ & - y^8 a_3 - 5x^4 a_2 - 4x^4 b_2 + x^4 b_3 - 16x^3 y a_2 - 4x^3 y a_3 - 12x^3 y b_2 - 18x^2 y^2 a_2 - 12x^2 y^2 a_3 \\ & - 12x^2 y^2 b_2 - 6x^2 y^2 b_3 - 8x y^3 a_2 - 12x y^3 a_3 - 4x y^3 b_2 - 8x y^3 b_3 - y^4 a_2 - 4y^4 a_3 - 3y^4 b_3 \\ & - 4x^3 a_1 - 4x^3 b_1 - 12x^2 y a_1 - 12x^2 y b_1 - 12x y^2 a_1 - 12x y^2 b_1 - 4y^3 a_1 - 4y^3 b_1 + b_2 = 0 \end{aligned}$$

Setting the numerator to zero gives

$$\begin{aligned}
 & -x^8 a_3 - 8x^7 y a_3 - 28x^6 y^2 a_3 - 56x^5 y^3 a_3 - 70x^4 y^4 a_3 - 56x^3 y^5 a_3 \\
 & - 28x^2 y^6 a_3 - 8x y^7 a_3 - y^8 a_3 - 5x^4 a_2 - 4x^4 b_2 + x^4 b_3 - 16x^3 y a_2 - 4x^3 y a_3 \\
 & - 12x^3 y b_2 - 18x^2 y^2 a_2 - 12x^2 y^2 a_3 - 12x^2 y^2 b_2 - 6x^2 y^2 b_3 - 8x y^3 a_2 \\
 & - 12x y^3 a_3 - 4x y^3 b_2 - 8x y^3 b_3 - y^4 a_2 - 4y^4 a_3 - 3y^4 b_3 - 4x^3 a_1 - 4x^3 b_1 \\
 & - 12x^2 y a_1 - 12x^2 y b_1 - 12x y^2 a_1 - 12x y^2 b_1 - 4y^3 a_1 - 4y^3 b_1 + b_2 = 0
 \end{aligned} \tag{6E}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
 & -a_3 v_1^8 - 8a_3 v_1^7 v_2 - 28a_3 v_1^6 v_2^2 - 56a_3 v_1^5 v_2^3 - 70a_3 v_1^4 v_2^4 - 56a_3 v_1^3 v_2^5 - 28a_3 v_1^2 v_2^6 \\
 & - 8a_3 v_1 v_2^7 - a_3 v_2^8 - 5a_2 v_1^4 - 16a_2 v_1^2 v_2^2 - 18a_2 v_1^2 v_2^2 - 8a_2 v_1 v_2^3 - a_2 v_2^4 \\
 & - 4a_3 v_1^3 v_2 - 12a_3 v_1^2 v_2^2 - 12a_3 v_1 v_2^3 - 4a_3 v_2^4 - 4b_2 v_1^4 - 12b_2 v_1^3 v_2 - 12b_2 v_1^2 v_2^2 \\
 & - 4b_2 v_1 v_2^3 + b_3 v_1^4 - 6b_3 v_1^2 v_2^2 - 8b_3 v_1 v_2^3 - 3b_3 v_2^4 - 4a_1 v_1^3 - 12a_1 v_1^2 v_2 \\
 & - 12a_1 v_1 v_2^2 - 4a_1 v_2^3 - 4b_1 v_1^3 - 12b_1 v_1^2 v_2 - 12b_1 v_1 v_2^2 - 4b_1 v_2^3 + b_2 = 0
 \end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned}
 & -a_3 v_1^8 - 8a_3 v_1^7 v_2 - 28a_3 v_1^6 v_2^2 - 56a_3 v_1^5 v_2^3 - 70a_3 v_1^4 v_2^4 + (-5a_2 - 4b_2 + b_3) v_1^4 \\
 & - 56a_3 v_1^3 v_2^5 + (-16a_2 - 4a_3 - 12b_2) v_1^3 v_2 + (-4a_1 - 4b_1) v_1^3 \\
 & - 28a_3 v_1^2 v_2^6 + (-18a_2 - 12a_3 - 12b_2 - 6b_3) v_1^2 v_2^2 + (-12a_1 - 12b_1) v_1^2 v_2 \\
 & - 8a_3 v_1 v_2^7 + (-8a_2 - 12a_3 - 4b_2 - 8b_3) v_1 v_2^3 + (-12a_1 - 12b_1) v_1 v_2^2 \\
 & - a_3 v_2^8 + (-a_2 - 4a_3 - 3b_3) v_2^4 + (-4a_1 - 4b_1) v_2^3 + b_2 = 0
 \end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 b_2 &= 0 \\
 -70a_3 &= 0 \\
 -56a_3 &= 0 \\
 -28a_3 &= 0 \\
 -8a_3 &= 0 \\
 -a_3 &= 0 \\
 -12a_1 - 12b_1 &= 0 \\
 -4a_1 - 4b_1 &= 0 \\
 -16a_2 - 4a_3 - 12b_2 &= 0 \\
 -5a_2 - 4b_2 + b_3 &= 0 \\
 -a_2 - 4a_3 - 3b_3 &= 0 \\
 -18a_2 - 12a_3 - 12b_2 - 6b_3 &= 0 \\
 -8a_2 - 12a_3 - 4b_2 - 8b_3 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= -b_1 \\ a_2 &= 0 \\ a_3 &= 0 \\ b_1 &= b_1 \\ b_2 &= 0 \\ b_3 &= 0 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= -1 \\ \eta &= 1 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned} \frac{dy}{dx} &= \frac{\eta}{\xi} \\ &= \frac{1}{-1} \\ &= -1 \end{aligned}$$

This is easily solved to give

$$y = -x + c_1$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = x + y$$

And S is found from

$$\begin{aligned} dS &= \frac{dx}{\xi} \\ &= \frac{dx}{-1} \end{aligned}$$

Integrating gives

$$\begin{aligned} S &= \int \frac{dx}{-1} \\ &= -x \end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = (x + y)^4$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 1 \\ S_x &= -1 \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{1 + (x + y)^4} \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R^4 + 1}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S .



Since the ode has the form $\frac{d}{dR}S(R) = f(R)$, then we only need to integrate $f(R)$.

$$\begin{aligned} \int dS &= \int -\frac{1}{R^4 + 1} dR \\ S(R) &= -\frac{\sqrt{2} \left(\ln \left(\frac{R^2 + R\sqrt{2} + 1}{R^2 - R\sqrt{2} + 1} \right) + 2 \arctan (R\sqrt{2} + 1) + 2 \arctan (R\sqrt{2} - 1) \right)}{8} + c_2 \end{aligned}$$

To complete the solution, we just need to transform the above back to x, y coordinates. This results in

$$-x = -\frac{\sqrt{2} \left(\ln \left(\frac{(x+y)^2 + (x+y)\sqrt{2} + 1}{(x+y)^2 - (x+y)\sqrt{2} + 1} \right) + 2 \arctan ((x + y) \sqrt{2} + 1) + 2 \arctan ((x + y) \sqrt{2} - 1) \right)}{8} + c_2$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = (x + y)^4$ 	$\begin{aligned} R &= x + y \\ S &= -x \end{aligned}$	$\frac{dS}{dR} = -\frac{1}{R^4 + 1}$ 




Figure 2.159: Slope field plot
 $y' = (x + y)^4$

Summary of solutions found

$$\begin{aligned}
 & -x \\
 & = -\frac{\sqrt{2} \left(\ln \left(\frac{(x+y)^2 + (x+y)\sqrt{2}+1}{(x+y)^2 - (x+y)\sqrt{2}+1} \right) + 2 \arctan \left((x+y)\sqrt{2}+1 \right) + 2 \arctan \left((x+y)\sqrt{2}-1 \right) \right)}{8} \\
 & + c_2
 \end{aligned}$$

Maple step by step solution

Let's solve

$$\frac{d}{dx}y(x) = (x + y(x))^4$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx}y(x)$$

- Solve for the highest derivative

$$\frac{d}{dx}y(x) = (x + y(x))^4$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous C
1st order, trying the canonical coordinates of the invariance group
<- 1st order, canonical coordinates successful
<- homogeneous successful`

```

Maple dsolve solution

Solving time : 0.912 (sec)

Leaf size : 882

```

dsolve(diff(y(x),x) = (x+y(x))^4,
       y(x),singsol=all)

```

Expression too large to display

Mathematica DSolve solution

Solving time : 0.168 (sec)

Leaf size : 88

```
DSolve[{D[y[x],x] == (x + y[x])^4,{}},
  y[x],x,IncludeSingularSolutions->True]
```

$$\text{Solve}\left[\frac{1}{4}\text{RootSum}\left[\#1^4 + 4\#1^3y(x) + 6\#1^2y(x)^2 + 4\#1y(x)^3 + y(x)^4\right.\right. \\ \left.\left.+ 1\&, \frac{\log(x - \#1)}{\#1^3 + 3\#1^2y(x) + 3\#1y(x)^2 + y(x)^3}\&\right] - x = c_1, y(x)\right]$$

2.2.8 problem 9

Maple step by step solution	535
Maple trace	535
Maple dsolve solution	535
Mathematica DSolve solution	535

Internal problem ID [8807]

Book : Second order enumerated odes**Section** : section 2**Problem number** : 9**Date solved** : Wednesday, December 18, 2024 at 02:16:37 AM**CAS classification** :

[_Liouville, [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + (3 + x)y' + (3 + y^2)y'^2 = 0$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`

```

Maple dsolve solution

Solving time : 0.014 (sec)

Leaf size : 32

```

dsolve(diff(diff(y(x),x),x)+(x+3)*diff(y(x),x)+(y(x)^2+3)*diff(y(x),x)^2 = 0,
        y(x),singsol=all)

```

$$c_1 \operatorname{erf}\left(\frac{\sqrt{2}(x+3)}{2}\right) - c_2 + \int^y e^{\frac{-a(-a^2+9)}{3}} d_a = 0$$

Mathematica DSolve solution

Solving time : 0.609 (sec)

Leaf size : 61

```

DSolve[{D[y[x],{x,2}]+(3+x)*D[y[x],x]+(3+y[x]^2)*(D[y[x],x])^2==0,{x},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \operatorname{InverseFunction}\left[\int_1^{\#1} e^{\frac{K[1]^3}{3}+3K[1]} dK[1] \& \right] \left[c_2 - e^{9/2} \sqrt{\frac{\pi}{2}} c_1 \operatorname{erf}\left(\frac{x+3}{\sqrt{2}}\right) \right]$$

2.2.9 problem 10

Solved as second nonlinear ode solved by Mainardi Liouville method	536
Maple step by step solution	537
Maple trace	537
Maple dsolve solution	538
Mathematica DSolve solution	538

Internal problem ID [8808]

Book : Second order enumerated odes

Section : section 2

Problem number : 10

Date solved : Thursday, December 12, 2024 at 09:51:46 AM

CAS classification : [_Liouville, [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + xy' + yy'^2 = 0$$

Solved as second nonlinear ode solved by Mainardi Liouville method

Time used: 0.213 (sec)

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \tag{1A}$$

Where in this problem

$$\begin{aligned} f(x) &= x \\ g(y) &= y \end{aligned}$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \tag{2A}$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \tag{3A}$$

And the last term in Eq (2A) can be written as

$$\begin{aligned} g \frac{dy}{dx} &= \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx} \\ &= \frac{d}{dx} \int g dy \end{aligned} \tag{4A}$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \tag{5A}$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \tag{6A}$$

Where c_2 is a new arbitrary constant. But since $g = y$ and $f = x$, then

$$\begin{aligned}\int -g dy &= \int -y dy \\ &= -\frac{y^2}{2} \\ \int -f dx &= \int -x dx \\ &= -\frac{x^2}{2}\end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = c_2 e^{-\frac{y^2}{2}} e^{-\frac{x^2}{2}}$$

Which is now solved as first order separable ode. The ode $y' = c_2 e^{-\frac{y^2}{2}} e^{-\frac{x^2}{2}}$ is separable as it can be written as

$$\begin{aligned}y' &= c_2 e^{-\frac{y^2}{2}} e^{-\frac{x^2}{2}} \\ &= f(x)g(y)\end{aligned}$$

Where

$$\begin{aligned}f(x) &= e^{-\frac{x^2}{2}} c_2 \\ g(y) &= e^{-\frac{y^2}{2}}\end{aligned}$$

Integrating gives

$$\begin{aligned}\int \frac{1}{g(y)} dy &= \int f(x) dx \\ \int e^{\frac{y^2}{2}} dy &= \int e^{-\frac{x^2}{2}} c_2 dx \\ -\frac{i\sqrt{\pi} \sqrt{2} \operatorname{erf}\left(\frac{i\sqrt{2}y}{2}\right)}{2} &= \frac{c_2 \sqrt{\pi} \sqrt{2} \operatorname{erf}\left(\frac{\sqrt{2}x}{2}\right)}{2} + c_3\end{aligned}$$

Will add steps showing solving for IC soon.

Solving for y from the above solution(s) gives (after possible removing of solutions that do not verify)

$$y = -i \operatorname{RootOf}\left(i\sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{2}x}{2}\right) c_2 + i\sqrt{2} c_3 - \operatorname{erf}(_Z) \sqrt{\pi}\right) \sqrt{2}$$

Summary of solutions found

$$y = -i \operatorname{RootOf}\left(i\sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{2}x}{2}\right) c_2 + i\sqrt{2} c_3 - \operatorname{erf}(_Z) \sqrt{\pi}\right) \sqrt{2}$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.013 (sec)

Leaf size : 37

```
dsolve(diff(diff(y(x),x),x)+diff(y(x),x)*x+y(x)*diff(y(x),x)^2 = 0,
y(x),singsol=all)
```

$$y = -i \operatorname{RootOf} \left(i\sqrt{\pi} \operatorname{erf} \left(\frac{\sqrt{2}x}{2} \right) c_1 + i\sqrt{2} c_2 - \operatorname{erf}(_Z) \sqrt{\pi} \right) \sqrt{2}$$

Mathematica DSolve solution

Solving time : 0.128 (sec)

Leaf size : 44

```
DSolve[{D[y[x],{x,2}]+x*D[y[x],x]+y[x]*(D[y[x],x])^2==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -i\sqrt{2}\operatorname{erf}^{-1} \left(i \left(\sqrt{\frac{2}{\pi}} c_2 - c_1 \operatorname{erf} \left(\frac{x}{\sqrt{2}} \right) \right) \right)$$

2.2.10 problem 11

Solved as second order missing y ode	539
Solved as second nonlinear ode solved by Mainardi Liouville method	541
Maple step by step solution	543
Maple trace	543
Maple dsolve solution	543
Mathematica DSolve solution	543

Internal problem ID [8809]

Book : Second order enumerated odes

Section : section 2

Problem number : 11

Date solved : Thursday, December 12, 2024 at 09:51:47 AM

CAS classification :

[[_2nd_order, _missing_y], _Liouville, [_2nd_order, _reducible, _mu_xy]]

Solve

$$y'' + \sin(x)y' + y'^2 = 0$$

Solved as second order missing y ode

Time used: 0.753 (sec)

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + \sin(x)p(x) + p(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode.

In canonical form, the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= -\sin(x)p - p^2 \end{aligned}$$

This is a Bernoulli ODE.

$$p' = (-\sin(x))p(x) + (-1)p^2 \tag{1}$$

The standard Bernoulli ODE has the form

$$p' = f_0(x)p + f_1(x)p^n \tag{2}$$

Comparing this to (1) shows that

$$\begin{aligned} f_0 &= -\sin(x) \\ f_1 &= -1 \end{aligned}$$

The first step is to divide the above equation by p^n which gives

$$\frac{p'}{p^n} = f_0(x)p^{1-n} + f_1(x) \tag{3}$$

The next step is use the substitution $v = p^{1-n}$ in equation (3) which generates a new ODE in $v(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $p(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= -\sin(x) \\ f_1(x) &= -1 \\ n &= 2 \end{aligned}$$

Dividing both sides of ODE (1) by $p^n = p^2$ gives

$$p' \frac{1}{p^2} = -\frac{\sin(x)}{p} - 1 \quad (4)$$

Let

$$\begin{aligned} v &= p^{1-n} \\ &= \frac{1}{p} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$v' = -\frac{1}{p^2} p' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -v'(x) &= -\sin(x)v(x) - 1 \\ v' &= \sin(x)v + 1 \end{aligned} \quad (7)$$

The above now is a linear ODE in $v(x)$ which is now solved.

In canonical form a linear first order is

$$v'(x) + q(x)v(x) = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\sin(x) \\ p(x) &= 1 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\sin(x) dx} \\ &= e^{\cos(x)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu v) &= \mu \\ \frac{d}{dx}(v e^{\cos(x)}) &= e^{\cos(x)} \\ d(v e^{\cos(x)}) &= e^{\cos(x)} dx \end{aligned}$$

Integrating gives

$$\begin{aligned} v e^{\cos(x)} &= \int e^{\cos(x)} dx \\ &= \int e^{\cos(x)} dx + c_1 \end{aligned}$$

Dividing throughout by the integrating factor $e^{\cos(x)}$ gives the final solution

$$v(x) = e^{-\cos(x)} \left(\int e^{\cos(x)} dx + c_1 \right)$$

The substitution $v = p^{1-n}$ is now used to convert the above solution back to $p(x)$ which results in

$$\frac{1}{p(x)} = e^{-\cos(x)} \left(\int e^{\cos(x)} dx + c_1 \right)$$

Solving for $p(x)$ gives

$$p(x) = \frac{e^{\cos(x)}}{\int e^{\cos(x)} dx + c_1}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{e^{\cos(x)}}{\int e^{\cos(x)} dx + c_1}$$

Since the ode has the form $y' = f(x)$, then we only need to integrate $f(x)$.

$$\int dy = \int \frac{e^{\cos(x)}}{\int e^{\cos(x)} dx + c_1} dx$$

$$y = \ln \left(\int e^{\cos(x)} dx + c_1 \right) + c_2$$

$$y = \int \frac{e^{\cos(x)}}{\int e^{\cos(x)} dx + c_1} dx + c_2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \int \frac{e^{\cos(x)}}{\int e^{\cos(x)} dx + c_1} dx + c_2$$

Solved as second nonlinear ode solved by Mainardi Liouville method

Time used: 0.253 (sec)

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \tag{1A}$$

Where in this problem

$$f(x) = \sin(x)$$

$$g(y) = 1$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \tag{2A}$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \tag{3A}$$

And the last term in Eq (2A) can be written as

$$\begin{aligned} g \frac{dy}{dx} &= \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx} \\ &= \frac{d}{dx} \int g dy \end{aligned} \quad (4A)$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \quad (5A)$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \quad (6A)$$

Where c_2 is a new arbitrary constant. But since $g = 1$ and $f = \sin(x)$, then

$$\begin{aligned} \int -g dy &= \int (-1) dy \\ &= -y \\ \int -f dx &= \int -\sin(x) dx \\ &= \cos(x) \end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = c_2 e^{-y} e^{\cos(x)}$$

Which is now solved as first order separable ode. The ode $y' = c_2 e^{-y} e^{\cos(x)}$ is separable as it can be written as

$$\begin{aligned} y' &= c_2 e^{-y} e^{\cos(x)} \\ &= f(x)g(y) \end{aligned}$$

Where

$$\begin{aligned} f(x) &= e^{\cos(x)} c_2 \\ g(y) &= e^{-y} \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(y)} dy &= \int f(x) dx \\ \int e^y dy &= \int e^{\cos(x)} c_2 dx \\ e^y &= \int e^{\cos(x)} c_2 dx + 2c_3 \end{aligned}$$

Solving for y gives

$$y = \ln \left(c_2 \int e^{\cos(x)} dx + 2c_3 \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \ln \left(c_2 \int e^{\cos(x)} dx + 2c_3 \right)$$

Maple step by step solution**Maple trace**

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.009 (sec)

Leaf size : 14

```
dsolve(diff(diff(y(x),x),x)+sin(x)*diff(y(x),x)+diff(y(x),x)^2 = 0,
        y(x),singsol=all)
```

$$y = \ln \left(c_1 \left(\int e^{\cos(x)} dx \right) + c_2 \right)$$

Mathematica DSolve solution

Solving time : 60.135 (sec)

Leaf size : 43

```
DSolve[{D[y[x],{x,2}]+Sin[x]*D[y[x],x]+(D[y[x],x])^2==0,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \int_1^x \frac{e^{\cos(K[2])}}{c_1 - \int_1^{K[2]} -e^{\cos(K[1])} dK[1]} dK[2] + c_2$$

2.2.11 problem 12

Solved as second nonlinear ode solved by Mainardi Liouville method	544
Maple step by step solution	545
Maple trace	545
Maple dsolve solution	545
Mathematica DSolve solution	546

Internal problem ID [8810]

Book : Second order enumerated odes

Section : section 2

Problem number : 12

Date solved : Thursday, December 12, 2024 at 09:51:48 AM

CAS classification :

[_Liouville, [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]

Solve

$$3y'' + \cos(x)y' + \sin(y)y'^2 = 0$$

Solved as second nonlinear ode solved by Mainardi Liouville method

Time used: 0.339 (sec)

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \tag{1A}$$

Where in this problem

$$f(x) = \frac{\cos(x)}{3}$$

$$g(y) = \frac{\sin(y)}{3}$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \tag{2A}$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \tag{3A}$$

And the last term in Eq (2A) can be written as

$$g \frac{dy}{dx} = \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx}$$

$$= \frac{d}{dx} \int g dy \tag{4A}$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \tag{5A}$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \tag{6A}$$

Where c_2 is a new arbitrary constant. But since $g = \frac{\sin(y)}{3}$ and $f = \frac{\cos(x)}{3}$, then

$$\begin{aligned}\int -g dy &= \int -\frac{\sin(y)}{3} dy \\ &= \frac{\cos(y)}{3} \\ \int -f dx &= \int -\frac{\cos(x)}{3} dx \\ &= -\frac{\sin(x)}{3}\end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = c_2 e^{\frac{\cos(y)}{3}} e^{-\frac{\sin(x)}{3}}$$

Which is now solved as first order separable ode. The ode $y' = c_2 e^{\frac{\cos(y)}{3}} e^{-\frac{\sin(x)}{3}}$ is separable as it can be written as

$$\begin{aligned}y' &= c_2 e^{\frac{\cos(y)}{3}} e^{-\frac{\sin(x)}{3}} \\ &= f(x)g(y)\end{aligned}$$

Where

$$\begin{aligned}f(x) &= e^{-\frac{\sin(x)}{3}} c_2 \\ g(y) &= e^{\frac{\cos(y)}{3}}\end{aligned}$$

Integrating gives

$$\begin{aligned}\int \frac{1}{g(y)} dy &= \int f(x) dx \\ \int e^{-\frac{\cos(y)}{3}} dy &= \int e^{-\frac{\sin(x)}{3}} c_2 dx \\ \int^y e^{-\frac{\cos(\tau)}{3}} d\tau &= \int e^{-\frac{\sin(x)}{3}} c_2 dx + 2c_3\end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\int^y e^{-\frac{\cos(\tau)}{3}} d\tau = \int e^{-\frac{\sin(x)}{3}} c_2 dx + 2c_3$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.008 (sec)

Leaf size : 27

```
dsolve(3*diff(diff(y(x),x),x)+cos(x)*diff(y(x),x)+sin(y(x))*diff(y(x),x)^2 = 0,
y(x),singsol=all)
```

$$\int^y e^{-\frac{\cos(a)}{3}} da - c_1 \left(\int e^{-\frac{\sin(x)}{3}} dx \right) - c_2 = 0$$

Mathematica DSolve solution

Solving time : 0.855 (sec)

Leaf size : 47

```
DSolve[{3*D[y[x],{x,2}]+Cos[x]*D[y[x],x]+Sin[y[x]]*(D[y[x],x])^2==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} e^{-\frac{1}{3} \cos(K[1])} dK[1] \& \right] \left[\int_1^x -e^{-\frac{1}{3} \sin(K[2])} c_1 dK[2] + c_2 \right]$$

2.2.12 problem 13

Solved as second nonlinear ode solved by Mainardi Liouville method	547
Maple step by step solution	548
Maple trace	548
Maple dsolve solution	549
Mathematica DSolve solution	549

Internal problem ID [8811]

Book : Second order enumerated odes

Section : section 2

Problem number : 13

Date solved : Thursday, December 12, 2024 at 09:51:49 AM

CAS classification :

[_Liouville, [_2nd_order, _with_linear_symmetries], [_2nd_order, _reducible, _mu_x_y1]

Solve

$$10y'' + x^2y' + \frac{3y'^2}{y} = 0$$

Solved as second nonlinear ode solved by Mainardi Liouville method

Time used: 0.214 (sec)

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \quad (1A)$$

Where in this problem

$$f(x) = \frac{x^2}{10}$$

$$g(y) = \frac{3}{10y}$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \quad (2A)$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \quad (3A)$$

And the last term in Eq (2A) can be written as

$$g \frac{dy}{dx} = \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx}$$

$$= \frac{d}{dx} \int g dy \quad (4A)$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \quad (5A)$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \quad (6A)$$

Where c_2 is a new arbitrary constant. But since $g = \frac{3}{10y}$ and $f = \frac{x^2}{10}$, then

$$\begin{aligned} \int -g dy &= \int -\frac{3}{10y} dy \\ &= -\frac{3 \ln(y)}{10} \\ \int -f dx &= \int -\frac{x^2}{10} dx \\ &= -\frac{x^3}{30} \end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = \frac{c_2 e^{-\frac{x^3}{30}}}{y^{3/10}}$$

Which is now solved as first order separable ode. The ode $y' = \frac{c_2 e^{-\frac{x^3}{30}}}{y^{3/10}}$ is separable as it can be written as

$$\begin{aligned} y' &= \frac{c_2 e^{-\frac{x^3}{30}}}{y^{3/10}} \\ &= f(x)g(y) \end{aligned}$$

Where

$$\begin{aligned} f(x) &= c_2 e^{-\frac{x^3}{30}} \\ g(y) &= \frac{1}{y^{3/10}} \end{aligned}$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(y)} dy &= \int f(x) dx \\ \int y^{3/10} dy &= \int c_2 e^{-\frac{x^3}{30}} dx \\ \frac{10y^{13/10}}{13} &= \int c_2 e^{-\frac{x^3}{30}} dx + 2c_3 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\frac{10y^{13/10}}{13} = \int c_2 e^{-\frac{x^3}{30}} dx + 2c_3$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

Maple dsolve solution

Solving time : 0.020 (sec)

Leaf size : 55

```
dsolve(10*diff(diff(y(x),x),x)+diff(y(x),x)*x^2+3*diff(y(x),x)^2/y(x) = 0,
y(x),singsol=all)
```

$$\frac{3 \left(c_1 x \operatorname{WhittakerM} \left(\frac{1}{6}, \frac{2}{3}, \frac{x^3}{30} \right) e^{-\frac{x^3}{60}} 30^{1/6} + \frac{4 \left(c_1 x e^{-\frac{x^3}{30}} + c_2 - \frac{10y^{13/10}}{13} \right) (x^3)^{1/6}}{3} \right)}{4(x^3)^{1/6}} = 0$$

Mathematica DSolve solution

Solving time : 66.802 (sec)

Leaf size : 73

```
DSolve[{10*D[y[x],{x,2}]+x^2*D[y[x],x]+3/y[x]*(D[y[x],x])^2==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_2 \exp \left(\int_1^x \frac{30e^{-\frac{1}{30}K[1]^3} \sqrt[3]{K[1]^3}}{30c_1 \sqrt[3]{K[1]^3} - 13\sqrt[3]{30}\Gamma \left(\frac{1}{3}, \frac{K[1]^3}{30} \right) K[1]} dK[1] \right)$$

2.2.13 problem 14

Maple step by step solution	550
Maple trace	550
Maple dsolve solution	550
Mathematica DSolve solution	550

Internal problem ID [8812]

Book : Second order enumerated odes

Section : section 2

Problem number : 14

Date solved : Wednesday, December 18, 2024 at 02:16:38 AM

CAS classification :

[_Liouville, [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]

Solve

$$10y'' + (e^x + 3x)y' + \frac{3e^y y'^2}{\sin(y)} = 0$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
    
```

Maple dsolve solution

Solving time : 0.010 (sec)

Leaf size : 38

```

dsolve(10*diff(diff(y(x),x),x)+(exp(x)+3*x)*diff(y(x),x)+3/sin(y(x))*exp(y(x))*diff(y(x),x),singsol=all)
    
```

$$\int^y e^{\frac{3(\int \csc(\frac{b}{10})e^{-b}d_b)}{10}} d_b - c_1 \left(\int e^{-\frac{3x^2}{20} - \frac{e^x}{10}} dx \right) - c_2 = 0$$

Mathematica DSolve solution

Solving time : 0.265 (sec)

Leaf size : 90

```

DSolve[{10*D[y[x],{x,2}]+(Exp[x]+3*x)*D[y[x],x]+3/Sin[y[x]]*Exp[y[x]]*(D[y[x],x])^2==0,{}},
y[x],x,IncludeSingularSolutions->True]
    
```

$y(x)$

$$\rightarrow \text{InverseFunction} \left[\int_1^{\#1} \exp \left(\left(-\frac{3}{10} - \frac{3i}{10} \right) e^{(1+i)K[1]} \text{Hypergeometric2F1} \left(\frac{1}{2} - \frac{i}{2}, 1, \frac{3}{2} - \frac{i}{2}, e^{2iK[1]} \right) \right) dK[1] \& - e^{\frac{1}{20}(-3K[2]^2 - 2e^{K[2]})} c_1 dK[2] + c_2 \right]$$

2.2.14 problem 15

Solved as second order ode using Kovacic algorithm	551
Solved as second order ode adjoint method	557
Maple step by step solution	562
Maple trace	562
Maple dsolve solution	562
Mathematica DSolve solution	563

Internal problem ID [8813]

Book : Second order enumerated odes

Section : section 2

Problem number : 15

Date solved : Thursday, December 12, 2024 at 09:51:50 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' - \frac{2y}{x^2} = x e^{-\sqrt{x}}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.194 (sec)

Writing the ode as

$$y'' - \frac{2y}{x^2} = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \end{aligned} \tag{3}$$

$$C = -\frac{2}{x^2}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.68: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{x} + (-)(0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \tag{2A}$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \frac{1}{x} \end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \frac{1}{x} \int \frac{1}{\frac{1}{x^2}} dx \\ &= \frac{1}{x} \left(\frac{x^3}{3}\right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x}\right) + c_2 \left(\frac{1}{x} \left(\frac{x^3}{3}\right)\right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - \frac{2y}{x^2} = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{x} + \frac{c_2 x^2}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x}$$

$$y_2 = \frac{x^2}{3}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & \frac{x^2}{3} \\ \frac{d}{dx} \left(\frac{1}{x} \right) & \frac{d}{dx} \left(\frac{x^2}{3} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & \frac{x^2}{3} \\ -\frac{1}{x^2} & \frac{2x}{3} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x} \right) \left(\frac{2x}{3} \right) - \left(\frac{x^2}{3} \right) \left(-\frac{1}{x^2} \right)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 e^{-\sqrt{x}}}{\frac{3}{1}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^3 e^{-\sqrt{x}}}{3} dx$$

Hence

$$u_1 = \frac{2x^{7/2}e^{-\sqrt{x}}}{3} + \frac{14x^3e^{-\sqrt{x}}}{3} + 28x^{5/2}e^{-\sqrt{x}} + 140x^2e^{-\sqrt{x}} + 560x^{3/2}e^{-\sqrt{x}} + 1680xe^{-\sqrt{x}} + 3360\sqrt{x}e^{-\sqrt{x}} + 3360e^{-\sqrt{x}}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-\sqrt{x}}}{1} dx$$

Which simplifies to

$$u_2 = \int e^{-\sqrt{x}} dx$$

Hence

$$u_2 = -2\sqrt{x}e^{-\sqrt{x}} - 2e^{-\sqrt{x}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\frac{2x^{7/2}e^{-\sqrt{x}}}{3} + \frac{14x^3e^{-\sqrt{x}}}{3} + 28x^{5/2}e^{-\sqrt{x}} + 140x^2e^{-\sqrt{x}} + 560x^{3/2}e^{-\sqrt{x}} + 1680xe^{-\sqrt{x}} + 3360\sqrt{x}e^{-\sqrt{x}} + 3360e^{-\sqrt{x}}}{x} + \frac{x^2(-2\sqrt{x}e^{-\sqrt{x}} - 2e^{-\sqrt{x}})}{3}$$

Which simplifies to

$$y_p(x) = \frac{4e^{-\sqrt{x}}(7x^{5/2} + 140x^{3/2} + x^3 + 35x^2 + 840\sqrt{x} + 420x + 840)}{x}$$

Therefore the general solution is

$$y = y_h + y_p = \left(\frac{c_1}{x} + \frac{c_2 x^2}{3} \right) + \left(\frac{4e^{-\sqrt{x}}(7x^{5/2} + 140x^{3/2} + x^3 + 35x^2 + 840\sqrt{x} + 420x + 840)}{x} \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1}{x} + \frac{c_2 x^2}{3} + \frac{4e^{-\sqrt{x}}(7x^{5/2} + 140x^{3/2} + x^3 + 35x^2 + 840\sqrt{x} + 420x + 840)}{x}$$

Solved as second order ode adjoint method

Time used: 0.795 (sec)

In normal form the ode

$$y'' - \frac{2y}{x^2} = x e^{-\sqrt{x}} \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= 0 \\ q(x) &= -\frac{2}{x^2} \\ r(x) &= x e^{-\sqrt{x}} \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (0)' + \left(-\frac{2\xi(x)}{x^2}\right) &= 0 \\ \xi''(x) - \frac{2\xi(x)}{x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi'' - \frac{2\xi}{x^2} = 0 \quad (1)$$

$$A\xi'' + B\xi' + C\xi = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \end{aligned} \quad (3)$$

$$C = -\frac{2}{x^2}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = \xi e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then ξ is found using the inverse transformation

$$\xi = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.69: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{x} + (-)(0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \tag{2A}$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

The first solution to the original ode in ξ is found from

$$\xi_1 = z_1 e^{\int -\frac{B}{2A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} \xi_1 &= z_1 \\ &= \frac{1}{x} \end{aligned}$$

Which simplifies to

$$\xi_1 = \frac{1}{x}$$

The second solution ξ_2 to the original ode is found using reduction of order

$$\xi_2 = \xi_1 \int \frac{e^{\int -\frac{B}{A} dx}}{\xi_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} \xi_2 &= \xi_1 \int \frac{1}{\xi_1^2} dx \\ &= \frac{1}{x} \int \frac{1}{\frac{1}{x^2}} dx \\ &= \frac{1}{x} \left(\frac{x^3}{3}\right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} \xi &= c_1 \xi_1 + c_2 \xi_2 \\ &= c_1 \left(\frac{1}{x}\right) + c_2 \left(\frac{1}{x} \left(\frac{x^3}{3}\right)\right) \end{aligned}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' - \frac{y \left(-\frac{c_1}{x^2} + \frac{2c_2 x}{3} \right)}{\frac{c_1}{x} + \frac{c_2 x^2}{3}} = \frac{2c_2 \left(-x^{7/2} e^{-\sqrt{x}} - 7x^3 e^{-\sqrt{x}} - 42x^{5/2} e^{-\sqrt{x}} - 210x^2 e^{-\sqrt{x}} - 840x^{3/2} e^{-\sqrt{x}} - 2520x e^{-\sqrt{x}} - 5040\sqrt{x} e^{-\sqrt{x}} - 5040 e^{-\sqrt{x}} \right)}{3 \left(\frac{c_1}{x} + \frac{c_2 x^2}{3} \right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{2c_2 x^3 - 3c_1}{x(c_2 x^3 + 3c_1)} \\ p(x) &= -\frac{14 \left(120c_2 x^{3/2} + 6x^{5/2} c_2 + \frac{x^{7/2} c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720) c_2 + \frac{3c_1}{7} \right) x e^{-\sqrt{x}}}{c_2 x^3 + 3c_1}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{2c_2 x^3 - 3c_1}{x(c_2 x^3 + 3c_1)} dx} \\ &= \frac{x}{c_2 x^3 + 3c_1}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(-\frac{14 \left(120c_2 x^{3/2} + 6x^{5/2} c_2 + \frac{x^{7/2} c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720) c_2 + \frac{3c_1}{7} \right) x e^{-\sqrt{x}}}{c_2 x^3 + 3c_1} \right) \\ \frac{d}{dx} \left(\frac{yx}{c_2 x^3 + 3c_1} \right) &= \left(\frac{x}{c_2 x^3 + 3c_1} \right) \left(-\frac{14 \left(120c_2 x^{3/2} + 6x^{5/2} c_2 + \frac{x^{7/2} c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720) c_2 + \frac{3c_1}{7} \right) x e^{-\sqrt{x}}}{c_2 x^3 + 3c_1} \right) \\ d \left(\frac{yx}{c_2 x^3 + 3c_1} \right) &= \left(-\frac{14 \left(120c_2 x^{3/2} + 6x^{5/2} c_2 + \frac{x^{7/2} c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720) c_2 + \frac{3c_1}{7} \right) x^2 e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{yx}{c_2 x^3 + 3c_1} &= \int -\frac{14 \left(120c_2 x^{3/2} + 6x^{5/2} c_2 + \frac{x^{7/2} c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720) c_2 + \frac{3c_1}{7} \right) x^2 e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx \\ &= \int -\frac{14 \left(120c_2 x^{3/2} + 6x^{5/2} c_2 + \frac{x^{7/2} c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720) c_2 + \frac{3c_1}{7} \right) x^2 e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx\end{aligned}$$

Dividing throughout by the integrating factor $\frac{x}{c_2 x^3 + 3c_1}$ gives the final solution

$$y = \frac{(c_2 x^3 + 3c_1) \left(\int - \frac{14 \left(120c_2 x^{3/2} + 6x^{5/2}c_2 + \frac{x^{7/2}c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720)c_2 + \frac{3c_1}{7} \right) x^2 e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + c_3 \right)}{x}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{(c_2 x^3 + 3c_1) \left(\int - \frac{14 \left(120c_2 x^{3/2} + 6x^{5/2}c_2 + \frac{x^{7/2}c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720)c_2 + \frac{3c_1}{7} \right) x^2 e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + c_3 \right)}{x}$$

The constants can be merged to give

$$y = \frac{(c_2 x^3 + 3c_1) \left(\int - \frac{14 \left(120c_2 x^{3/2} + 6x^{5/2}c_2 + \frac{x^{7/2}c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720)c_2 + \frac{3c_1}{7} \right) x^2 e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + 1 \right)}{x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{(c_2 x^3 + 3c_1) \left(\int - \frac{14 \left(120c_2 x^{3/2} + 6x^{5/2}c_2 + \frac{x^{7/2}c_2}{7} + \left(\frac{3c_1}{7} + 720c_2 \right) \sqrt{x} + (x^3 + 30x^2 + 360x + 720)c_2 + \frac{3c_1}{7} \right) x^2 e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + 1 \right)}{x}$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    <- LODE of Euler type successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.015 (sec)

Leaf size : 51

```

dsolve(diff(diff(y(x),x),x)-2/x^2*y(x) = x*exp(-x^(1/2)),
        y(x),singsol=all)

```

$$y = \frac{4e^{-\sqrt{x}}(7x^{5/2} + 140x^{3/2} + x^3 + 35x^2 + 840\sqrt{x} + 420x + 840) + c_1x^3 + c_2}{x}$$

Mathematica DSolve solution

Solving time : 0.067 (sec)

Leaf size : 54

```
DSolve[{D[y[x],{x,2}]-2/x^2*y[x] == x*Exp[-x^(1/2)],{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{-2e^{-\sqrt{x}}(\sqrt{x} + 1)x^3 + 3(c_2x^3 + c_1) + 2\Gamma(8, \sqrt{x})}{3x}$$

2.2.15 problem 16

Solved as second order ode using Kovacic algorithm 564
 Solved as second order ode adjoint method 570
 Maple step by step solution 575
 Maple trace 575
 Maple dsolve solution 575
 Mathematica DSolve solution 576

Internal problem ID [8814]

Book : Second order enumerated odes

Section : section 2

Problem number : 16

Date solved : Thursday, December 12, 2024 at 09:51:52 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' - \frac{y'}{\sqrt{x}} + \frac{(x + \sqrt{x} - 8)y}{4x^2} = x$$

Solved as second order ode using Kovacic algorithm

Time used: 0.286 (sec)

Writing the ode as

$$y'' - \frac{y'}{\sqrt{x}} + \left(\frac{1}{4x} + \frac{1}{4x^{3/2}} - \frac{2}{x^2} \right) y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -\frac{1}{\sqrt{x}} \\ C &= \frac{1}{4x} + \frac{1}{4x^{3/2}} - \frac{2}{x^2} \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.70: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{x} + (-)(0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \tag{1A}$$

Let

$$p(x) = 1 \tag{2A}$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-\frac{1}{\sqrt{x}}}{1} dx} \\ &= z_1 e^{\sqrt{x}} \\ &= z_1 \left(e^{\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{e^{\sqrt{x}}}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-\frac{1}{\sqrt{x}}}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2\sqrt{x}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{e^{\sqrt{x}}}{x} \right) + c_2 \left(\frac{e^{\sqrt{x}}}{x} \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - \frac{y'}{\sqrt{x}} + \left(\frac{1}{4x} + \frac{1}{4x^{3/2}} - \frac{2}{x^2} \right) y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1 e^{\sqrt{x}}}{x} + \frac{c_2 x^2 e^{\sqrt{x}}}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{e^{\sqrt{x}}}{x}$$

$$y_2 = \frac{x^2 e^{\sqrt{x}}}{3}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{e^{\sqrt{x}}}{x} & \frac{x^2 e^{\sqrt{x}}}{3} \\ \frac{d}{dx} \left(\frac{e^{\sqrt{x}}}{x} \right) & \frac{d}{dx} \left(\frac{x^2 e^{\sqrt{x}}}{3} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{e^{\sqrt{x}}}{x} & \frac{x^2 e^{\sqrt{x}}}{3} \\ \frac{e^{\sqrt{x}}}{2x^{3/2}} - \frac{e^{\sqrt{x}}}{x^2} & \frac{2x e^{\sqrt{x}}}{3} + \frac{x^{3/2} e^{\sqrt{x}}}{6} \end{vmatrix}$$

Therefore

$$W = \left(\frac{e^{\sqrt{x}}}{x} \right) \left(\frac{2x e^{\sqrt{x}}}{3} + \frac{x^{3/2} e^{\sqrt{x}}}{6} \right) - \left(\frac{x^2 e^{\sqrt{x}}}{3} \right) \left(\frac{e^{\sqrt{x}}}{2x^{3/2}} - \frac{e^{\sqrt{x}}}{x^2} \right)$$

Which simplifies to

$$W = e^{2\sqrt{x}}$$

Which simplifies to

$$W = e^{2\sqrt{x}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 e^{\sqrt{x}}}{e^{2\sqrt{x}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^3 e^{-\sqrt{x}}}{3} dx$$

Hence

$$u_1 = \frac{2x^{7/2}e^{-\sqrt{x}}}{3} + \frac{14x^3e^{-\sqrt{x}}}{3} + 28x^{5/2}e^{-\sqrt{x}} + 140x^2e^{-\sqrt{x}} + 560x^{3/2}e^{-\sqrt{x}} + 1680xe^{-\sqrt{x}} + 3360e^{-\sqrt{x}}\sqrt{x} + 3360e^{-\sqrt{x}}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{\sqrt{x}}}{e^{2\sqrt{x}}} dx$$

Which simplifies to

$$u_2 = \int e^{-\sqrt{x}} dx$$

Hence

$$u_2 = -2e^{-\sqrt{x}}\sqrt{x} - 2e^{-\sqrt{x}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\frac{2x^{7/2}e^{-\sqrt{x}}}{3} + \frac{14x^3e^{-\sqrt{x}}}{3} + 28x^{5/2}e^{-\sqrt{x}} + 140x^2e^{-\sqrt{x}} + 560x^{3/2}e^{-\sqrt{x}} + 1680xe^{-\sqrt{x}} + 3360e^{-\sqrt{x}}\sqrt{x} + 3360e^{-\sqrt{x}} \right) - x^2e^{\sqrt{x}}(-2e^{-\sqrt{x}}\sqrt{x} - 2e^{-\sqrt{x}})}{3x}$$

Which simplifies to

$$y_p(x) = \frac{28x^{5/2} + 560x^{3/2} + 4x^3 + 140x^2 + 3360\sqrt{x} + 1680x + 3360}{x}$$

Therefore the general solution is

$$y = y_h + y_p = \left(\frac{c_1 e^{\sqrt{x}}}{x} + \frac{c_2 x^2 e^{\sqrt{x}}}{3} \right) + \left(\frac{28x^{5/2} + 560x^{3/2} + 4x^3 + 140x^2 + 3360\sqrt{x} + 1680x + 3360}{x} \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 e^{\sqrt{x}}}{x} + \frac{c_2 x^2 e^{\sqrt{x}}}{3} + \frac{28x^{5/2} + 560x^{3/2} + 4x^3 + 140x^2 + 3360\sqrt{x} + 1680x + 3360}{x}$$

Solved as second order ode adjoint method

Time used: 1.194 (sec)

In normal form the ode

$$y'' - \frac{y'}{\sqrt{x}} + \frac{(x + \sqrt{x} - 8)y}{4x^2} = x \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{1}{\sqrt{x}} \\ q(x) &= \frac{1}{4x} + \frac{1}{4x^{3/2}} - \frac{2}{x^2} \\ r(x) &= x \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(-\frac{\xi(x)}{\sqrt{x}}\right)' + \left(\left(\frac{1}{4x} + \frac{1}{4x^{3/2}} - \frac{2}{x^2}\right)\xi(x)\right) &= 0 \\ \xi''(x) + \frac{\xi'(x)}{\sqrt{x}} - \frac{(\sqrt{x} - x + 8)\xi(x)}{4x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi'' + \frac{\xi'}{\sqrt{x}} + \left(-\frac{1}{4x^{3/2}} + \frac{1}{4x} - \frac{2}{x^2}\right)\xi = 0 \quad (1)$$

$$A\xi'' + B\xi' + C\xi = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= \frac{1}{\sqrt{x}} \\ C &= -\frac{1}{4x^{3/2}} + \frac{1}{4x} - \frac{2}{x^2} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = \xi e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then ξ is found using the inverse transformation

$$\xi = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.71: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{x} + (-)(0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \tag{1A}$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

The first solution to the original ode in ξ is found from

$$\begin{aligned} \xi_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{\sqrt{x}} dx} \\ &= z_1 e^{-\sqrt{x}} \\ &= z_1 \left(e^{-\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$\xi_1 = \frac{e^{-\sqrt{x}}}{x}$$

The second solution ξ_2 to the original ode is found using reduction of order

$$\xi_2 = \xi_1 \int \frac{e^{\int -\frac{B}{A} dx}}{\xi_1^2} dx$$

Substituting gives

$$\begin{aligned} \xi_2 &= \xi_1 \int \frac{e^{\int -\frac{1}{2} \frac{1}{\sqrt{x}} dx}}{(\xi_1)^2} dx \\ &= \xi_1 \int \frac{e^{-2\sqrt{x}}}{(\xi_1)^2} dx \\ &= \xi_1 \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} \xi &= c_1 \xi_1 + c_2 \xi_2 \\ &= c_1 \left(\frac{e^{-\sqrt{x}}}{x} \right) + c_2 \left(\frac{e^{-\sqrt{x}}}{x} \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(-\frac{1}{\sqrt{x}} - \frac{-\frac{c_1 e^{-\sqrt{x}}}{2x^{3/2}} - \frac{c_1 e^{-\sqrt{x}}}{x^2} + \frac{2c_2 x e^{-\sqrt{x}}}{3} - \frac{c_2 x^{3/2} e^{-\sqrt{x}}}{6}}{\frac{c_1 e^{-\sqrt{x}}}{x} + \frac{c_2 x^2 e^{-\sqrt{x}}}{3}} \right) = \frac{2c_2 (-x^{7/2} e^{-\sqrt{x}} - 7x^3 e^{-\sqrt{x}} - 42x^{5/2} e^{-\sqrt{x}} - 210x^2 e^{-\sqrt{x}} - 840x e^{-\sqrt{x}} - 420 e^{-\sqrt{x}})}{3}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{4x^{7/2}c_2 - 6c_1\sqrt{x} + x(c_2x^3 + 3c_1)}{x^{3/2}(2c_2x^3 + 6c_1)} \\ p(x) &= -\frac{2\sqrt{x}(2520c_2x^{3/2} + 210c_2x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1))}{c_2x^3 + 3c_1}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{4x^{7/2}c_2 - 6c_1\sqrt{x} + x(c_2x^3 + 3c_1)}{x^{3/2}(2c_2x^3 + 6c_1)} dx} \\ &= \frac{x e^{-\sqrt{x}}}{c_2 x^3 + 3c_1}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(-\frac{2\sqrt{x}(2520c_2x^{3/2} + 210c_2x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1))}{c_2x^3 + 3c_1} \right) \\ \frac{d}{dx} \left(\frac{yx e^{-\sqrt{x}}}{c_2x^3 + 3c_1} \right) &= \left(\frac{x e^{-\sqrt{x}}}{c_2x^3 + 3c_1} \right) \left(-\frac{2\sqrt{x}(2520c_2x^{3/2} + 210c_2x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1))}{c_2x^3 + 3c_1} \right) \\ \frac{d}{dx} \left(\frac{yx e^{-\sqrt{x}}}{c_2x^3 + 3c_1} \right) &= \left(-\frac{2x^{3/2}(2520c_2x^{3/2} + 210c_2x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1))}{(c_2x^3 + 3c_1)^2} \right)\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{yx e^{-\sqrt{x}}}{c_2x^3 + 3c_1} &= \int -\frac{2x^{3/2}(2520c_2x^{3/2} + 210c_2x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1))}{(c_2x^3 + 3c_1)^2} dx \\ &= \int -\frac{2x^{3/2}(2520c_2x^{3/2} + 210c_2x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1))}{(c_2x^3 + 3c_1)^2} dx\end{aligned}$$

Dividing throughout by the integrating factor $\frac{x e^{-\sqrt{x}}}{c_2 x^3 + 3c_1}$ gives the final solution

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}} \left(\int -\frac{2x^{3/2}(2520c_2 x^{3/2} + 210c_2 x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1)x)e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + C \right)}{x}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}} \left(\int -\frac{2x^{3/2}(2520c_2 x^{3/2} + 210c_2 x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1)x)e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + C \right)}{x}$$

The constants can be merged to give

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}} \left(\int -\frac{2x^{3/2}(2520c_2 x^{3/2} + 210c_2 x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1)x)e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + C \right)}{x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}} \left(\int -\frac{2x^{3/2}(2520c_2 x^{3/2} + 210c_2 x^{5/2} + 7x^{7/2}c_2 + (3c_1 + 5040c_2)\sqrt{x} + ((x^3 + 42x^2 + 840x + 5040)c_2 + 3c_1)x)e^{-\sqrt{x}}}{(c_2 x^3 + 3c_1)^2} dx + C \right)}{x}$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
        A Liouvillian solution exists
        Reducible group (found an exponential solution)
    <- Kovacic's algorithm successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.014 (sec)

Leaf size : 50

```

dsolve(diff(diff(y(x), x), x) - 1/x^(1/2)*diff(y(x), x) + 1/4/x^2*(x+x^(1/2)-8)*y(x) = x,
        y(x), singsol=all)

```

$$y = \frac{560x^{3/2} + 28x^{5/2} + (c_1 x^3 + c_2) e^{\sqrt{x}} + 4x^3 + 140x^2 + 1680x + 3360\sqrt{x} + 3360}{x}$$

Mathematica DSolve solution

Solving time : 0.08 (sec)

Leaf size : 63

```
DSolve[{D[y[x],{x,2}]-1/Sqrt[x]*D[y[x],x]+1/(4*x^2)*(x+Sqrt[x]-8)*y[x]==x,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{-2x^{7/2} + x^3(-2 + c_2 e^{\sqrt{x}}) + 2e^{\sqrt{x}}\Gamma(8, \sqrt{x}) + 3c_1 e^{\sqrt{x}}}{3x}$$

2.2.16 problem 17

Solved as second order ode using change of variable on x method 2	577
Solved as second order ode using change of variable on x method 1	579
Solved as second order Bessel ode	580
Solved as second order ode using Kovacic algorithm	581
Solved as second order ode adjoint method	585
Maple step by step solution	587
Maple trace	588
Maple dsolve solution	589
Mathematica DSolve solution	589

Internal problem ID [8815]

Book : Second order enumerated odes

Section : section 2

Problem number : 17

Date solved : Thursday, December 12, 2024 at 09:51:54 AM

CAS classification :

[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F(x)]']]

Solve

$$y'' + \frac{2y'}{x} + \frac{a^2y}{x^4} = 0$$

Solved as second order ode using change of variable on x method 2

Time used: 0.346 (sec)

In normal form the ode

$$y'' + \frac{2y'}{x} + \frac{a^2y}{x^4} = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{2}{x}$$

$$q(x) = \frac{a^2}{x^4}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned}
 \tau &= \int e^{-\int p(x)dx} dx \\
 &= \int e^{-\int \frac{2}{x} dx} dx \\
 &= \int e^{-2\ln(x)} dx \\
 &= \int \frac{1}{x^2} dx \\
 &= -\frac{1}{x}
 \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}
 q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\
 &= \frac{a^2}{\frac{1}{x^4}} \\
 &= a^2
 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}
 \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) + a^2y(\tau) &= 0
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = a^2$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} + a^2 e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$a^2 + \lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = a^2$ into the above gives

$$\begin{aligned}
 \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(a^2)} \\
 &= \pm \sqrt{-a^2}
 \end{aligned}$$

Hence

$$\begin{aligned}
 \lambda_1 &= +\sqrt{-a^2} \\
 \lambda_2 &= -\sqrt{-a^2}
 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
 \lambda_1 &= \sqrt{-a^2} \\
 \lambda_2 &= -\sqrt{-a^2}
 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(\sqrt{-a^2})\tau} + c_2 e^{(-\sqrt{-a^2})\tau}$$

Or

$$y(\tau) = c_1 e^{\tau\sqrt{-a^2}} + c_2 e^{-\tau\sqrt{-a^2}}$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 e^{-\frac{\sqrt{-a^2}}{x}} + c_2 e^{\frac{\sqrt{-a^2}}{x}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{\sqrt{-a^2}}{x}} + c_2 e^{\frac{\sqrt{-a^2}}{x}}$$

Solved as second order ode using change of variable on x method 1

Time used: 0.091 (sec)

In normal form the ode

$$y'' + \frac{2y'}{x} + \frac{a^2 y}{x^4} = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$

$$q(x) = \frac{a^2}{x^4}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{\sqrt{\frac{a^2}{x^4}}}{c} \quad (6)$$

$$\tau'' = -\frac{2a^2}{c\sqrt{\frac{a^2}{x^4}}x^5}$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{2a^2}{c\sqrt{\frac{a^2}{x^4}}x^5} + \frac{2}{x}\frac{\sqrt{\frac{a^2}{x^4}}}{c}}{\left(\frac{\sqrt{\frac{a^2}{x^4}}}{c}\right)^2} \\ &= 0 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{\frac{a^2}{x^4}} dx}{c} \\ &= -\frac{x\sqrt{\frac{a^2}{x^4}}}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos\left(x\sqrt{\frac{a^2}{x^4}}\right) - c_2 \sin\left(x\sqrt{\frac{a^2}{x^4}}\right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos\left(x\sqrt{\frac{a^2}{x^4}}\right) - c_2 \sin\left(x\sqrt{\frac{a^2}{x^4}}\right)$$

Solved as second order Bessel ode

Time used: 0.070 (sec)

Writing the ode as

$$x^2 y'' + 2y'x + \frac{a^2 y}{x^2} = 0 \quad (1)$$

Bessel ode has the form

$$x^2 y'' + y'x + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= -\frac{1}{2} \\ \beta &= a \\ n &= \frac{1}{2} \\ \gamma &= -1 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1 \sqrt{2} \sin\left(\frac{a}{x}\right)}{\sqrt{x} \sqrt{\pi} \sqrt{\frac{a}{x}}} - \frac{c_2 \sqrt{2} \cos\left(\frac{a}{x}\right)}{\sqrt{x} \sqrt{\pi} \sqrt{\frac{a}{x}}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 \sqrt{2} \sin\left(\frac{a}{x}\right)}{\sqrt{x} \sqrt{\pi} \sqrt{\frac{a}{x}}} - \frac{c_2 \sqrt{2} \cos\left(\frac{a}{x}\right)}{\sqrt{x} \sqrt{\pi} \sqrt{\frac{a}{x}}}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.227 (sec)

Writing the ode as

$$y'' + \frac{2y'}{x} + \frac{a^2 y}{x^4} = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= \frac{2}{x} \\ C &= \frac{a^2}{x^4} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-a^2}{x^4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -a^2 \\ t &= x^4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{a^2}{x^4}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.72: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 4 - 0 \\ &= 4 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^4$. There is a pole at $x = 0$ of order 4. Since there is no odd order pole larger than 2 and the order at ∞ is 4 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Looking at higher order poles of order $2v \geq 4$ (must be even order for case one). Then for each pole c , $[\sqrt{r}]_c$ is the sum of terms $\frac{1}{(x-c)^i}$ for $2 \leq i \leq v$ in the Laurent series expansion of \sqrt{r} expanded around each pole c . Hence

$$[\sqrt{r}]_c = \sum_2^v \frac{a_i}{(x-c)^i} \quad (1B)$$

Let a be the coefficient of the term $\frac{1}{(x-c)^v}$ in the above where v is the pole order divided by 2. Let b be the coefficient of $\frac{1}{(x-c)^{v+1}}$ in r minus the coefficient of $\frac{1}{(x-c)^{v+1}}$ in $[\sqrt{r}]_c$. Then

$$\begin{aligned} \alpha_c^+ &= \frac{1}{2} \left(\frac{b}{a} + v \right) \\ \alpha_c^- &= \frac{1}{2} \left(-\frac{b}{a} + v \right) \end{aligned}$$

The partial fraction decomposition of r is

$$r = -\frac{a^2}{x^4}$$

There is pole in r at $x = 0$ of order 4, hence $v = 2$. Expanding \sqrt{r} as Laurent series about this pole $c = 0$ gives

$$[\sqrt{r}]_c \approx \frac{ia}{x^2} + \dots \quad (2B)$$

Using eq. (1B), taking the sum up to $v = 2$ the above becomes

$$[\sqrt{r}]_c = \frac{ia}{x^2} \quad (3B)$$

The above shows that the coefficient of $\frac{1}{(x-0)^2}$ is

$$a = ia$$

Now we need to find b . let b be the coefficient of the term $\frac{1}{(x-c)^{v+1}}$ in r minus the coefficient of the same term but in the sum $[\sqrt{r}]_c$ found in eq. (3B). Here c is current pole which is $c = 0$. This term becomes $\frac{1}{x^3}$. The coefficient of this term in the sum $[\sqrt{r}]_c$ is seen to be 0 and the coefficient of this term r is found from the partial fraction decomposition from above to be 0. Therefore

$$\begin{aligned} b &= (0) - (0) \\ &= 0 \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_c &= \frac{ia}{x^2} \\ \alpha_c^+ &= \frac{1}{2} \left(\frac{b}{a} + v \right) = \frac{1}{2} \left(\frac{0}{ia} + 2 \right) = 1 \\ \alpha_c^- &= \frac{1}{2} \left(-\frac{b}{a} + v \right) = \frac{1}{2} \left(-\frac{0}{ia} + 2 \right) = 1 \end{aligned}$$

Since the order of r at ∞ is $4 > 2$ then

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= 0 \\ \alpha_\infty^- &= 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{a^2}{x^4}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	4	$\frac{ia}{x^2}$	1	1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
4	0	0	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= 1 - (1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= -\frac{ia}{x^2} + \frac{1}{x} + (-) (0) \\ &= -\frac{ia}{x^2} + \frac{1}{x} \\ &= \frac{-ia + x}{x^2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-\frac{ia}{x^2} + \frac{1}{x} \right) (0) + \left(\left(\frac{2ia}{x^3} - \frac{1}{x^2} \right) + \left(-\frac{ia}{x^2} + \frac{1}{x} \right)^2 - \left(-\frac{a^2}{x^4} \right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \left(-\frac{ia}{x^2} + \frac{1}{x} \right) dx} \\ &= x e^{\frac{ia}{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\ &= z_1 e^{-\ln(x)} \\ &= z_1 \left(\frac{1}{x} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{\frac{ia}{x}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{x} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{ie^{-\frac{2ia}{x}}}{2a} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{\frac{ia}{x}} \right) + c_2 \left(e^{\frac{ia}{x}} \left(-\frac{ie^{-\frac{2ia}{x}}}{2a} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{\frac{ia}{x}} - \frac{ic_2 e^{-\frac{ia}{x}}}{2a}$$

Solved as second order ode adjoint method

Time used: 0.441 (sec)

In normal form the ode

$$y'' + \frac{2y'}{x} + \frac{a^2 y}{x^4} = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= \frac{2}{x} \\ q(x) &= \frac{a^2}{x^4} \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(\frac{2\xi(x)}{x} \right)' + \left(\frac{a^2 \xi(x)}{x^4} \right) &= 0 \\ \xi''(x) - \frac{2\xi'(x)}{x} + \frac{(a^2 + 2x^2)\xi(x)}{x^4} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$x^2 \xi'' - 2\xi' x + \left(2 + \frac{a^2}{x^2} \right) \xi = 0 \tag{1}$$

Bessel ode has the form

$$x^2 \xi'' + \xi' x + (-n^2 + x^2) \xi = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 \xi'' + (1 - 2\alpha) x \xi' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) \xi = 0 \quad (3)$$

With the standard solution

$$\xi = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= \frac{3}{2} \\ \beta &= a \\ n &= -\frac{1}{2} \\ \gamma &= -1 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$\xi = \frac{c_1 x^{3/2} \sqrt{2} \cos\left(\frac{a}{x}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x}}} + \frac{c_2 x^{3/2} \sqrt{2} \sin\left(\frac{a}{x}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x}}}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(\frac{2}{x} - \frac{\frac{3c_1 \sqrt{x} \sqrt{2} \cos\left(\frac{a}{x}\right)}{2\sqrt{\pi} \sqrt{\frac{a}{x}}} + \frac{c_1 \sqrt{2} \cos\left(\frac{a}{x}\right) a}{2\sqrt{x} \sqrt{\pi} \left(\frac{a}{x}\right)^{3/2}} + \frac{c_1 \sqrt{2} a \sin\left(\frac{a}{x}\right)}{\sqrt{x} \sqrt{\pi} \sqrt{\frac{a}{x}}} + \frac{3c_2 \sqrt{x} \sqrt{2} \sin\left(\frac{a}{x}\right)}{2\sqrt{\pi} \sqrt{\frac{a}{x}}} + \frac{c_2 \sqrt{2} \sin\left(\frac{a}{x}\right) a}{2\sqrt{x} \sqrt{\pi} \left(\frac{a}{x}\right)^{3/2}} - \frac{c_2 \sqrt{2} a \cos\left(\frac{a}{x}\right)}{\sqrt{x} \sqrt{\pi} \sqrt{\frac{a}{x}}}}{\frac{c_1 x^{3/2} \sqrt{2} \cos\left(\frac{a}{x}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x}}} + \frac{c_2 x^{3/2} \sqrt{2} \sin\left(\frac{a}{x}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x}}}} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{(-\cos\left(\frac{a}{x}\right) c_2 + \sin\left(\frac{a}{x}\right) c_1) a}{(c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)) x^2} \\ p(x) &= 0 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(-\cos\left(\frac{a}{x}\right) c_2 + \sin\left(\frac{a}{x}\right) c_1) a}{(c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)) x^2} dx} \\ &= \frac{1}{c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$

$$\frac{d}{dx} \left(\frac{y}{c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)} \right) = 0$$

Integrating gives

$$\frac{y}{c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)} = \int 0 dx + c_3$$

$$= c_3$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)}$ gives the final solution

$$y = \left(c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right) \right) c_3$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \left(c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right) \right) c_3$$

The constants can be merged to give

$$y = c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos\left(\frac{a}{x}\right) + c_2 \sin\left(\frac{a}{x}\right)$$

Maple step by step solution

Let's solve

$$\frac{d^2}{dx^2} y(x) + \frac{2\left(\frac{d}{dx} y(x)\right)}{x} + \frac{a^2 y(x)}{x^4} = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Multiply by denominators of the ODE

$$\left(\frac{d^2}{dx^2} y(x) \right) x^4 + 2 \left(\frac{d}{dx} y(x) \right) x^3 + a^2 y(x) = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$\frac{d}{dx} y(x) = \left(\frac{d}{dt} y(t) \right) \left(\frac{d}{dx} t(x) \right)$$

- Compute derivative

$$\frac{d}{dx} y(x) = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$\frac{d^2}{dx^2} y(x) = \left(\frac{d^2}{dt^2} y(t) \right) \left(\frac{d}{dx} t(x) \right)^2 + \left(\frac{d^2}{dx^2} t(x) \right) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$\frac{d^2}{dx^2} y(x) = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$\left(\frac{d^2 y(t)}{dt^2} - \frac{d y(t)}{dt} \right) x^4 + 2 \left(\frac{d y(t)}{dt} \right) x^2 + a^2 y(t) = 0$$

- Simplify

$$x^2 \left(\frac{d^2 y(t)}{dt^2} \right) + \left(\frac{d y(t)}{dt} \right) x^2 + a^2 y(t) = 0$$

- Isolate 2nd derivative

$$\frac{d^2 y(t)}{dt^2} = -\frac{a^2 y(t)}{x^2} - \frac{d y(t)}{dt}$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2 y(t)}{dt^2} + \frac{d y(t)}{dt} + \frac{a^2 y(t)}{x^2} = 0$$

- Characteristic polynomial of ODE

$$r^2 + r + \frac{a^2}{x^2} = 0$$

- Factor the characteristic polynomial

$$\frac{r^2 x^2 + r x^2 + a^2}{x^2} = 0$$

- Roots of the characteristic polynomial

$$r = \left(\frac{-\frac{x}{2} + \frac{\sqrt{-4a^2 + x^2}}{2}}{x}, \frac{-\frac{x}{2} - \frac{\sqrt{-4a^2 + x^2}}{2}}{x} \right)$$

- 1st solution of the ODE

$$y_1(t) = e^{\frac{\left(-\frac{x}{2} + \frac{\sqrt{-4a^2 + x^2}}{2}\right)t}{x}}$$

- 2nd solution of the ODE

$$y_2(t) = e^{\frac{\left(-\frac{x}{2} - \frac{\sqrt{-4a^2 + x^2}}{2}\right)t}{x}}$$

- General solution of the ODE

$$y(t) = C1 y_1(t) + C2 y_2(t)$$

- Substitute in solutions

$$y(t) = C1 e^{\frac{\left(-\frac{x}{2} + \frac{\sqrt{-4a^2 + x^2}}{2}\right)t}{x}} + C2 e^{\frac{\left(-\frac{x}{2} - \frac{\sqrt{-4a^2 + x^2}}{2}\right)t}{x}}$$

- Change variables back using $t = \ln(x)$

$$y(x) = C1 e^{\frac{\left(-\frac{x}{2} + \frac{\sqrt{-4a^2 + x^2}}{2}\right)\ln(x)}{x}} + C2 e^{\frac{\left(-\frac{x}{2} - \frac{\sqrt{-4a^2 + x^2}}{2}\right)\ln(x)}{x}}$$

- Simplify

$$y(x) = C1 x^{-\frac{x + \sqrt{-4a^2 + x^2}}{2x}} + C2 x^{-\frac{x - \sqrt{-4a^2 + x^2}}{2x}}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

Maple dsolve solution

Solving time : 0.004 (sec)

Leaf size : 21

```
dsolve(diff(diff(y(x),x),x)+2/x*diff(y(x),x)+a^2/x^4*y(x) = 0,  
y(x),singsol=all)
```

$$y = c_1 \sin\left(\frac{a}{x}\right) + c_2 \cos\left(\frac{a}{x}\right)$$

Mathematica DSolve solution

Solving time : 0.031 (sec)

Leaf size : 25

```
DSolve[{D[y[x],{x,2}]+2/x*D[y[x],x]+a^2/x^4*y[x]==0,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_1 \cos\left(\frac{a}{x}\right) - c_2 \sin\left(\frac{a}{x}\right)$$

2.2.17 problem 18

Solved as second order ode using change of variable on x
 method 2 590
 Solved as second order ode using change of variable on x
 method 1 592
 Solved as second order ode using Kovacic algorithm 594
 Solved as second order ode adjoint method 598
 Maple step by step solution 603
 Maple trace 605
 Maple dsolve solution 605
 Mathematica DSolve solution 605

Internal problem ID [8816]

Book : Second order enumerated odes

Section : section 2

Problem number : 18

Date solved : Thursday, December 12, 2024 at 09:51:56 AM

CAS classification :

[_Gegenbauer, [_2nd_order, _linear, ‘_with_symmetry_[0,F(x)]’]]

Solve

$$(-x^2 + 1) y'' - xy' - c^2y = 0$$

Solved as second order ode using change of variable on x method 2

Time used: 0.484 (sec)

In normal form the ode

$$(-x^2 + 1) y'' - xy' - c^2y = 0 \tag{1}$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$p(x) = \frac{x}{x^2 - 1}$$

$$q(x) = \frac{c^2}{x^2 - 1}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned}
 \tau &= \int e^{-\int p(x)dx} dx \\
 &= \int e^{-\int \frac{x}{x^2-1} dx} dx \\
 &= \int e^{-\frac{\ln(x-1)}{2} - \frac{\ln(x+1)}{2}} dx \\
 &= \int \frac{1}{\sqrt{x-1}\sqrt{x+1}} dx \\
 &= \frac{\sqrt{(x-1)(x+1)} \ln(x + \sqrt{x^2-1})}{\sqrt{x-1}\sqrt{x+1}}
 \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}
 q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\
 &= \frac{\frac{c^2}{x^2-1}}{\frac{1}{(x-1)(x+1)}} \\
 &= c^2
 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}
 \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = c^2$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\tau\lambda} + c^2 e^{\tau\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$c^2 + \lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = c^2$ into the above gives

$$\begin{aligned}
 \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(c^2)} \\
 &= \pm \sqrt{-c^2}
 \end{aligned}$$

Hence

$$\begin{aligned}
 \lambda_1 &= +\sqrt{-c^2} \\
 \lambda_2 &= -\sqrt{-c^2}
 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
 \lambda_1 &= \sqrt{-c^2} \\
 \lambda_2 &= -\sqrt{-c^2}
 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(\sqrt{-c^2})\tau} + c_2 e^{(-\sqrt{-c^2})\tau}$$

Or

$$y(\tau) = c_1 e^{\tau\sqrt{-c^2}} + c_2 e^{-\tau\sqrt{-c^2}}$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 e^{\frac{\sqrt{(x-1)(x+1)} \ln(x+\sqrt{x^2-1}) \sqrt{-c^2}}{\sqrt{x-1}\sqrt{x+1}}} + c_2 e^{-\frac{\sqrt{(x-1)(x+1)} \ln(x+\sqrt{x^2-1}) \sqrt{-c^2}}{\sqrt{x-1}\sqrt{x+1}}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{\frac{\sqrt{(x-1)(x+1)} \ln(x+\sqrt{x^2-1}) \sqrt{-c^2}}{\sqrt{x-1}\sqrt{x+1}}} + c_2 e^{-\frac{\sqrt{(x-1)(x+1)} \ln(x+\sqrt{x^2-1}) \sqrt{-c^2}}{\sqrt{x-1}\sqrt{x+1}}}$$

Solved as second order ode using change of variable on x method 1

Time used: 0.161 (sec)

In normal form the ode

$$(-x^2 + 1)y'' - xy' - c^2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{x}{x^2 - 1}$$

$$q(x) = \frac{c^2}{x^2 - 1}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{\sqrt{\frac{c^2}{x^2-1}}}{c} \quad (6)$$

$$\tau'' = -\frac{c^2x}{c\sqrt{\frac{c^2}{x^2-1}}(x^2-1)^2}$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{c^2 x}{c \sqrt{\frac{c^2}{x^2-1}} (x^2-1)^2} + \frac{x}{x^2-1} \frac{\sqrt{\frac{c^2}{x^2-1}}}{c}}{\left(\frac{\sqrt{\frac{c^2}{x^2-1}}}{c}\right)^2} \\ &= 0 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{\frac{c^2}{x^2-1}} dx}{c} \\ &= \frac{\sqrt{\frac{c^2}{x^2-1}} \sqrt{x^2-1} \ln(x + \sqrt{x^2-1})}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$\begin{aligned} y &= c_1 \cos\left(\sqrt{\frac{c^2}{x^2-1}} \sqrt{x^2-1} \ln(x + \sqrt{x^2-1})\right) \\ &\quad + c_2 \sin\left(\sqrt{\frac{c^2}{x^2-1}} \sqrt{x^2-1} \ln(x + \sqrt{x^2-1})\right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned} y &= c_1 \cos\left(\sqrt{\frac{c^2}{x^2-1}} \sqrt{x^2-1} \ln(x + \sqrt{x^2-1})\right) \\ &\quad + c_2 \sin\left(\sqrt{\frac{c^2}{x^2-1}} \sqrt{x^2-1} \ln(x + \sqrt{x^2-1})\right) \end{aligned}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.320 (sec)

Writing the ode as

$$(-x^2 + 1)y'' - xy' - c^2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= -x^2 + 1 \\ B &= -x \\ C &= -c^2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4c^2x^2 + 4c^2 - x^2 - 2}{4(x^2 - 1)^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4c^2x^2 + 4c^2 - x^2 - 2 \\ t &= 4(x^2 - 1)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{-4c^2x^2 + 4c^2 - x^2 - 2}{4(x^2 - 1)^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.74: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 4 - 2 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4(x^2 - 1)^2$. There is a pole at $x = 1$ of order 2. There is a pole at $x = -1$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Unable to find solution using case one

Attempting to find a solution using case $n = 2$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{3}{16(x-1)^2} + \frac{\frac{1}{16} - \frac{c^2}{2}}{x-1} - \frac{3}{16(x+1)^2} + \frac{\frac{c^2}{2} - \frac{1}{16}}{x+1}$$

For the pole at $x = 1$ let b be the coefficient of $\frac{1}{(x-1)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{3}{16}$. Hence

$$\begin{aligned} E_c &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{1, 2, 3\} \end{aligned}$$

For the pole at $x = -1$ let b be the coefficient of $\frac{1}{(x+1)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{3}{16}$. Hence

$$\begin{aligned} E_c &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{1, 2, 3\} \end{aligned}$$

Since the order of r at ∞ is 2 then let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{-4c^2x^2 + 4c^2 - x^2 - 2}{4(x^2 - 1)^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -1$. Hence

$$\begin{aligned} E_\infty &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{2\} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ for case 2 of Kovacic algorithm.

pole c location	pole order	E_c
1	2	{1, 2, 3}
-1	2	{1, 2, 3}
Order of r at ∞	E_∞	
2	{2}	

Using the family $\{e_1, e_2, \dots, e_\infty\}$ given by

$$e_1 = 1, e_2 = 1, e_\infty = 2$$

Gives a non negative integer d (the degree of the polynomial $p(x)$), which is generated using

$$\begin{aligned} d &= \frac{1}{2} \left(e_\infty - \sum_{c \in \Gamma} e_c \right) \\ &= \frac{1}{2} (2 - (1 + (1))) \\ &= 0 \end{aligned}$$

We now form the following rational function

$$\begin{aligned} \theta &= \frac{1}{2} \sum_{c \in \Gamma} \frac{e_c}{x - c} \\ &= \frac{1}{2} \left(\frac{1}{(x - (1))} + \frac{1}{(x - (-1))} \right) \\ &= \frac{1}{2x - 2} + \frac{1}{2x + 2} \end{aligned}$$

Now we search for a monic polynomial $p(x)$ of degree $d = 0$ such that

$$p''' + 3\theta p'' + (3\theta^2 + 3\theta' - 4r) p' + (\theta'' + 3\theta\theta' + \theta^3 - 4r\theta - 2r') p = 0 \quad (1A)$$

Since $d = 0$, then letting

$$p = 1 \quad (2A)$$

Substituting p and θ into Eq. (1A) gives

$$0 = 0$$

And solving for p gives

$$p = 1$$

Now that $p(x)$ is found let

$$\begin{aligned} \phi &= \theta + \frac{p'}{p} \\ &= \frac{1}{2x - 2} + \frac{1}{2x + 2} \end{aligned}$$

Let ω be the solution of

$$\omega^2 - \phi\omega + \left(\frac{1}{2}\phi' + \frac{1}{2}\phi^2 - r \right) = 0$$

Substituting the values for ϕ and r into the above equation gives

$$w^2 - \left(\frac{1}{2x - 2} + \frac{1}{2x + 2} \right) w + \frac{4c^2 x^2 - 4c^2 + x^2}{4(x^2 - 1)^2} = 0$$

Solving for ω gives

$$\omega = \frac{x + 2c\sqrt{-x^2 + 1}}{2(x - 1)(x + 1)}$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= e^{\int \omega dx} \\ &= e^{\int \frac{x + 2c\sqrt{-x^2 + 1}}{2(x - 1)(x + 1)} dx} \\ &= (x^2 - 1)^{1/4} e^{-c \arcsin(x)} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{-x^2+1} dx} \\ &= z_1 e^{-\frac{\ln(x-1)}{4} - \frac{\ln(x+1)}{4}} \\ &= z_1 \left(\frac{1}{(x-1)^{1/4} (x+1)^{1/4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{(x^2 - 1)^{1/4} e^{-c \arcsin(x)}}{(x-1)^{1/4} (x+1)^{1/4}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{-x^2+1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\frac{\ln(x-1)}{2} - \frac{\ln(x+1)}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\int \frac{e^{-\frac{\ln(x-1)}{2} - \frac{\ln(x+1)}{2}} \sqrt{x-1} \sqrt{x+1} e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 \left(\frac{(x^2 - 1)^{1/4} e^{-c \arcsin(x)}}{(x-1)^{1/4} (x+1)^{1/4}} \right) + c_2 \left(\frac{(x^2 - 1)^{1/4} e^{-c \arcsin(x)}}{(x-1)^{1/4} (x+1)^{1/4}} \left(\int \frac{e^{-\frac{\ln(x-1)}{2} - \frac{\ln(x+1)}{2}} \sqrt{x-1} \sqrt{x+1} e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx \right) \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 (x^2 - 1)^{1/4} e^{-c \arcsin(x)}}{(x-1)^{1/4} (x+1)^{1/4}} + \frac{c_2 (x^2 - 1)^{1/4} e^{-c \arcsin(x)} \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx}{(x-1)^{1/4} (x+1)^{1/4}}$$

Solved as second order ode adjoint method

Time used: 0.891 (sec)

In normal form the ode

$$(-x^2 + 1)y'' - xy' - c^2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{x}{x^2 - 1} \\ q(x) &= \frac{c^2}{x^2 - 1} \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(\frac{x\xi(x)}{x^2 - 1}\right)' + \left(\frac{c^2\xi(x)}{x^2 - 1}\right) &= 0 \\ \xi''(x) - \frac{x\xi'(x)}{x^2 - 1} + \frac{(x^2 + 1 + c^2(x^2 - 1))\xi(x)}{(x^2 - 1)^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi'' - \frac{x\xi'}{x^2 - 1} + \frac{(x^2 + 1 + c^2(x^2 - 1))\xi}{(x^2 - 1)^2} = 0 \quad (1)$$

$$A\xi'' + B\xi' + C\xi = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -\frac{x}{x^2 - 1} \\ C &= \frac{x^2 + 1 + c^2(x^2 - 1)}{(x^2 - 1)^2} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = \xi e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4c^2x^2 + 4c^2 - x^2 - 2}{4(x^2 - 1)^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4c^2x^2 + 4c^2 - x^2 - 2 \\ t &= 4(x^2 - 1)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{-4c^2x^2 + 4c^2 - x^2 - 2}{4(x^2 - 1)^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then ξ is found using the inverse transformation

$$\xi = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.75: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 4 - 2 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4(x^2 - 1)^2$. There is a pole at $x = 1$ of order 2. There is a pole at $x = -1$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Unable to find solution using case one

Attempting to find a solution using case $n = 2$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{3}{16(x-1)^2} + \frac{\frac{1}{16} - \frac{c^2}{2}}{x-1} - \frac{3}{16(x+1)^2} + \frac{\frac{c^2}{2} - \frac{1}{16}}{x+1}$$

For the pole at $x = 1$ let b be the coefficient of $\frac{1}{(x-1)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{3}{16}$. Hence

$$\begin{aligned} E_c &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{1, 2, 3\} \end{aligned}$$

For the pole at $x = -1$ let b be the coefficient of $\frac{1}{(x+1)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{3}{16}$. Hence

$$\begin{aligned} E_c &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{1, 2, 3\} \end{aligned}$$

Since the order of r at ∞ is 2 then let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{-4c^2x^2 + 4c^2 - x^2 - 2}{4(x^2 - 1)^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -1$. Hence

$$\begin{aligned} E_\infty &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{2\} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ for case 2 of Kovacic algorithm.

pole c location	pole order	E_c
1	2	$\{1, 2, 3\}$
-1	2	$\{1, 2, 3\}$
Order of r at ∞		E_∞
2		$\{2\}$

Using the family $\{e_1, e_2, \dots, e_\infty\}$ given by

$$e_1 = 1, e_2 = 1, e_\infty = 2$$

Gives a non negative integer d (the degree of the polynomial $p(x)$), which is generated using

$$\begin{aligned} d &= \frac{1}{2} \left(e_\infty - \sum_{c \in \Gamma} e_c \right) \\ &= \frac{1}{2} (2 - (1 + (1))) \\ &= 0 \end{aligned}$$

We now form the following rational function

$$\begin{aligned} \theta &= \frac{1}{2} \sum_{c \in \Gamma} \frac{e_c}{x - c} \\ &= \frac{1}{2} \left(\frac{1}{(x - (1))} + \frac{1}{(x - (-1))} \right) \\ &= \frac{1}{2x - 2} + \frac{1}{2x + 2} \end{aligned}$$

Now we search for a monic polynomial $p(x)$ of degree $d = 0$ such that

$$p''' + 3\theta p'' + (3\theta^2 + 3\theta' - 4r) p' + (\theta'' + 3\theta\theta' + \theta^3 - 4r\theta - 2r') p = 0 \quad (1A)$$

Since $d = 0$, then letting

$$p = 1 \tag{2A}$$

Substituting p and θ into Eq. (1A) gives

$$0 = 0$$

And solving for p gives

$$p = 1$$

Now that $p(x)$ is found let

$$\begin{aligned} \phi &= \theta + \frac{p'}{p} \\ &= \frac{1}{2x-2} + \frac{1}{2x+2} \end{aligned}$$

Let ω be the solution of

$$\omega^2 - \phi\omega + \left(\frac{1}{2}\phi' + \frac{1}{2}\phi^2 - r\right) = 0$$

Substituting the values for ϕ and r into the above equation gives

$$w^2 - \left(\frac{1}{2x-2} + \frac{1}{2x+2}\right)w + \frac{4c^2x^2 - 4c^2 + x^2}{4(x^2-1)^2} = 0$$

Solving for ω gives

$$\omega = \frac{x + 2c\sqrt{-x^2+1}}{2(x-1)(x+1)}$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= e^{\int \omega dx} \\ &= e^{\int \frac{x+2c\sqrt{-x^2+1}}{2(x-1)(x+1)} dx} \\ &= (x^2-1)^{1/4} e^{-c \arcsin(x)} \end{aligned}$$

The first solution to the original ode in ξ is found from

$$\begin{aligned} \xi_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{x^2-1} dx} \\ &= z_1 e^{\frac{\ln(x-1)}{4} + \frac{\ln(x+1)}{4}} \\ &= z_1 \left((x-1)^{1/4} (x+1)^{1/4} \right) \end{aligned}$$

Which simplifies to

$$\xi_1 = (x-1)^{1/4} (x+1)^{1/4} (x^2-1)^{1/4} e^{-c \arcsin(x)}$$

The second solution ξ_2 to the original ode is found using reduction of order

$$\xi_2 = \xi_1 \int \frac{e^{\int -\frac{B}{A} dx}}{\xi_1^2} dx$$

Substituting gives

$$\begin{aligned} \xi_2 &= \xi_1 \int \frac{e^{\int -\frac{x}{x^2-1} dx}}{(\xi_1)^2} dx \\ &= \xi_1 \int \frac{e^{\frac{\ln(x-1)}{2} + \frac{\ln(x+1)}{2}}}{(\xi_1)^2} dx \\ &= \xi_1 \left(\int \frac{e^{\frac{\ln(x-1)}{2} + \frac{\ln(x+1)}{2}} e^{2c \arcsin(x)}}{\sqrt{x-1} \sqrt{x+1} \sqrt{x^2-1}} dx \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} \xi &= c_1 \xi_1 + c_2 \xi_2 \\ &= c_1 \left((x-1)^{1/4} (x+1)^{1/4} (x^2-1)^{1/4} e^{-c \arcsin(x)} \right) + c_2 \left((x-1)^{1/4} (x+1)^{1/4} (x^2-1)^{1/4} e^{-c \arcsin(x)} \left(\int \frac{e^{\frac{\ln(x-1)}{2} + \frac{\ln(x+1)}{2}} e^{2c \arcsin(x)}}{\sqrt{x-1} \sqrt{x+1} \sqrt{x^2-1}} dx \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(\frac{x}{x^2-1} - \frac{\frac{c_1(x+1)^{1/4}(x^2-1)^{1/4}e^{-c \arcsin(x)}}{4(x-1)^{3/4}} + \frac{c_1(x-1)^{1/4}(x^2-1)^{1/4}e^{-c \arcsin(x)}}{4(x+1)^{3/4}} + \frac{c_1(x-1)^{1/4}(x+1)^{1/4}e^{-c \arcsin(x)}x}{2(x^2-1)^{3/4}} - \frac{c_1(x-1)^{1/4}(x+1)^{1/4}e^{-c \arcsin(x)}}{2(x^2-1)^{3/4}}}{\xi(x)} \right) = \frac{\int \xi(x) r(x) dx}{\xi(x)}$$

Which is now a first order ode. This is now solved for y . The ode $y' = -\frac{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c \right) y}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1 \right) \sqrt{-x^2+1} (x-1)(x+1)}$

is separable as it can be written as

$$\begin{aligned} y' &= -\frac{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c \right) y}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1 \right) \sqrt{-x^2+1} (x-1)(x+1)} \\ &= f(x)g(y) \end{aligned}$$

Where

$$f(x) = -\frac{\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1 \right) \sqrt{-x^2+1} (x-1)(x+1)}$$

$$g(y) = y$$

Integrating gives

$$\begin{aligned} \int \frac{1}{g(y)} dy &= \int f(x) dx \\ \int \frac{1}{y} dy &= \int -\frac{\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1 \right) \sqrt{-x^2+1} (x-1)(x+1)} dx \\ \ln(y) &= \int -\frac{\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1 \right) \sqrt{-x^2+1} (x-1)(x+1)} dx \end{aligned}$$

We now need to find the singular solutions, these are found by finding for what values $g(y)$ is zero, since we had to divide by this above. Solving $g(y) = 0$ or $y = 0$ for y gives

$$y = 0$$

Now we go over each such singular solution and check if it verifies the ode itself and any initial conditions given. If it does not then the singular solution will not be used.

Therefore the solutions found are

$$\ln(y) = \int -\frac{\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1\right) \sqrt{-x^2+1} (x-1)(x+1)} dx +$$

y

Hence, the solution found using Lagrange adjoint equation method is

$$\begin{aligned} \ln(y) &= \int \\ & -\frac{\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1\right) \sqrt{-x^2+1} (x-1)(x+1)} dx \\ & + 2c_3 \\ & y = 0 \end{aligned}$$

The constants can be merged to give

$$\begin{aligned} \ln(y) &= \int \\ & -\frac{\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1\right) \sqrt{-x^2+1} (x-1)(x+1)} dx \\ & + 2 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned} \ln(y) &= \int \\ & -\frac{\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 c x^2 - c_2 e^{2c \arcsin(x)} \sqrt{x^2-1} \sqrt{-x^2+1} + c_1 c x^2 - c_2 c \int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx - c_1 c}{\left(\int \frac{e^{2c \arcsin(x)}}{\sqrt{x^2-1}} dx c_2 + c_1\right) \sqrt{-x^2+1} (x-1)(x+1)} dx \\ & + 2 \\ & y = 0 \end{aligned}$$

Maple step by step solution

Let's solve

$$(-x^2 + 1) \left(\frac{d^2}{dx^2} y(x) \right) - x \left(\frac{d}{dx} y(x) \right) - c^2 y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = -\frac{c^2 y(x)}{x^2-1} - \frac{\left(\frac{d}{dx} y(x)\right) x}{x^2-1}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is lin

$$\frac{d^2}{dx^2}y(x) + \frac{\left(\frac{d}{dx}y(x)\right)x}{x^2-1} + \frac{c^2y(x)}{x^2-1} = 0$$

□ Check to see if x_0 is a regular singular point

○ Define functions

$$\left[P_2(x) = \frac{x}{x^2-1}, P_3(x) = \frac{c^2}{x^2-1} \right]$$

○ $(x+1) \cdot P_2(x)$ is analytic at $x = -1$

$$\left. ((x+1) \cdot P_2(x)) \right|_{x=-1} = \frac{1}{2}$$

○ $(x+1)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$\left. ((x+1)^2 \cdot P_3(x)) \right|_{x=-1} = 0$$

○ $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

• Multiply by denominators

$$(x^2 - 1) \left(\frac{d^2}{dx^2}y(x) \right) + x \left(\frac{d}{dx}y(x) \right) + c^2y(x) = 0$$

• Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2}y(u) \right) + (u - 1) \left(\frac{d}{du}y(u) \right) + c^2y(u) = 0$$

• Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $u^m \cdot \left(\frac{d}{du}y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du}y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

○ Shift index using $k- > k+1-m$

$$u^m \cdot \left(\frac{d}{du}y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

○ Convert $u^m \cdot \left(\frac{d^2}{du^2}y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2}y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

○ Shift index using $k- > k+2-m$

$$u^m \cdot \left(\frac{d^2}{du^2}y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-a_0r(-1+2r)u^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{k+1}(k+1+r)(2k+1+2r) + a_k(c^2+k^2+2kr+r^2)) u^{k+r} \right) = 0$$

• a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-1+2r) = 0$$

• Values of r that satisfy the indicial equation

$$r \in \left\{ 0, \frac{1}{2} \right\}$$

• Each term in the series must be 0, giving the recursion relation

$$-2(k+1+r) \left(k + \frac{1}{2} + r \right) a_{k+1} + a_k(c^2+k^2+2kr+r^2) = 0$$

• Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(c^2+k^2+2kr+r^2)}{(k+1+r)(2k+1+2r)}$$

• Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k(c^2+k^2)}{(k+1)(2k+1)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k(c^2+k^2)}{(k+1)(2k+1)} \right]$$

- Revert the change of variables $u = x + 1$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k (x+1)^k, a_{k+1} = \frac{a_k(c^2+k^2)}{(k+1)(2k+1)} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+1} = \frac{a_k(c^2+k^2+k+\frac{1}{4})}{(k+\frac{3}{2})(2k+2)}$$

- Solution for $r = \frac{1}{2}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k(c^2+k^2+k+\frac{1}{4})}{(k+\frac{3}{2})(2k+2)} \right]$$

- Revert the change of variables $u = x + 1$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k (x+1)^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k(c^2+k^2+k+\frac{1}{4})}{(k+\frac{3}{2})(2k+2)} \right]$$

- Combine solutions and rename parameters

$$\left[y(x) = \left(\sum_{k=0}^{\infty} a_k (x+1)^k \right) + \left(\sum_{k=0}^{\infty} b_k (x+1)^{k+\frac{1}{2}} \right), a_{k+1} = \frac{a_k(c^2+k^2)}{(k+1)(2k+1)}, b_{k+1} = \frac{b_k(c^2+k^2+k+\frac{1}{4})}{(k+\frac{3}{2})(2k+2)} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`
```

Maple dsolve solution

Solving time : 0.002 (sec)

Leaf size : 35

```
dsolve((-x^2+1)*diff(diff(y(x),x),x)-diff(y(x),x)*x-c^2*y(x) = 0,
y(x),singsol=all)
```

$$y = c_1 \left(x + \sqrt{x^2 - 1} \right)^{ic} + c_2 \left(x + \sqrt{x^2 - 1} \right)^{-ic}$$

Mathematica DSolve solution

Solving time : 0.086 (sec)

Leaf size : 42

```
DSolve[{(1-x^2)*D[y[x],{x,2}]-x*D[y[x],x]-c^2*y[x]==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_1 \cos \left(\operatorname{carctanh} \left(\frac{x}{\sqrt{x^2 - 1}} \right) \right) + c_2 \sin \left(\operatorname{carctanh} \left(\frac{x}{\sqrt{x^2 - 1}} \right) \right)$$

2.2.18 problem 19

Solved as second order ode using change of variable on x
 method 2 606
 Solved as second order ode using change of variable on x
 method 1 610
 Solved as second order Bessel ode 614
 Solved as second order ode adjoint method 616
 Maple step by step solution 619
 Maple trace 619
 Maple dsolve solution 619
 Mathematica DSolve solution 619

Internal problem ID [8817]

Book : Second order enumerated odes

Section : section 2

Problem number : 19

Date solved : Thursday, December 12, 2024 at 09:51:58 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$x^6 y'' + 3x^5 y' + a^2 y = \frac{1}{x^2}$$

Solved as second order ode using change of variable on x method 2

Time used: 0.490 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^6 y'' + 3x^5 y' + a^2 y = 0$$

In normal form the ode

$$x^6 y'' + 3x^5 y' + a^2 y = 0 \tag{1}$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$p(x) = \frac{3}{x}$$

$$q(x) = \frac{a^2}{x^6}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned}\tau &= \int e^{-\int p(x)dx} dx \\ &= \int e^{-\int \frac{3}{x} dx} dx \\ &= \int e^{-3\ln(x)} dx \\ &= \int \frac{1}{x^3} dx \\ &= -\frac{1}{2x^2}\end{aligned}\tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{a^2}{\frac{1}{x^6}} \\ &= a^2\end{aligned}\tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + a^2y(\tau) &= 0\end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = a^2$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\tau\lambda} + a^2 e^{\tau\lambda} = 0\tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$a^2 + \lambda^2 = 0\tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = a^2$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(a^2)} \\ &= \pm \sqrt{-a^2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +\sqrt{-a^2} \\ \lambda_2 &= -\sqrt{-a^2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \sqrt{-a^2} \\ \lambda_2 &= -\sqrt{-a^2}\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y(\tau) &= c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau} \\ y(\tau) &= c_1 e^{(\sqrt{-a^2})\tau} + c_2 e^{(-\sqrt{-a^2})\tau}\end{aligned}$$

Or

$$y(\tau) = c_1 e^{\tau\sqrt{-a^2}} + c_2 e^{-\tau\sqrt{-a^2}}$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 e^{-\frac{\sqrt{-a^2}}{2x^2}} + c_2 e^{\frac{\sqrt{-a^2}}{2x^2}}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{-\frac{\sqrt{-a^2}}{2x^2}} + c_2 e^{\frac{\sqrt{-a^2}}{2x^2}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= e^{-\frac{\sqrt{-a^2}}{2x^2}} \\ y_2 &= e^{\frac{\sqrt{-a^2}}{2x^2}}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-\frac{\sqrt{-a^2}}{2x^2}} & e^{\frac{\sqrt{-a^2}}{2x^2}} \\ \frac{d}{dx} \left(e^{-\frac{\sqrt{-a^2}}{2x^2}} \right) & \frac{d}{dx} \left(e^{\frac{\sqrt{-a^2}}{2x^2}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-\frac{\sqrt{-a^2}}{2x^2}} & e^{\frac{\sqrt{-a^2}}{2x^2}} \\ \frac{\sqrt{-a^2} e^{-\frac{\sqrt{-a^2}}{2x^2}}}{x^3} & -\frac{\sqrt{-a^2} e^{\frac{\sqrt{-a^2}}{2x^2}}}{x^3} \end{vmatrix}$$

Therefore

$$W = \left(e^{-\frac{\sqrt{-a^2}}{2x^2}} \right) \left(-\frac{\sqrt{-a^2} e^{\frac{\sqrt{-a^2}}{2x^2}}}{x^3} \right) - \left(e^{\frac{\sqrt{-a^2}}{2x^2}} \right) \left(\frac{\sqrt{-a^2} e^{-\frac{\sqrt{-a^2}}{2x^2}}}{x^3} \right)$$

Which simplifies to

$$W = -\frac{2 e^{-\frac{\sqrt{-a^2}}{2x^2}} \sqrt{-a^2} e^{\frac{\sqrt{-a^2}}{2x^2}}}{x^3}$$

Which simplifies to

$$W = -\frac{2\sqrt{-a^2}}{x^3}$$

Therefore Eq. (2) becomes

$$u_1 = -\int \frac{\frac{e^{\frac{\sqrt{-a^2}}{2x^2}}}{x^2}}{-2x^3\sqrt{-a^2}} dx$$

Which simplifies to

$$u_1 = -\int -\frac{e^{\frac{\sqrt{-a^2}}{2x^2}}}{2x^5\sqrt{-a^2}} dx$$

Hence

$$u_1 = \frac{-\frac{e^{\frac{\sqrt{-a^2}}{2x^2}}}{x^2\sqrt{-a^2}} - \frac{2e^{\frac{\sqrt{-a^2}}{2x^2}}}{a^2}}{2\sqrt{-a^2}}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{e^{-\frac{\sqrt{-a^2}}{2x^2}}}{x^2}}{-2x^3\sqrt{-a^2}} dx$$

Which simplifies to

$$u_2 = \int -\frac{e^{-\frac{\sqrt{-a^2}}{2x^2}}}{2x^5\sqrt{-a^2}} dx$$

Hence

$$u_2 = -\frac{\frac{e^{-\frac{\sqrt{-a^2}}{2x^2}}}{x^2\sqrt{-a^2}} - \frac{2e^{-\frac{\sqrt{-a^2}}{2x^2}}}{a^2}}{2\sqrt{-a^2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(-\frac{e^{\frac{\sqrt{-a^2}}{2x^2}}}{x^2\sqrt{-a^2}} - \frac{2e^{\frac{\sqrt{-a^2}}{2x^2}}}{a^2} \right) e^{-\frac{\sqrt{-a^2}}{2x^2}}}{2\sqrt{-a^2}} - \frac{e^{\frac{\sqrt{-a^2}}{2x^2}} \left(\frac{e^{-\frac{\sqrt{-a^2}}{2x^2}}}{x^2\sqrt{-a^2}} - \frac{2e^{-\frac{\sqrt{-a^2}}{2x^2}}}{a^2} \right)}{2\sqrt{-a^2}}$$

Which simplifies to

$$y_p(x) = \frac{1}{a^2 x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{\sqrt{-a^2}}{2x^2}} + c_2 e^{\frac{\sqrt{-a^2}}{2x^2}} \right) + \left(\frac{1}{a^2 x^2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{1}{a^2 x^2} + c_1 e^{-\frac{\sqrt{-a^2}}{2x^2}} + c_2 e^{\frac{\sqrt{-a^2}}{2x^2}}$$

Solved as second order ode using change of variable on x method 1

Time used: 0.830 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^6$, $B = 3x^5$, $C = a^2$, $f(x) = \frac{1}{x^2}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^6 y'' + 3x^5 y' + a^2 y = 0$$

In normal form the ode

$$x^6 y'' + 3x^5 y' + a^2 y = 0 \tag{1}$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$\begin{aligned} p(x) &= \frac{3}{x} \\ q(x) &= \frac{a^2}{x^6} \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{\frac{a^2}{x^6}}}{c} \\ \tau'' &= -\frac{3a^2}{c\sqrt{\frac{a^2}{x^6}}x^7}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{3a^2}{c\sqrt{\frac{a^2}{x^6}}x^7} + \frac{3}{x}\frac{\sqrt{\frac{a^2}{x^6}}}{c}}{\left(\frac{\sqrt{\frac{a^2}{x^6}}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int \sqrt{\frac{a^2}{x^6}} dx}{c} \\ &= -\frac{x\sqrt{\frac{a^2}{x^6}}}{2c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) - c_2 \sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2\tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= \cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) \\ y_2 &= -\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) & -\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) \\ \frac{d}{dx}\left(\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\right) & \frac{d}{dx}\left(-\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) & -\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) \\ -\left(\frac{\sqrt{\frac{a^2}{x^6}}}{2} - \frac{3a^2}{2x^6\sqrt{\frac{a^2}{x^6}}}\right)\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) & -\left(\frac{\sqrt{\frac{a^2}{x^6}}}{2} - \frac{3a^2}{2x^6\sqrt{\frac{a^2}{x^6}}}\right)\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) \end{vmatrix}$$

Therefore

$$W = \left(\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\right) \left(-\left(\frac{\sqrt{\frac{a^2}{x^6}}}{2} - \frac{3a^2}{2x^6\sqrt{\frac{a^2}{x^6}}}\right)\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\right) \\ - \left(-\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\right) \left(-\left(\frac{\sqrt{\frac{a^2}{x^6}}}{2} - \frac{3a^2}{2x^6\sqrt{\frac{a^2}{x^6}}}\right)\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\right)$$

Which simplifies to

$$W = \frac{a^2 \left(\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)^2 + \sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)^2 \right)}{x^6 \sqrt{\frac{a^2}{x^6}}}$$

Which simplifies to

$$W = \frac{a^2}{x^6 \sqrt{\frac{a^2}{x^6}}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)}{\frac{x^2}{\frac{a^2}{\sqrt{\frac{a^2}{x^6}}}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) \sqrt{\frac{a^2}{x^6}}}{x^2 a^2} dx$$

Hence

$$u_1 = -\frac{4x^3\sqrt{\pi}\sqrt{\frac{a^2}{x^6}}\left(-\frac{x^7\left(\frac{a^2}{x^6}\right)^{3/2}\cos\left(\frac{a}{2x^2}\right)}{4\sqrt{\pi}a^2} + \frac{x^9\left(\frac{a^2}{x^6}\right)^{3/2}\sin\left(\frac{a}{2x^2}\right)}{2\sqrt{\pi}a^3}\right)}{a^4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)}{\frac{x^2}{\frac{a^2}{\sqrt{\frac{a^2}{x^6}}}}} dx$$

Which simplifies to

$$u_2 = \int \frac{\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\sqrt{\frac{a^2}{x^6}}}{x^2a^2} dx$$

Hence

$$u_2 = -\frac{4\sqrt{\pi}\sqrt{\frac{a^2}{x^6}}x^3\left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos\left(\frac{a}{2x^2}\right)}{2\sqrt{\pi}} + \frac{a\sin\left(\frac{a}{2x^2}\right)}{4\sqrt{\pi}x^2}\right)}{a^4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{4x^3\sqrt{\pi}\sqrt{\frac{a^2}{x^6}}\left(-\frac{x^7\left(\frac{a^2}{x^6}\right)^{3/2}\cos\left(\frac{a}{2x^2}\right)}{4\sqrt{\pi}a^2} + \frac{x^9\left(\frac{a^2}{x^6}\right)^{3/2}\sin\left(\frac{a}{2x^2}\right)}{2\sqrt{\pi}a^3}\right)\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)}{a^4} \\ + \frac{4\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\sqrt{\pi}\sqrt{\frac{a^2}{x^6}}x^3\left(-\frac{1}{2\sqrt{\pi}} + \frac{\cos\left(\frac{a}{2x^2}\right)}{2\sqrt{\pi}} + \frac{a\sin\left(\frac{a}{2x^2}\right)}{4\sqrt{\pi}x^2}\right)}{a^4}$$

Which simplifies to

$$y_p(x) = \frac{x^3\sqrt{\frac{a^2}{x^6}}\left(2x^2\cos\left(\frac{a}{2x^2}\right) + a\sin\left(\frac{a}{2x^2}\right) - 2x^2\right)\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) + a\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\left(-2x^2\sin\left(\frac{a}{2x^2}\right) + a\cos\left(\frac{a}{2x^2}\right)\right)}{x^2a^4}$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) - c_2\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\right) \\ + \left(\frac{x^3\sqrt{\frac{a^2}{x^6}}\left(2x^2\cos\left(\frac{a}{2x^2}\right) + a\sin\left(\frac{a}{2x^2}\right) - 2x^2\right)\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) + a\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\left(-2x^2\sin\left(\frac{a}{2x^2}\right) + a\cos\left(\frac{a}{2x^2}\right)\right)}{x^2a^4}\right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) - c_2\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) \\ + \frac{x^3\sqrt{\frac{a^2}{x^6}}\left(2x^2\cos\left(\frac{a}{2x^2}\right) + a\sin\left(\frac{a}{2x^2}\right) - 2x^2\right)\sin\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right) + a\cos\left(\frac{x\sqrt{\frac{a^2}{x^6}}}{2}\right)\left(-2x^2\sin\left(\frac{a}{2x^2}\right) + a\cos\left(\frac{a}{2x^2}\right)\right)}{x^2a^4}$$

Solved as second order Bessel ode

Time used: 0.270 (sec)

Writing the ode as

$$x^2 y'' + 3y'x + \frac{a^2 y}{x^4} = \frac{1}{x^6} \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + y'x + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= -1 \\ \beta &= \frac{a}{2} \\ n &= \frac{1}{2} \\ \gamma &= -2 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = \frac{2c_1 \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}} - \frac{2c_2 \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{2c_1 \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}} - \frac{2c_2 \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \frac{2 \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}} \\ y_2 &= -\frac{2 \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{2 \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} & -\frac{2 \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \\ \frac{d}{dx} \left(\frac{2 \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \right) & \frac{d}{dx} \left(-\frac{2 \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{2 \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} & -\frac{2 \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \\ -\frac{2 \sin\left(\frac{a}{2x^2}\right)}{x^2\sqrt{\pi}\sqrt{\frac{a}{x^2}}} + \frac{2 \sin\left(\frac{a}{2x^2}\right)a}{x^4\sqrt{\pi}\left(\frac{a}{x^2}\right)^{3/2}} - \frac{2a \cos\left(\frac{a}{2x^2}\right)}{x^4\sqrt{\pi}\sqrt{\frac{a}{x^2}}} & \frac{2 \cos\left(\frac{a}{2x^2}\right)}{x^2\sqrt{\pi}\sqrt{\frac{a}{x^2}}} - \frac{2 \cos\left(\frac{a}{2x^2}\right)a}{x^4\sqrt{\pi}\left(\frac{a}{x^2}\right)^{3/2}} - \frac{2a \sin\left(\frac{a}{2x^2}\right)}{x^4\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \end{vmatrix}$$

Therefore

$$W = \left(\frac{2 \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \right) \left(\frac{2 \cos\left(\frac{a}{2x^2}\right)}{x^2\sqrt{\pi}\sqrt{\frac{a}{x^2}}} - \frac{2 \cos\left(\frac{a}{2x^2}\right)a}{x^4\sqrt{\pi}\left(\frac{a}{x^2}\right)^{3/2}} - \frac{2a \sin\left(\frac{a}{2x^2}\right)}{x^4\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \right) \\ - \left(-\frac{2 \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \right) \left(-\frac{2 \sin\left(\frac{a}{2x^2}\right)}{x^2\sqrt{\pi}\sqrt{\frac{a}{x^2}}} + \frac{2 \sin\left(\frac{a}{2x^2}\right)a}{x^4\sqrt{\pi}\left(\frac{a}{x^2}\right)^{3/2}} - \frac{2a \cos\left(\frac{a}{2x^2}\right)}{x^4\sqrt{\pi}\sqrt{\frac{a}{x^2}}} \right)$$

Which simplifies to

$$W = -\frac{4\left(\cos\left(\frac{a}{2x^2}\right)^2 + \sin\left(\frac{a}{2x^2}\right)^2\right)}{x^3\pi}$$

Which simplifies to

$$W = -\frac{4}{x^3\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{2 \cos\left(\frac{a}{2x^2}\right)}{x^2\sqrt{\pi}\sqrt{\frac{a}{x^2}}}}{-\frac{4}{x\pi}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sqrt{\pi} \cos\left(\frac{a}{2x^2}\right)}{2x^6\sqrt{\frac{a}{x^2}}} dx$$

Hence

$$u_1 = \frac{\sqrt{\pi} \cos\left(\frac{a}{2x^2}\right)}{x a^2 \sqrt{\frac{a}{x^2}}} + \frac{\sqrt{\pi} \sin\left(\frac{a}{2x^2}\right)}{2a x^3 \sqrt{\frac{a}{x^2}}}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2 \sin\left(\frac{a}{2x^2}\right) x^7 \sqrt{\pi} \sqrt{\frac{a}{x^2}}}{-\frac{4}{x\pi}} dx$$

Which simplifies to

$$u_2 = \int -\frac{\sqrt{\pi} \sin\left(\frac{a}{2x^2}\right)}{2x^6 \sqrt{\frac{a}{x^2}}} dx$$

Hence

$$u_2 = -\frac{\sqrt{\pi} \cos\left(\frac{a}{2x^2}\right)}{2a x^3 \sqrt{\frac{a}{x^2}}} + \frac{\sqrt{\pi} \sin\left(\frac{a}{2x^2}\right)}{x a^2 \sqrt{\frac{a}{x^2}}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{2 \left(\frac{\sqrt{\pi} \cos\left(\frac{a}{2x^2}\right)}{x a^2 \sqrt{\frac{a}{x^2}}} + \frac{\sqrt{\pi} \sin\left(\frac{a}{2x^2}\right)}{2a x^3 \sqrt{\frac{a}{x^2}}} \right) \sin\left(\frac{a}{2x^2}\right) - 2 \cos\left(\frac{a}{2x^2}\right) \left(-\frac{\sqrt{\pi} \cos\left(\frac{a}{2x^2}\right)}{2a x^3 \sqrt{\frac{a}{x^2}}} + \frac{\sqrt{\pi} \sin\left(\frac{a}{2x^2}\right)}{x a^2 \sqrt{\frac{a}{x^2}}} \right)}{x \sqrt{\pi} \sqrt{\frac{a}{x^2}}}$$

Which simplifies to

$$y_p(x) = \frac{1}{a^2 x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{2c_1 \sin\left(\frac{a}{2x^2}\right)}{x \sqrt{\pi} \sqrt{\frac{a}{x^2}}} - \frac{2c_2 \cos\left(\frac{a}{2x^2}\right)}{x \sqrt{\pi} \sqrt{\frac{a}{x^2}}} \right) + \left(\frac{1}{a^2 x^2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{2c_1 \sin\left(\frac{a}{2x^2}\right)}{x \sqrt{\pi} \sqrt{\frac{a}{x^2}}} - \frac{2c_2 \cos\left(\frac{a}{2x^2}\right)}{x \sqrt{\pi} \sqrt{\frac{a}{x^2}}} + \frac{1}{a^2 x^2}$$

Solved as second order ode adjoint method

Time used: 2.273 (sec)

In normal form the ode

$$x^6 y'' + 3x^5 y' + a^2 y = \frac{1}{x^2} \tag{1}$$

Becomes

$$y'' + p(x) y' + q(x) y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= \frac{3}{x} \\ q(x) &= \frac{a^2}{x^6} \\ r(x) &= \frac{1}{x^8} \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned}\xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(\frac{3\xi(x)}{x}\right)' + \left(\frac{a^2\xi(x)}{x^6}\right) &= 0 \\ \xi''(x) - \frac{3\xi'(x)}{x} + \frac{(3x^4 + a^2)\xi(x)}{x^6} &= 0\end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$x^2\xi'' - 3\xi'x + \left(3 + \frac{a^2}{x^4}\right)\xi = 0 \quad (1)$$

Bessel ode has the form

$$x^2\xi'' + \xi'x + (-n^2 + x^2)\xi = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2\xi'' + (1 - 2\alpha)x\xi' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)\xi = 0 \quad (3)$$

With the standard solution

$$\xi = x^\alpha(c_3 \text{BesselJ}(n, \beta x^\gamma) + c_4 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= 2 \\ \beta &= \frac{a}{2} \\ n &= -\frac{1}{2} \\ \gamma &= -2\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$\xi = \frac{2c_3 x^2 \cos\left(\frac{a}{2x^2}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x^2}}} + \frac{2c_4 x^2 \sin\left(\frac{a}{2x^2}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x^2}}}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x)y' - y\xi'(x) + \xi(x)p(x)y &= \int \xi(x)r(x)dx \\ y' + y\left(p(x) - \frac{\xi'(x)}{\xi(x)}\right) &= \frac{\int \xi(x)r(x)dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y\left(\frac{3}{x} - \frac{\frac{4c_3x \cos\left(\frac{a}{2x^2}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x^2}}} + \frac{2c_3 \cos\left(\frac{a}{2x^2}\right)a}{x\sqrt{\pi}\left(\frac{a}{x^2}\right)^{3/2}} + \frac{2c_3a \sin\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}} + \frac{4c_4x \sin\left(\frac{a}{2x^2}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x^2}}} + \frac{2c_4 \sin\left(\frac{a}{2x^2}\right)a}{x\sqrt{\pi}\left(\frac{a}{x^2}\right)^{3/2}} - \frac{2c_4a \cos\left(\frac{a}{2x^2}\right)}{x\sqrt{\pi} \sqrt{\frac{a}{x^2}}}}{\frac{2c_3 x^2 \cos\left(\frac{a}{2x^2}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x^2}}} + \frac{2c_4 x^2 \sin\left(\frac{a}{2x^2}\right)}{\sqrt{\pi} \sqrt{\frac{a}{x^2}}}}\right) = \frac{2(-2c_3x^2 - c_3a) \sin\left(\frac{a}{2x^2}\right) + (-2c_4x^2 + c_4a) \cos\left(\frac{a}{2x^2}\right)}{a^2x^5 \left(c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)\right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{a(-\cos\left(\frac{a}{2x^2}\right)c_4 + \sin\left(\frac{a}{2x^2}\right)c_3)}{x^3(c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right))} \\ p(x) &= \frac{(-2c_4x^2 - c_3a) \sin\left(\frac{a}{2x^2}\right) + (-2c_3x^2 + c_4a) \cos\left(\frac{a}{2x^2}\right)}{a^2x^5(c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right))}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{a\left(-\cos\left(\frac{a}{2x^2}\right)c_4 + \sin\left(\frac{a}{2x^2}\right)c_3\right)}{x^3\left(c_3\cos\left(\frac{a}{2x^2}\right) + c_4\sin\left(\frac{a}{2x^2}\right)\right)} dx} \\ &= \frac{1}{c_3\cos\left(\frac{a}{2x^2}\right) + c_4\sin\left(\frac{a}{2x^2}\right)}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{(-2c_4 x^2 - c_3 a) \sin\left(\frac{a}{2x^2}\right) + (-2c_3 x^2 + c_4 a) \cos\left(\frac{a}{2x^2}\right)}{a^2 x^5 (c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right))} \right) \\ \frac{d}{dx} \left(\frac{y}{c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)} \right) &= \left(\frac{1}{c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)} \right) \left(\frac{(-2c_4 x^2 - c_3 a) \sin\left(\frac{a}{2x^2}\right) + (-2c_3 x^2 + c_4 a) \cos\left(\frac{a}{2x^2}\right)}{a^2 x^5 (c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right))} \right) \\ d \left(\frac{y}{c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)} \right) &= \left(\frac{(-2c_4 x^2 - c_3 a) \sin\left(\frac{a}{2x^2}\right) + (-2c_3 x^2 + c_4 a) \cos\left(\frac{a}{2x^2}\right)}{a^2 x^5 (c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right))^2} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)} &= \int \frac{(-2c_4 x^2 - c_3 a) \sin\left(\frac{a}{2x^2}\right) + (-2c_3 x^2 + c_4 a) \cos\left(\frac{a}{2x^2}\right)}{a^2 x^5 (c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right))^2} dx \\ &= \frac{2e^{\frac{ia}{2x^2}}}{a^2 x^2 \left(-ic_4 e^{\frac{ia}{x^2}} + e^{\frac{ia}{x^2}} c_3 + ic_4 + c_3\right)} + c_5\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)}$ gives the final solution

$$y = \frac{(c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)) \left(-2e^{\frac{ia}{2x^2}} + a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2 c_5\right)}{a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{(c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)) \left(-2e^{\frac{ia}{2x^2}} + a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2 c_5\right)}{a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2}$$

The constants can be merged to give

$$y = \frac{(c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)) \left(-2e^{\frac{ia}{2x^2}} + a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2\right)}{a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{(c_3 \cos\left(\frac{a}{2x^2}\right) + c_4 \sin\left(\frac{a}{2x^2}\right)) \left(-2e^{\frac{ia}{2x^2}} + a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2\right)}{a^2 \left((ic_4 - c_3) e^{\frac{ia}{x^2}} - ic_4 - c_3\right) x^2}$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    <- linear_1 successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.008 (sec)

Leaf size : 30

```

dsolve(diff(diff(y(x),x),x)*x^6+3*x^5*diff(y(x),x)+a^2*y(x) = 1/x^2,
        y(x),singsol=all)

```

$$y = \sin\left(\frac{a}{2x^2}\right) c_2 + \cos\left(\frac{a}{2x^2}\right) c_1 + \frac{1}{a^2 x^2}$$

Mathematica DSolve solution

Solving time : 0.092 (sec)

Leaf size : 38

```

DSolve[{x^6*D[y[x],{x,2}]+3*x^5*D[y[x],x]+a^2*y[x]==1/x^2,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{1}{a^2 x^2} + c_1 \cos\left(\frac{a}{2x^2}\right) - c_2 \sin\left(\frac{a}{2x^2}\right)$$

2.2.19 problem 20

Solved as second order Euler type ode	620
Solved as second order ode using change of variable on x method 2	623
Solved as second order ode using change of variable on x method 1	629
Solved as second order ode using change of variable on y method 2	632
Solved as second order ode using non constant coeff trans- formation on B method	636
Solved as second order ode using Kovacic algorithm	639
Solved as second order ode adjoint method	645
Maple step by step solution	647
Maple trace	647
Maple dsolve solution	647
Mathematica DSolve solution	647

Internal problem ID [8818]

Book : Second order enumerated odes

Section : section 2

Problem number : 20

Date solved : Thursday, December 12, 2024 at 09:52:03 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$x^2y'' - 3xy' + 3y = 2x^3 - x^2$$

Solved as second order Euler type ode

Time used: 0.178 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -3x$, $C = 3$, $f(x) = 2x^3 - x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 3xy' + 3y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 3rx^{r-1} + 3x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 3rx^r + 3x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - 3r + 3 = 0$$

Or

$$r^2 - 4r + 3 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 1$$

$$r_2 = 3$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_2 x^3 + c_1 x$$

Next, we find the particular solution to the ODE

$$x^2 y'' - 3xy' + 3y = 2x^3 - x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = x^3$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^3 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^3) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^3 \\ 1 & 3x^2 \end{vmatrix}$$

Therefore

$$W = (x)(3x^2) - (x^3)(1)$$

Which simplifies to

$$W = 2x^3$$

Which simplifies to

$$W = 2x^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3(2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_1 = - \int \left(x - \frac{1}{2} \right) dx$$

Hence

$$u_1 = -\frac{1}{2}x^2 + \frac{1}{2}x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_2 = \int \frac{2x - 1}{2x^2} dx$$

Hence

$$u_2 = \ln(x) + \frac{1}{2x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{1}{2}x^2 + \frac{1}{2}x \right) x + x^3 \left(\ln(x) + \frac{1}{2x} \right)$$

Which simplifies to

$$y_p(x) = x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= c_2 x^3 + c_1 x + x^3 \ln(x) - \frac{x^3}{2} + x^2 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2 x^3 + c_1 x + x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Solved as second order ode using change of variable on x method 2

Time used: 0.663 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' - 3xy' + 3y = 0$$

In normal form the ode

$$x^2y'' - 3xy' + 3y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$

$$q(x) = \frac{3}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\int p(x)dx} dx \\ &= \int e^{-\int -\frac{3}{x}dx} dx \\ &= \int e^{3\ln(x)} dx \\ &= \int x^3 dx \\ &= \frac{x^4}{4} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{3}{x^2}}{x^6} \\ &= \frac{3}{x^8} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{3y(\tau)}{x^8} &= 0\end{aligned}$$

But in terms of τ

$$\frac{3}{x^8} = \frac{3}{16\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{3y(\tau)}{16\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. Writing the ode as

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{3y(\tau)}{16\tau^2} = 0 \quad (1)$$

$$A\frac{d^2}{d\tau^2}y(\tau) + B\frac{d}{d\tau}y(\tau) + Cy(\tau) = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\ B &= 0 \\ C &= \frac{3}{16\tau^2}\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(\tau) = y(\tau) e^{\int \frac{B}{2A} d\tau}$$

Then (2) becomes

$$z''(\tau) = rz(\tau) \quad (4)$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{16\tau^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -3 \\ t &= 16\tau^2\end{aligned}$$

Therefore eq. (4) becomes

$$z''(\tau) = \left(-\frac{3}{16\tau^2}\right)z(\tau) \quad (7)$$

Equation (7) is now solved. After finding $z(\tau)$ then $y(\tau)$ is found using the inverse transformation

$$y(\tau) = z(\tau) e^{-\int \frac{B}{2A} d\tau}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.77: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 16\tau^2$. There is a pole at $\tau = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{3}{16\tau^2}$$

For the pole at $\tau = 0$ let b be the coefficient of $\frac{1}{\tau^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{3}{16}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{4} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{4} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{\tau^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{3}{16\tau^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{3}{16}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{4} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{4} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{3}{16\tau^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{4}$	$\frac{1}{4}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{4}$	$\frac{1}{4}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{4}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{4} - \left(\frac{1}{4}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{\tau - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{\tau - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= \frac{1}{4\tau} + (-)(0) \\ &= \frac{1}{4\tau} \\ &= \frac{1}{4\tau} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(\tau)$ of degree $d = 0$ to solve the ode. The polynomial $p(\tau)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(\tau) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{4\tau}\right)(0) + \left(\left(-\frac{1}{4\tau^2}\right) + \left(\frac{1}{4\tau}\right)^2 - \left(-\frac{3}{16\tau^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(\tau) &= pe^{\int \omega d\tau} \\ &= e^{\int \frac{1}{4\tau} d\tau} \\ &= \tau^{1/4} \end{aligned}$$

The first solution to the original ode in $y(\tau)$ is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} d\tau}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \tau^{1/4} \end{aligned}$$

Which simplifies to

$$y_1 = \tau^{1/4}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} d\tau}}{y_1^2} d\tau$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} d\tau \\ &= \tau^{1/4} \int \frac{1}{\sqrt{\tau}} d\tau \\ &= \tau^{1/4} (2\sqrt{\tau}) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y(\tau) &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\tau^{1/4}) + c_2 (\tau^{1/4} (2\sqrt{\tau})) \end{aligned}$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 4^{3/4} (x^4)^{1/4}}{4} + \frac{c_2 4^{1/4} (x^4)^{3/4}}{2}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 4^{3/4} (x^4)^{1/4}}{4} + \frac{c_2 4^{1/4} (x^4)^{3/4}}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= (x^4)^{1/4} \\ y_2 &= (x^4)^{3/4} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x^4)^{1/4} & (x^4)^{3/4} \\ \frac{d}{dx}((x^4)^{1/4}) & \frac{d}{dx}((x^4)^{3/4}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x^4)^{1/4} & (x^4)^{3/4} \\ \frac{x^3}{(x^4)^{3/4}} & \frac{3x^3}{(x^4)^{1/4}} \end{vmatrix}$$

Therefore

$$W = ((x^4)^{1/4}) \left(\frac{3x^3}{(x^4)^{1/4}} \right) - ((x^4)^{3/4}) \left(\frac{x^3}{(x^4)^{3/4}} \right)$$

Which simplifies to

$$W = 2x^3$$

Which simplifies to

$$W = 2x^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(x^4)^{3/4} (2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_1 = - \int \frac{(x^4)^{3/4} (2x - 1)}{2x^3} dx$$

Hence

$$u_1 = - \frac{(x - 1)(x^4)^{3/4}}{2x^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x^4)^{1/4} (2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_2 = \int \frac{(x^4)^{1/4} (2x - 1)}{2x^3} dx$$

Hence

$$u_2 = \frac{(x^4)^{1/4}}{2x^2} + \frac{(x^4)^{1/4} \ln(x)}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(x-1)}{2} + (x^4)^{3/4} \left(\frac{(x^4)^{1/4}}{2x^2} + \frac{(x^4)^{1/4} \ln(x)}{x} \right)$$

Which simplifies to

$$y_p(x) = x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1 4^{3/4} (x^4)^{1/4}}{4} + \frac{c_2 4^{1/4} (x^4)^{3/4}}{2} \right) + \left(x^3 \ln(x) - \frac{x^3}{2} + x^2 \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^3 \ln(x) - \frac{x^3}{2} + x^2 + \frac{c_1 4^{3/4} (x^4)^{1/4}}{4} + \frac{c_2 4^{1/4} (x^4)^{3/4}}{2}$$

Solved as second order ode using change of variable on x method 1

Time used: 0.244 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -3x$, $C = 3$, $f(x) = 2x^3 - x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' - 3xy' + 3y = 0$$

In normal form the ode

$$x^2 y'' - 3xy' + 3y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$\begin{aligned} p(x) &= -\frac{3}{x} \\ q(x) &= \frac{3}{x^2} \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{3}\sqrt{\frac{1}{x^2}}}{c} \\ \tau'' &= -\frac{\sqrt{3}}{c\sqrt{\frac{1}{x^2}}x^3} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{\sqrt{3}}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{3}{x}\frac{\sqrt{3}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{3}\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\ &= -\frac{4c\sqrt{3}}{3} \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - \frac{4c\sqrt{3}}{3}\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{\frac{2\sqrt{3}c\tau}{3}} \left(c_1 \cosh\left(\frac{\sqrt{3}c\tau}{3}\right) + ic_2 \sinh\left(\frac{\sqrt{3}c\tau}{3}\right) \right)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int \sqrt{3}\sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{3} \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = x^2 \left(c_1 \left(\frac{x}{2} + \frac{1}{2x} \right) + ic_2 \left(\frac{x}{2} - \frac{1}{2x} \right) \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^2 \left(\frac{x}{2} + \frac{1}{2x} \right)$$

$$y_2 = ix^2 \left(\frac{x}{2} - \frac{1}{2x} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^2 \left(\frac{x}{2} + \frac{1}{2x} \right) & ix^2 \left(\frac{x}{2} - \frac{1}{2x} \right) \\ \frac{d}{dx} \left(x^2 \left(\frac{x}{2} + \frac{1}{2x} \right) \right) & \frac{d}{dx} \left(ix^2 \left(\frac{x}{2} - \frac{1}{2x} \right) \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^2 \left(\frac{x}{2} + \frac{1}{2x} \right) & ix^2 \left(\frac{x}{2} - \frac{1}{2x} \right) \\ 2x \left(\frac{x}{2} + \frac{1}{2x} \right) + x^2 \left(\frac{1}{2} - \frac{1}{2x^2} \right) & 2ix \left(\frac{x}{2} - \frac{1}{2x} \right) + ix^2 \left(\frac{1}{2} + \frac{1}{2x^2} \right) \end{vmatrix}$$

Therefore

$$W = \left(x^2 \left(\frac{x}{2} + \frac{1}{2x} \right) \right) \left(2ix \left(\frac{x}{2} - \frac{1}{2x} \right) + ix^2 \left(\frac{1}{2} + \frac{1}{2x^2} \right) \right) - \left(ix^2 \left(\frac{x}{2} - \frac{1}{2x} \right) \right) \left(2x \left(\frac{x}{2} + \frac{1}{2x} \right) + x^2 \left(\frac{1}{2} - \frac{1}{2x^2} \right) \right)$$

Which simplifies to

$$W = ix^3$$

Which simplifies to

$$W = ix^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{ix^2 \left(\frac{x}{2} - \frac{1}{2x} \right) (2x^3 - x^2)}{ix^5} dx$$

Which simplifies to

$$u_1 = - \int \frac{(x^2 - 1)(2x - 1)}{2x^2} dx$$

Hence

$$u_1 = -\frac{x^2}{2} + \frac{x}{2} + \ln(x) + \frac{1}{2x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^2 \left(\frac{x}{2} + \frac{1}{2x}\right) (2x^3 - x^2)}{ix^5} dx$$

Which simplifies to

$$u_2 = \int -\frac{i(x^2 + 1)(2x - 1)}{2x^2} dx$$

Hence

$$u_2 = -\frac{i(x^2 - x + 2 \ln(x) + \frac{1}{x})}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{x^2}{2} + \frac{x}{2} + \ln(x) + \frac{1}{2x}\right) x^2 \left(\frac{x}{2} + \frac{1}{2x}\right) + \frac{x^2 \left(\frac{x}{2} - \frac{1}{2x}\right) (x^2 - x + 2 \ln(x) + \frac{1}{x})}{2}$$

Which simplifies to

$$y_p(x) = x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(x^2 \left(c_1 \left(\frac{x}{2} + \frac{1}{2x}\right) + ic_2 \left(\frac{x}{2} - \frac{1}{2x}\right)\right)\right) + \left(x^3 \ln(x) - \frac{x^3}{2} + x^2\right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^2 \left(c_1 \left(\frac{x}{2} + \frac{1}{2x}\right) + ic_2 \left(\frac{x}{2} - \frac{1}{2x}\right)\right) + x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Solved as second order ode using change of variable on y method 2

Time used: 0.253 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -3x$, $C = 3$, $f(x) = 2x^3 - x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' - 3xy' + 3y = 0$$

In normal form the ode

$$x^2 y'' - 3xy' + 3y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$

$$q(x) = \frac{3}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{3n}{x^2} + \frac{3}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 3 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{3v'(x)}{x} = 0$$

$$v''(x) + \frac{3v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{3u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form a linear first order is

$$u'(x) + q(x)u(x) = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = \frac{3}{x}$$

$$p(x) = 0$$

The integrating factor μ is

$$\mu = e^{\int q dx}$$

$$= e^{\int \frac{3}{x} dx}$$

$$= x^3$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu u &= 0 \\ \frac{d}{dx}(u x^3) &= 0\end{aligned}$$

Integrating gives

$$\begin{aligned}u x^3 &= \int 0 dx + c_1 \\ &= c_1\end{aligned}$$

Dividing throughout by the integrating factor x^3 gives the final solution

$$u(x) = \frac{c_1}{x^3}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{2x^2} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{2x^2} + c_2\right) x^3 \\ &= c_2 x^3 - \frac{1}{2}c_1 x\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' - 3xy' + 3y = 2x^3 - x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= x \\ y_2 &= x^3\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^3 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^3) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^3 \\ 1 & 3x^2 \end{vmatrix}$$

Therefore

$$W = (x)(3x^2) - (x^3)(1)$$

Which simplifies to

$$W = 2x^3$$

Which simplifies to

$$W = 2x^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3(2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_1 = - \int \left(x - \frac{1}{2} \right) dx$$

Hence

$$u_1 = -\frac{1}{2}x^2 + \frac{1}{2}x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_2 = \int \frac{2x - 1}{2x^2} dx$$

Hence

$$u_2 = \ln(x) + \frac{1}{2x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{1}{2}x^2 + \frac{1}{2}x \right) x + x^3 \left(\ln(x) + \frac{1}{2x} \right)$$

Which simplifies to

$$y_p(x) = x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\left(-\frac{c_1}{2x^2} + c_2 \right) x^3 \right) + \left(x^3 \ln(x) - \frac{x^3}{2} + x^2 \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(-\frac{c_1}{2x^2} + c_2 \right) x^3 + x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Solved as second order ode using non constant coeff transformation on B method

Time used: 0.118 (sec)

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned} A &= x^2 \\ B &= -3x \\ C &= 3 \\ F &= 2x^3 - x^2 \end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x^2)(0) + (-3x)(-3) + (3)(-3x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-3x^3v'' + (3x^2)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-3x^2(u'(x)x - u(x)) = 0$$

Which is now solved for u . In canonical form a linear first order is

$$u'(x) + q(x)u(x) = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{1}{x}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu u &= 0 \\ \frac{d}{dx}\left(\frac{u}{x}\right) &= 0\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{u}{x} &= \int 0 dx + c_1 \\ &= c_1\end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{x}$ gives the final solution

$$u(x) = c_1x$$

The ode for v now becomes

$$v'(x) = c_1x$$

Which is now solved for v . Since the ode has the form $v'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned}\int dv &= \int c_1x dx \\ v(x) &= \frac{c_1x^2}{2} + c_2\end{aligned}$$

Replacing $v(x)$ above by $-\frac{y}{3x}$, then the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\ &= -\frac{3(c_1x^2 + 2c_2)x}{2}\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of

parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = x^3$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^3 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^3) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^3 \\ 1 & 3x^2 \end{vmatrix}$$

Therefore

$$W = (x)(3x^2) - (x^3)(1)$$

Which simplifies to

$$W = 2x^3$$

Which simplifies to

$$W = 2x^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3(2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_1 = - \int \left(x - \frac{1}{2} \right) dx$$

Hence

$$u_1 = -\frac{1}{2}x^2 + \frac{1}{2}x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(2x^3 - x^2)}{2x^5} dx$$

Which simplifies to

$$u_2 = \int \frac{2x - 1}{2x^2} dx$$

Hence

$$u_2 = \ln(x) + \frac{1}{2x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{1}{2}x^2 + \frac{1}{2}x\right)x + x^3\left(\ln(x) + \frac{1}{2x}\right)$$

Which simplifies to

$$y_p(x) = x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Hence the complete solution is

$$\begin{aligned} y(x) &= y_h + y_p \\ &= \left(-\frac{3(c_1 x^2 + 2c_2)x}{2}\right) + \left(x^3 \ln(x) - \frac{x^3}{2} + x^2\right) \\ &= x^3 \ln(x) + \frac{(-3c_1 - 1)x^3}{2} + x^2 - 3c_2x \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^3 \ln(x) + \frac{(-3c_1 - 1)x^3}{2} + x^2 - 3c_2x$$

Solved as second order ode using Kovacic algorithm

Time used: 0.230 (sec)

Writing the ode as

$$x^2 y'' - 3xy' + 3y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= -3x \\ C &= 3 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 3 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.78: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{3}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-)(0) \\ &= -\frac{1}{2x} \\ &= -\frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2x}\right)(0) + \left(\left(\frac{1}{2x^2}\right) + \left(-\frac{1}{2x}\right)^2 - \left(\frac{3}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{2x} dx} \\ &= \frac{1}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-3x}{x^2} dx} \\ &= z_1 e^{\frac{3 \ln(x)}{2}} \\ &= z_1 (x^{3/2}) \end{aligned}$$

Which simplifies to

$$y_1 = x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-3x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{3 \ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(\frac{x^2}{2}\right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(x) + c_2 \left(x \left(\frac{x^2}{2}\right)\right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' - 3xy' + 3y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1x + \frac{1}{2}c_2x^3$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = \frac{x^3}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & \frac{x^3}{2} \\ \frac{d}{dx}(x) & \frac{d}{dx}\left(\frac{x^3}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & \frac{x^3}{2} \\ 1 & \frac{3x^2}{2} \end{vmatrix}$$

Therefore

$$W = (x) \left(\frac{3x^2}{2} \right) - \left(\frac{x^3}{2} \right) \quad (1)$$

Which simplifies to

$$W = x^3$$

Which simplifies to

$$W = x^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3(2x^3-x^2)}{x^5} dx$$

Which simplifies to

$$u_1 = - \int \left(x - \frac{1}{2} \right) dx$$

Hence

$$u_1 = -\frac{1}{2}x^2 + \frac{1}{2}x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(2x^3-x^2)}{x^5} dx$$

Which simplifies to

$$u_2 = \int \frac{2x-1}{x^2} dx$$

Hence

$$u_2 = 2 \ln(x) + \frac{1}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{1}{2}x^2 + \frac{1}{2}x \right) x + \frac{x^3(2 \ln(x) + \frac{1}{x})}{2}$$

Which simplifies to

$$y_p(x) = x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1x + \frac{1}{2}c_2x^3 \right) + \left(x^3 \ln(x) - \frac{x^3}{2} + x^2 \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1x + \frac{c_2x^3}{2} + x^3 \ln(x) - \frac{x^3}{2} + x^2$$

Solved as second order ode adjoint method

Time used: 0.209 (sec)

In normal form the ode

$$x^2 y'' - 3xy' + 3y = 2x^3 - x^2 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{3}{x} \\ q(x) &= \frac{3}{x^2} \\ r(x) &= 2x - 1 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(-\frac{3\xi(x)}{x}\right)' + \left(\frac{3\xi(x)}{x^2}\right) &= 0 \\ \xi''(x) + \frac{3\xi'(x)}{x} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is second order ode with missing dependent variable ξ . Let

$$p(x) = \xi'$$

Then

$$p'(x) = \xi''$$

Hence the ode becomes

$$p'(x) + \frac{3p(x)}{x} = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form a linear first order is

$$p'(x) + q(x)p(x) = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= \frac{3}{x} \\ p(x) &= 0 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int \frac{3}{x} dx} \\ &= x^3 \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx} \mu p &= 0 \\ \frac{d}{dx} (p x^3) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned} p x^3 &= \int 0 dx + c_1 \\ &= c_1 \end{aligned}$$

Dividing throughout by the integrating factor x^3 gives the final solution

$$p(x) = \frac{c_1}{x^3}$$

For solution (1) found earlier, since $p = \xi'$ then we now have a new first order ode to solve which is

$$\xi' = \frac{c_1}{x^3}$$

Since the ode has the form $\xi' = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned} \int d\xi &= \int \frac{c_1}{x^3} dx \\ \xi &= -\frac{c_1}{2x^2} + c_2 \end{aligned}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(-\frac{3}{x} - \frac{c_1}{x^3 \left(-\frac{c_1}{2x^2} + c_2 \right)} \right) = \frac{c_2 x^2 - c_2 x - c_1 \ln(x) - \frac{c_1}{2x}}{-\frac{c_1}{2x^2} + c_2}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{-6c_2 x^2 + c_1}{x(-2c_2 x^2 + c_1)} \\ p(x) &= \frac{x(-2c_2 x^3 + 2c_1 \ln(x) x + 2c_2 x^2 + c_1)}{-2c_2 x^2 + c_1} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{-6c_2 x^2 + c_1}{x(-2c_2 x^2 + c_1)} dx} \\ &= \frac{1}{x(-2c_2 x^2 + c_1)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{x(-2c_2 x^3 + 2c_1 \ln(x) x + 2c_2 x^2 + c_1)}{-2c_2 x^2 + c_1} \right) \\ \frac{d}{dx} \left(-\frac{y}{x(-2c_2 x^2 + c_1)} \right) &= \left(-\frac{1}{x(-2c_2 x^2 + c_1)} \right) \left(\frac{x(-2c_2 x^3 + 2c_1 \ln(x) x + 2c_2 x^2 + c_1)}{-2c_2 x^2 + c_1} \right) \\ d \left(-\frac{y}{x(-2c_2 x^2 + c_1)} \right) &= \left(-\frac{-2c_2 x^3 + 2c_1 \ln(x) x + 2c_2 x^2 + c_1}{(-2c_2 x^2 + c_1)^2} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} -\frac{y}{x(-2c_2x^2 + c_1)} &= \int \frac{-2c_2x^3 + 2c_1 \ln(x)x + 2c_2x^2 + c_1}{(-2c_2x^2 + c_1)^2} dx \\ &= \frac{\frac{x}{2} - \frac{c_1}{8c_2}}{-\frac{c_1}{2} + c_2x^2} + \frac{\ln(x)x^2}{2c_2x^2 - c_1} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $-\frac{1}{x(-2c_2x^2 + c_1)}$ gives the final solution

$$y = \frac{x(8c_2^2c_3x^2 + 4x^2 \ln(x)c_2 - 4c_1c_2c_3 + 4c_2x - c_1)}{4c_2}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{x(8c_2^2c_3x^2 + 4x^2 \ln(x)c_2 - 4c_1c_2c_3 + 4c_2x - c_1)}{4c_2}$$

The constants can be merged to give

$$y = \frac{x(8c_2^2x^2 + 4x^2 \ln(x)c_2 - 4c_1c_2 + 4c_2x - c_1)}{4c_2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x(8c_2^2x^2 + 4x^2 \ln(x)c_2 - 4c_1c_2 + 4c_2x - c_1)}{4c_2}$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful`
```

Maple dsolve solution

Solving time : 0.005 (sec)

Leaf size : 28

```
dsolve(x^2*diff(diff(y(x),x),x)-3*diff(y(x),x)*x+3*y(x) = 2*x^3-x^2,
y(x),singsol=all)
```

$$y = \frac{x(2 \ln(x)x^2 + (c_1 - 1)x^2 + 2x + 2c_2)}{2}$$

Mathematica DSolve solution

Solving time : 0.026 (sec)

Leaf size : 27

```
DSolve[{x^2*D[y[x],{x,2}]-3*x*D[y[x],x]+3*y[x]==2*x^3-x^2,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x \left(x^2 \log(x) + \left(-\frac{1}{2} + c_2 \right) x^2 + x + c_1 \right)$$

2.2.20 problem 21

Solved as second order ode using change of variable on x
 method 2 648
 Maple step by step solution 650
 Maple trace 650
 Maple dsolve solution 650
 Mathematica DSolve solution 650

Internal problem ID [8819]

Book : Second order enumerated odes

Section : section 2

Problem number : 21

Date solved : Thursday, December 12, 2024 at 09:52:05 AM

CAS classification :

[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear, ‘_with_symmetry_[0,F(x)]’]

Solve

$$y'' + \cot(x) y' + 4y \csc(x)^2 = 0$$

Solved as second order ode using change of variable on x method 2

Time used: 0.506 (sec)

In normal form the ode

$$y'' + \cot(x) y' + 4y \csc(x)^2 = 0 \tag{1}$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$p(x) = \cot(x)$$

$$q(x) = 4 \csc(x)^2$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\int p(x)dx} dx \\ &= \int e^{-\int \cot(x)dx} dx \\ &= \int e^{-\ln(\sin(x))} dx \\ &= \int \csc(x) dx \\ &= -\ln(\csc(x) + \cot(x)) \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{4 \csc(x)^2}{\csc(x)^2} \\ &= 4 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + 4y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\tau\lambda} + 4 e^{\tau\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +2i \\ \lambda_2 &= -2i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2i \\ \lambda_2 &= -2i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau} (c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0 (c_1 \cos(2\tau) + c_2 \sin(2\tau))$$

Or

$$y(\tau) = c_1 \cos(2\tau) + c_2 \sin(2\tau)$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(2 \ln(\csc(x) + \cot(x))) - c_2 \sin(2 \ln(\csc(x) + \cot(x)))$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos(2 \ln(\csc(x) + \cot(x))) - c_2 \sin(2 \ln(\csc(x) + \cot(x)))$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`
```

Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 23

```
dsolve(diff(diff(y(x),x),x)+cot(x)*diff(y(x),x)+4*y(x)*csc(x)^2 = 0,
        y(x),singsol=all)
```

$$y = c_1(\csc(x) + \cot(x))^{-2i} + c_2(\csc(x) + \cot(x))^{2i}$$

Mathematica DSolve solution

Solving time : 0.071 (sec)

Leaf size : 25

```
DSolve[{D[y[x],{x,2}]+Cot[x]*D[y[x],x]+4*y[x]*Csc[x]^2==0,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_1 \cos(2\arctanh(\cos(x))) - c_2 \sin(2\arctanh(\cos(x)))$$

2.2.21 problem 22

Maple step by step solution	651
Maple trace	651
Maple dsolve solution	652
Mathematica DSolve solution	652

Internal problem ID [8820]

Book : Second order enumerated odes

Section : section 2

Problem number : 22

Date solved : Wednesday, December 18, 2024 at 02:16:39 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$(x^2 + 1) y'' + (1 + x) y' + y = 4 \cos(\ln(1 + x))$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
        -> heuristic approach
            <- heuristic approach successful
        <- hypergeometric successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.108 (sec)

Leaf size : 280

```
dsolve((x^2+1)*diff(diff(y(x),x),x)+(x+1)*diff(y(x),x)+y(x) = 4*cos(ln(x+1))),
      y(x),singsol=all)
```

$$\begin{aligned}
y = & \text{hypergeom} \left([i, -i], \left[\frac{1}{2} + \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right) c_2 \\
& + (x+i)^{\frac{1}{2}-\frac{i}{2}} \text{hypergeom} \left(\left[\frac{1}{2} + \frac{i}{2}, \frac{1}{2} - \frac{3i}{2} \right], \left[\frac{3}{2} - \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right) c_1 \\
& - 80 \left(\int \frac{10 \text{hypergeom} \left(\left[\frac{1}{2} + \frac{i}{2}, \frac{1}{2} - \frac{3i}{2} \right], \left[\frac{3}{2} - \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right) ((-1-i + (-1+i)x) \text{hypergeom} \left([i, -i], \left[\frac{1}{2} + \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right) \right.}{(x^2+1)} \right. \\
& \left. - 80 \left(\int \frac{\text{hypergeom} \left([i, -i], \left[\frac{1}{2} + \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right)}{7 \left(\frac{10((1-i+(-1-i)x) \text{hypergeom} \left([1-i, 1+i], \left[\frac{3}{2} + \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right) + (-1+i) \text{hypergeom} \left([i, -i], \left[\frac{1}{2} + \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right)) \text{hypergeom} \left(\left[\frac{1}{2} + \frac{i}{2}, \frac{1}{2} - \frac{3i}{2} \right], \left[\frac{3}{2} - \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right) \right.}{7} \right. \right. \\
& \left. \left. + i)^{\frac{1}{2}-\frac{i}{2}} \text{hypergeom} \left(\left[\frac{1}{2} + \frac{i}{2}, \frac{1}{2} - \frac{3i}{2} \right], \left[\frac{3}{2} - \frac{i}{2} \right], \frac{1}{2} - \frac{ix}{2} \right) \right) \right)
\end{aligned}$$

Mathematica DSolve solution

Solving time : 0.0 (sec)

Leaf size : 0

```
DSolve[{(1+x^2)*D[y[x],{x,2}]+(1+x)*D[y[x],x]+y[x]==4*Cos[Log[1+x]],{}}],
      y[x],x,IncludeSingularSolutions->True]
```

Not solved

2.2.22 problem 23

Solved as second order ode using change of variable on x

method 2	653
Maple step by step solution	655
Maple trace	655
Maple dsolve solution	655
Mathematica DSolve solution	655

Internal problem ID [8821]

Book : Second order enumerated odes**Section** : section 2**Problem number** : 23**Date solved** : Thursday, December 12, 2024 at 09:52:15 AM**CAS classification** : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' + \tan(x) y' + \cos(x)^2 y = 0$$

Solved as second order ode using change of variable on x method 2

Time used: 0.397 (sec)

In normal form the ode

$$y'' + \tan(x) y' + \cos(x)^2 y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= \tan(x) \\ q(x) &= \cos(x)^2 \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\int p(x) dx} dx \\ &= \int e^{-\int \tan(x) dx} dx \\ &= \int e^{\ln(\cos(x))} dx \\ &= \int \cos(x) dx \\ &= \sin(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\cos(x)^2}{\cos(x)^2} \\ &= 1 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} + e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau} (c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0 (c_1 \cos(\tau) + c_2 \sin(\tau))$$

Or

$$y(\tau) = c_1 \cos(\tau) + c_2 \sin(\tau)$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(\sin(x)) + c_2 \sin(\sin(x))$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos(\sin(x)) + c_2 \sin(\sin(x))$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebiu
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) *
-> Trying changes of variables to rationalize or make the ODE simpler
    trying a quadrature
    checking if the LODE has constant coefficients
    <- constant coefficients successful
Change of variables used:
    [x = arcsin(t)]
Linear ODE actually solved:
    (-2*t^2+2)*u(t)+(-2*t^2+2)*diff(diff(u(t),t),t) = 0
<- change of variables successful`

```

Maple dsolve solution

Solving time : 0.105 (sec)

Leaf size : 15

```

dsolve(diff(diff(y(x),x),x)+tan(x)*diff(y(x),x)+cos(x)^2*y(x) = 0,
        y(x),singsol=all)

```

$$y = c_1 \sin(\sin(x)) + c_2 \cos(\sin(x))$$

Mathematica DSolve solution

Solving time : 0.093 (sec)

Leaf size : 18

```

DSolve[{D[y[x],{x,2}]+Tan[x]*D[y[x],x]+Cos[x]^2*y[x]==0,{x]},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow c_2 \sin(\sin(x)) + c_1 \cos(\sin(x))$$

2.2.23 problem 24

Solved as second order ode using change of variable on x
 method 2 656
 Solved as second order ode using change of variable on x
 method 1 661
 Solved as second order Bessel ode 665
 Solved as second order ode using Kovacic algorithm 668
 Solved as second order ode adjoint method 675
 Maple step by step solution 677
 Maple trace 677
 Maple dsolve solution 678
 Mathematica DSolve solution 678

Internal problem ID [8822]

Book : Second order enumerated odes

Section : section 2

Problem number : 24

Date solved : Thursday, December 12, 2024 at 09:52:18 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$xy'' - y' + 4x^3y = 8x^3 \sin(x)^2$$

Solved as second order ode using change of variable on x method 2

Time used: 9.489 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$xy'' - y' + 4x^3y = 0$$

In normal form the ode

$$xy'' - y' + 4x^3y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{1}{x}$$

$$q(x) = 4x^2$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\int p(x) dx} dx \\ &= \int e^{-\int -\frac{1}{x} dx} dx \\ &= \int e^{\ln(x)} dx \\ &= \int x dx \\ &= \frac{x^2}{2} \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{4x^2}{x^2} \\ &= 4 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2} y(\tau) + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + 4y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} + 4e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +2i \\ \lambda_2 &= -2i \end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 2i \\ \lambda_2 &= -2i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau}(c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0(c_1 \cos(2\tau) + c_2 \sin(2\tau))$$

Or

$$y(\tau) = c_1 \cos(2\tau) + c_2 \sin(2\tau)$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(x^2) + c_2 \sin(x^2)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x^2) + c_2 \sin(x^2)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x^2)$$

$$y_2 = \sin(x^2)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x^2) & \sin(x^2) \\ \frac{d}{dx}(\cos(x^2)) & \frac{d}{dx}(\sin(x^2)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x^2) & \sin(x^2) \\ -2x \sin(x^2) & 2x \cos(x^2) \end{vmatrix}$$

Therefore

$$W = (\cos(x^2))(2x \cos(x^2)) - (\sin(x^2))(-2x \sin(x^2))$$

Which simplifies to

$$W = 2 \cos(x^2)^2 x + 2 \sin(x^2)^2 x$$

Which simplifies to

$$W = 2x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{8 \sin(x^2) x^3 \sin(x)^2}{2x^2} dx$$

Which simplifies to

$$u_1 = - \int 4 \sin(x^2) x \sin(x)^2 dx$$

Hence

$$\begin{aligned} u_1 = & \cos(x^2) - \frac{\cos(x^2 - 2x)}{2} \\ & + \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \right)}{2} - \frac{\cos(x^2 + 2x)}{2} \\ & - \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \right)}{2} \end{aligned}$$

And Eq. (3) becomes

$$u_2 = \int \frac{8 \cos(x^2) x^3 \sin(x)^2}{2x^2} dx$$

Which simplifies to

$$u_2 = \int 4 \cos(x^2) x \sin(x)^2 dx$$

Hence

$$\begin{aligned} u_2 = & \sin(x^2) - \frac{\sin(x^2 - 2x)}{2} \\ & - \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) + \sin(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \right)}{2} - \frac{\sin(x^2 + 2x)}{2} \\ & + \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) + \sin(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \right)}{2} \end{aligned}$$

Therefore the particular solution, from equation (1) is

$$\begin{aligned}
 y_p(x) = & \left(\cos(x^2) - \frac{\cos(x^2 - 2x)}{2} \right. \\
 & + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) - \sin(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \right)}{2} \\
 & \left. - \frac{\cos(x^2 + 2x)}{2} \right. \\
 & \left. - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) - \sin(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \right)}{2} \right) \cos(x^2) \\
 & + \sin(x^2) \left(\sin(x^2) - \frac{\sin(x^2 - 2x)}{2} \right. \\
 & \left. - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) + \sin(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \right)}{2} \right. \\
 & \left. - \frac{\sin(x^2 + 2x)}{2} \right. \\
 & \left. + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) + \sin(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \right)}{2} \right)
 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
 y_p(x) = & 1 + \frac{\operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \cos(x^2 + 1)}{2} \\
 & - \frac{\operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \cos(x^2 + 1)}{2} \\
 & - \frac{\operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \sin(x^2 + 1)}{2} \\
 & + \frac{\operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \sin(x^2 + 1)}{2} - \cos(2x)
 \end{aligned}$$

Therefore the general solution is

$$\begin{aligned}
 y = & y_h + y_p \\
 = & (c_1 \cos(x^2) + c_2 \sin(x^2)) + \left(1 + \frac{\operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \cos(x^2 + 1)}{2} \right. \\
 & - \frac{\operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \cos(x^2 + 1)}{2} - \frac{\operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \sin(x^2 + 1)}{2} \\
 & \left. + \frac{\operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2}\sqrt{\pi} \sin(x^2 + 1)}{2} - \cos(2x) \right)
 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = 1 + \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \cos(x^2 + 1)}{2} - \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \cos(x^2 + 1)}{2} - \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \sin(x^2 + 1)}{2} + \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \sin(x^2 + 1)}{2} - \cos(2x) + c_1 \cos(x^2) + c_2 \sin(x^2)$$

Solved as second order ode using change of variable on x method 1

Time used: 8.948 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x$, $B = -1$, $C = 4x^3$, $f(x) = 8x^3 \sin(x)^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$xy'' - y' + 4x^3y = 0$$

In normal form the ode

$$xy'' - y' + 4x^3y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{1}{x}$$

$$q(x) = 4x^2$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{2\sqrt{x^2}}{c} \tag{6}$$

$$\tau'' = \frac{2x}{c\sqrt{x^2}}$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{2x}{c\sqrt{x^2}} - \frac{1}{x} \frac{2\sqrt{x^2}}{c}}{\left(\frac{2\sqrt{x^2}}{c}\right)^2} \\ &= 0 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 2\sqrt{x^2} dx}{c} \\ &= \frac{x\sqrt{x^2}}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos(x\sqrt{x^2}) + c_2 \sin(x\sqrt{x^2})$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \cos(x\sqrt{x^2}) \\ y_2 &= \sin(x\sqrt{x^2}) \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x\sqrt{x^2}) & \sin(x\sqrt{x^2}) \\ \frac{d}{dx}(\cos(x\sqrt{x^2})) & \frac{d}{dx}(\sin(x\sqrt{x^2})) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x\sqrt{x^2}) & \sin(x\sqrt{x^2}) \\ -\left(\sqrt{x^2} + \frac{x^2}{\sqrt{x^2}}\right) \sin(x\sqrt{x^2}) & \left(\sqrt{x^2} + \frac{x^2}{\sqrt{x^2}}\right) \cos(x\sqrt{x^2}) \end{vmatrix}$$

Therefore

$$W = \left(\cos(x\sqrt{x^2})\right) \left(\left(\sqrt{x^2} + \frac{x^2}{\sqrt{x^2}}\right) \cos(x\sqrt{x^2})\right) - \left(\sin(x\sqrt{x^2})\right) \left(-\left(\sqrt{x^2} + \frac{x^2}{\sqrt{x^2}}\right) \sin(x\sqrt{x^2})\right)$$

Which simplifies to

$$W = \frac{2x^2 \left(\cos(x\sqrt{x^2})^2 + \sin(x\sqrt{x^2})^2\right)}{\sqrt{x^2}}$$

Which simplifies for $0 < x$ to

$$W = 2x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{8 \sin(x\sqrt{x^2}) x^3 \sin(x)^2}{2x^2} dx$$

Which simplifies for $0 < x$ to

$$u_1 = - \int 4 \sin(x^2) x \sin(x)^2 dx$$

Hence

$$u_1 = \cos(x^2) - \frac{\cos(x^2 - 2x)}{2} + \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \text{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) - \sin(1) \text{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right)\right)}{2} - \frac{\cos(x^2 + 2x)}{2} - \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \text{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) - \sin(1) \text{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right)\right)}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{8 \cos(x\sqrt{x^2}) x^3 \sin(x)^2}{2x^2} dx$$

Which simplifies for $0 < x$ to

$$u_2 = \int 4 \cos(x^2) x \sin(x)^2 dx$$

Hence

$$u_2 = \sin(x^2) - \frac{\sin(x^2 - 2x)}{2} - \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \text{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) + \sin(1) \text{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right)\right)}{2} - \frac{\sin(x^2 + 2x)}{2} + \frac{\sqrt{2} \sqrt{\pi} \left(\cos(1) \text{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) + \sin(1) \text{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right)\right)}{2}$$

Therefore the particular solution, from equation (1) is

$$\begin{aligned}
 y_p(x) = & \left(\cos(x^2) - \frac{\cos(x^2 - 2x)}{2} \right. \\
 & + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) - \sin(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \right)}{2} \\
 & - \frac{\cos(x^2 + 2x)}{2} \\
 & \left. - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) - \sin(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \right)}{2} \right) \cos(x\sqrt{x^2}) \\
 & + \sin(x\sqrt{x^2}) \left(\sin(x^2) - \frac{\sin(x^2 - 2x)}{2} \right. \\
 & - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) + \sin(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \right)}{2} \\
 & - \frac{\sin(x^2 + 2x)}{2} \\
 & \left. + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) + \sin(1) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \right)}{2} \right)
 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
 y_p(x) = & - \frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) (\cos(1) \sin(x^2) + \cos(x^2) \sin(1))}{2} \\
 & + \frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) (\cos(1) \sin(x^2) + \cos(x^2) \sin(1))}{2} \\
 & - \frac{\sqrt{2}\sqrt{\pi} (-\cos(x^2) \cos(1) + \sin(1) \sin(x^2)) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right)}{2} \\
 & + \frac{\sqrt{2}\sqrt{\pi} (-\cos(x^2) \cos(1) + \sin(1) \sin(x^2)) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right)}{2} \\
 & - (\cos(x^2)^2 + \sin(x^2)^2) (-1 + \cos(2x))
 \end{aligned}$$

Therefore the general solution is

$$\begin{aligned}
 y = & y_h + y_p \\
 = & \left(c_1 \cos(x\sqrt{x^2}) + c_2 \sin(x\sqrt{x^2}) \right) \\
 & + \left(- \frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) (\cos(1) \sin(x^2) + \cos(x^2) \sin(1))}{2} \right. \\
 & + \frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) (\cos(1) \sin(x^2) + \cos(x^2) \sin(1))}{2} \\
 & - \frac{\sqrt{2}\sqrt{\pi} (-\cos(x^2) \cos(1) + \sin(1) \sin(x^2)) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right)}{2} \\
 & + \frac{\sqrt{2}\sqrt{\pi} (-\cos(x^2) \cos(1) + \sin(1) \sin(x^2)) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right)}{2} \\
 & \left. - (\cos(x^2)^2 + \sin(x^2)^2) (-1 + \cos(2x)) \right)
 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned}
 y = & c_1 \cos(x\sqrt{x^2}) + c_2 \sin(x\sqrt{x^2}) \\
 & - \frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) (\cos(1)\sin(x^2) + \cos(x^2)\sin(1))}{2} \\
 & + \frac{\sqrt{2}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) (\cos(1)\sin(x^2) + \cos(x^2)\sin(1))}{2} \\
 & - \frac{\sqrt{2}\sqrt{\pi} (-\cos(x^2)\cos(1) + \sin(1)\sin(x^2)) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right)}{2} \\
 & + \frac{\sqrt{2}\sqrt{\pi} (-\cos(x^2)\cos(1) + \sin(1)\sin(x^2)) \operatorname{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right)}{2} \\
 & - (\cos(x^2)^2 + \sin(x^2)^2) (-1 + \cos(2x))
 \end{aligned}$$

Solved as second order Bessel ode

Time used: 0.258 (sec)

Writing the ode as

$$x^2 y'' - y'x + 4x^4 y = 8x^4 \sin(x)^2 \quad (1)$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2 y'' + y'x + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \operatorname{BesselJ}(n, \beta x^\gamma) + c_2 \operatorname{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\alpha = 1$$

$$\beta = 1$$

$$n = \frac{1}{2}$$

$$\gamma = 2$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{x\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

$$y_2 = -\frac{x\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{x\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} & -\frac{x\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \\ \frac{d}{dx} \left(\frac{x\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) & \frac{d}{dx} \left(-\frac{x\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{x\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} & -\frac{x\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \\ \frac{\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} & -\frac{\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \end{vmatrix}$$

Therefore

$$W = \left(\frac{x\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) \left(-\frac{\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) - \left(-\frac{x\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) \left(\frac{\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right)$$

Which simplifies to

$$W = \frac{4x \left(\cos(x^2)^2 + \sin(x^2)^2 \right)}{\pi}$$

Which simplifies to

$$W = \frac{4x}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = -\int \frac{\frac{8x^5 \sqrt{2} \cos(x^2) \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}}{\frac{4x^3}{\pi}} dx$$

Which simplifies for $0 < x$ to

$$u_1 = - \int -2x\sqrt{2}\sqrt{\pi} \cos(x^2) \sin(x)^2 dx$$

Hence

$$u_1 = 2\sqrt{2}\sqrt{\pi} \left(\frac{\sin(x^2)}{4} - \frac{\sin(x^2 - 2x)}{8} - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) + \sin(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \right)}{8} - \frac{\sin(x^2 + 2x)}{8} + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) + \sin(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \right)}{8} \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{8x^5\sqrt{2}\sin(x^2)\sin(x)^2}{\frac{\sqrt{\pi}\sqrt{x^2}}{\frac{4x^3}{\pi}}} dx$$

Which simplifies for $0 < x$ to

$$u_2 = \int 2x\sqrt{2}\sqrt{\pi} \sin(x^2) \sin(x)^2 dx$$

Hence

$$u_2 = 2\sqrt{2}\sqrt{\pi} \left(-\frac{\cos(x^2)}{4} + \frac{\cos(x^2 - 2x)}{8} - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \right)}{8} + \frac{\cos(x^2 + 2x)}{8} + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \right)}{8} \right)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{4 \left(\frac{\sin(x^2)}{4} - \frac{\sin(x^2 - 2x)}{8} - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) + \sin(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \right)}{8} - \frac{\sin(x^2 + 2x)}{8} + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) + \sin(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \right)}{8} \right) \sqrt{x^2}}{\sqrt{x^2}} + \frac{4x \cos(x^2) \left(-\frac{\cos(x^2)}{4} + \frac{\cos(x^2 - 2x)}{8} - \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \right)}{8} + \frac{\cos(x^2 + 2x)}{8} + \frac{\sqrt{2}\sqrt{\pi} \left(\cos(1) \operatorname{FresnelS} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) - \sin(1) \operatorname{FresnelC} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \right)}{8} \right) \sqrt{x^2}}{\sqrt{x^2}} - \cos(2x)$$

Which simplifies to

$$y_p(x) = 1 + \frac{\operatorname{FresnelS} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \sqrt{2}\sqrt{\pi} \cos(x^2 + 1)}{2} - \frac{\operatorname{FresnelS} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \sqrt{2}\sqrt{\pi} \cos(x^2 + 1)}{2} - \frac{\operatorname{FresnelC} \left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}} \right) \sqrt{2}\sqrt{\pi} \sin(x^2 + 1)}{2} + \frac{\operatorname{FresnelC} \left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}} \right) \sqrt{2}\sqrt{\pi} \sin(x^2 + 1)}{2} - \cos(2x)$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(\frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) + \left(1 + \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \cos(x^2 + 1)}{2} \right. \\
 &\quad - \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \cos(x^2 + 1)}{2} - \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \sin(x^2 + 1)}{2} \\
 &\quad \left. + \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \sin(x^2 + 1)}{2} - \cos(2x) \right)
 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned}
 y &= \frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + 1 + \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \cos(x^2 + 1)}{2} \\
 &\quad - \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \cos(x^2 + 1)}{2} - \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \sin(x^2 + 1)}{2} \\
 &\quad + \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{2} \sqrt{\pi} \sin(x^2 + 1)}{2} - \cos(2x)
 \end{aligned}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.767 (sec)

Writing the ode as

$$xy'' - y' + 4x^3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}
 A &= x \\
 B &= -1 \\
 C &= 4x^3
 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}
 r &= \frac{s}{t} \\
 &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}
 \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16x^4 + 3}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -16x^4 + 3 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{-16x^4 + 3}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.79: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 4 \\ &= -2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Therefore

$$L = [1, 2]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -4x^2 + \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \quad (8)$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx 2ix - \frac{3i}{16x^3} - \frac{9i}{1024x^7} - \frac{27i}{32768x^{11}} - \frac{405i}{4194304x^{15}} - \frac{1701i}{134217728x^{19}} - \frac{15309i}{8589934592x^{23}} - \frac{72171i}{274877906944x^{27}} + \dots \quad (9)$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = 2i$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= 2ix \end{aligned} \quad (10)$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = -4x^2$$

This shows that the coefficient of 1 in the above is 0. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{-16x^4 + 3}{4x^2} \\ &= Q + \frac{R}{4x^2} \\ &= (-4x^2) + \left(\frac{3}{4x^2}\right) \\ &= -4x^2 + \frac{3}{4x^2} \end{aligned}$$

We see that the coefficient of the term x in the quotient is 0. Now b can be found.

$$\begin{aligned} b &= (0) - (0) \\ &= 0 \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 2ix \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{0}{2i} - 1 \right) = -\frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{0}{2i} - 1 \right) = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{-16x^4 + 3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$2ix$	$-\frac{1}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-) (2ix) \\ &= -\frac{1}{2x} - 2ix \\ &= -\frac{1}{2x} - 2ix \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2 \left(-\frac{1}{2x} - 2ix \right) (0) + \left(\left(\frac{1}{2x^2} - 2i \right) + \left(-\frac{1}{2x} - 2ix \right)^2 - \left(\frac{-16x^4 + 3}{4x^2} \right) \right) = 0$$

$0 = 0$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int (-\frac{1}{2x} - 2ix) dx} \\ &= \frac{e^{-ix^2}}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{x} dx} \\ &= z_1 e^{\frac{\ln(x)}{2}} \\ &= z_1 (\sqrt{x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-ix^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{x} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{ie^{2ix^2}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-ix^2}) + c_2 \left(e^{-ix^2} \left(-\frac{ie^{2ix^2}}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$xy'' - y' + 4x^3y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-ix^2} - \frac{ic_2 e^{ix^2}}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-ix^2}$$

$$y_2 = -\frac{ie^{ix^2}}{4}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-ix^2} & -\frac{ie^{ix^2}}{4} \\ \frac{d}{dx}(e^{-ix^2}) & \frac{d}{dx}\left(-\frac{ie^{ix^2}}{4}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-ix^2} & -\frac{ie^{ix^2}}{4} \\ -2ix e^{-ix^2} & \frac{x e^{ix^2}}{2} \end{vmatrix}$$

Therefore

$$W = (e^{-ix^2}) \left(\frac{x e^{ix^2}}{2} \right) - \left(-\frac{ie^{ix^2}}{4} \right) (-2ix e^{-ix^2})$$

Which simplifies to

$$W = e^{-ix^2} x e^{ix^2}$$

Which simplifies to

$$W = x$$

Therefore Eq. (2) becomes

$$u_1 = -\int \frac{-2ie^{ix^2} x^3 \sin(x)^2}{x^2} dx$$

Which simplifies to

$$u_1 = -\int -2ie^{ix^2} x \sin(x)^2 dx$$

Hence

$$u_1 = -\frac{e^{ix(2+x)}}{4} + \frac{i\sqrt{\pi} e^{-i} \operatorname{erf}\left(\sqrt{-i}x - \frac{i}{\sqrt{-i}}\right)}{4\sqrt{-i}} - \frac{e^{ix(-2+x)}}{4} - \frac{i\sqrt{\pi} e^{-i} \operatorname{erf}\left(\sqrt{-i}x + \frac{i}{\sqrt{-i}}\right)}{4\sqrt{-i}} + \frac{e^{ix^2}}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{8e^{-ix^2} x^3 \sin(x)^2}{x^2} dx$$

Which simplifies to

$$u_2 = \int 8e^{-ix^2} x \sin(x)^2 dx$$

Hence

$$u_2 = -ie^{-ix(-2+x)} + \sqrt{\pi} e^i (-1)^{3/4} \operatorname{erf}\left((-1)^{1/4} x - (-1)^{1/4}\right) - ie^{-ix(2+x)} - \sqrt{\pi} e^i (-1)^{3/4} \operatorname{erf}\left((-1)^{1/4} x + (-1)^{1/4}\right) + 2ie^{-ix^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{e^{ix(2+x)}}{4} + \frac{i\sqrt{\pi} e^{-i} \operatorname{erf}\left(\sqrt{-i}x - \frac{i}{\sqrt{-i}}\right)}{4\sqrt{-i}} - \frac{e^{ix(-2+x)}}{4} - \frac{i\sqrt{\pi} e^{-i} \operatorname{erf}\left(\sqrt{-i}x + \frac{i}{\sqrt{-i}}\right)}{4\sqrt{-i}} + \frac{e^{ix^2}}{2} \right) e^{-ix^2} - \frac{ie^{ix^2} \left(-ie^{-ix(-2+x)} + \sqrt{\pi} e^i (-1)^{3/4} \operatorname{erf}\left((-1)^{1/4} x - (-1)^{1/4}\right) - ie^{-ix(2+x)} - \sqrt{\pi} e^i (-1)^{3/4} \operatorname{erf}\left((-1)^{1/4} x + (-1)^{1/4}\right) + 2ie^{-ix^2} \right)}{4}$$

Which simplifies to

$$y_p(x) = \left(\frac{1}{8} + \frac{i}{8} \right) \left(i\sqrt{\pi} \left(\operatorname{erf}\left(\left(\frac{1}{2} - \frac{i}{2}\right)(x+1)\sqrt{2}\right) - \operatorname{erf}\left(\left(\frac{1}{2} - \frac{i}{2}\right)(x-1)\sqrt{2}\right) \right) e^{-ix^2-i} - \sqrt{\pi} \left(\operatorname{erf}\left(\left(\frac{1}{2} + \frac{i}{2}\right)(x+1)\sqrt{2}\right) - \operatorname{erf}\left(\left(\frac{1}{2} + \frac{i}{2}\right)(x-1)\sqrt{2}\right) \right) e^{ix^2+i} + (-2+2i)(-1+\cos(2x))\sqrt{2} \right) \sqrt{2}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 e^{-ix^2} - \frac{ic_2 e^{ix^2}}{4} \right) + \left(\left(\frac{1}{8} + \frac{i}{8} \right) \left(i\sqrt{\pi} \left(\operatorname{erf}\left(\left(\frac{1}{2} - \frac{i}{2}\right)(x+1)\sqrt{2}\right) - \operatorname{erf}\left(\left(\frac{1}{2} - \frac{i}{2}\right)(x-1)\sqrt{2}\right) \right) e^{-ix^2-i} - \sqrt{\pi} \left(\operatorname{erf}\left(\left(\frac{1}{2} + \frac{i}{2}\right)(x+1)\sqrt{2}\right) - \operatorname{erf}\left(\left(\frac{1}{2} + \frac{i}{2}\right)(x-1)\sqrt{2}\right) \right) e^{ix^2+i} + (-2+2i)(-1+\cos(2x))\sqrt{2} \right) \sqrt{2} \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned}
y = c_1 e^{-ix^2} - \frac{ic_2 e^{ix^2}}{4} \\
+ \left(\frac{1}{8} + \frac{i}{8}\right) \left(i\sqrt{\pi} \left(\operatorname{erf} \left(\left(\frac{1}{2} - \frac{i}{2}\right) (x+1)\sqrt{2} \right) - \operatorname{erf} \left(\left(\frac{1}{2} - \frac{i}{2}\right) (x-1)\sqrt{2} \right) \right) e^{-ix^2-i} \right. \\
\left. - \sqrt{\pi} \left(\operatorname{erf} \left(\left(\frac{1}{2} + \frac{i}{2}\right) (x+1)\sqrt{2} \right) - \operatorname{erf} \left(\left(\frac{1}{2} + \frac{i}{2}\right) (x-1)\sqrt{2} \right) \right) e^{ix^2+i} \right. \\
\left. + (-2 + 2i) (-1 + \cos(2x)) \sqrt{2} \right) \sqrt{2}
\end{aligned}$$

Solved as second order ode adjoint method

Time used: 7.800 (sec)

In normal form the ode

$$xy'' - y' + 4x^3y = 8x^3 \sin(x)^2 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned}
p(x) &= -\frac{1}{x} \\
q(x) &= 4x^2 \\
r(x) &= 8x^2 \sin(x)^2
\end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned}
\xi'' - (\xi p)' + \xi q &= 0 \\
\xi'' - \left(-\frac{\xi(x)}{x}\right)' + (4x^2\xi(x)) &= 0 \\
\xi''(x) + \frac{\xi'(x)}{x} + \frac{(4x^4 - 1)\xi(x)}{x^2} &= 0
\end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi''x^2 + \xi'x + (4x^4 - 1)\xi = 0 \quad (1)$$

Bessel ode has the form

$$\xi''x^2 + \xi'x + (-n^2 + x^2)\xi = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$\xi''x^2 + (1 - 2\alpha)x\xi' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)\xi = 0 \quad (3)$$

With the standard solution

$$\xi = x^\alpha (c_1 \operatorname{BesselJ}(n, \beta x^\gamma) + c_2 \operatorname{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}
\alpha &= 0 \\
\beta &= 1 \\
n &= -\frac{1}{2} \\
\gamma &= 2
\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$\xi = \frac{c_1 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{c_2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(-\frac{1}{x} - \frac{-\frac{c_1 \sqrt{2} \cos(x^2) x}{\sqrt{\pi} (x^2)^{3/2}} - \frac{2c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 \sqrt{2} \sin(x^2) x}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}}{\frac{c_1 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{c_2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}} \right) = \frac{-\frac{\sqrt{2} \sqrt{x^2} c_1 e^{-i} \operatorname{erf}(\sqrt{-i} x + \frac{i}{\sqrt{-i}})}{2x \sqrt{-i}} + \dots}{\dots}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{2(c_2 \cos(x^2) - c_1 \sin(x^2)) x}{c_1 \cos(x^2) + c_2 \sin(x^2)} \\ p(x) &= \frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))(ic_2+c_1) + \dots}{\dots} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{2(c_2 \cos(x^2) - c_1 \sin(x^2)) x}{c_1 \cos(x^2) + c_2 \sin(x^2)} dx} \\ &= \frac{1}{c_1 \cos(x^2) + c_2 \sin(x^2)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))(ic_2+c_1) + \dots}{\dots} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right) &= \left(\frac{1}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right) \left(\frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))(ic_2+c_1) + \dots}{\dots} \right) \\ d \left(\frac{y}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right) &= \left(\frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))(ic_2+c_1) + e^i \dots}{\dots} \right) \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_1 \cos(x^2) + c_2 \sin(x^2)} &= \int \frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))}{c_1 \cos(x^2) + c_2 \sin(x^2)} \\ &= \int \frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))}{c_1 \cos(x^2) + c_2 \sin(x^2)} \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x^2) + c_2 \sin(x^2)}$ gives the final solution

$$y = (c_1 \cos(x^2) + c_2 \sin(x^2)) \left(\int \frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right)$$

Hence, the solution found using Lagrange adjoint equation method is

$$\begin{aligned} y &= (c_1 \cos(x^2) \\ &+ c_2 \sin(x^2)) \left(\int \frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right. \\ &\qquad \qquad \qquad \left. + c_3 \right) \end{aligned}$$

The constants can be merged to give

$$\begin{aligned} y &= (c_1 \cos(x^2) \\ &+ c_2 \sin(x^2)) \left(\int \frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right. \\ &\qquad \qquad \qquad \left. + 1 \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned} y &= (c_1 \cos(x^2) \\ &+ c_2 \sin(x^2)) \left(\int \frac{(1-i)\sqrt{2}(-e^{-i}\sqrt{\pi} \operatorname{erf}((\frac{1}{2}-\frac{i}{2})(x-1)\sqrt{2}))(ic_1+c_2) - e^i\sqrt{\pi} \operatorname{erf}((\frac{1}{2}+\frac{i}{2})(x-1)\sqrt{2}))}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right. \\ &\qquad \qquad \qquad \left. + 1 \right) \end{aligned}$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    <- linear_1 successful
<- solving first the homogeneous part of the ODE successful`
```

Maple dsolve solution

Solving time : 0.009 (sec)

Leaf size : 124

```
dsolve(x*diff(diff(y(x),x),x)-diff(y(x),x)+4*y(x)*x^3 = 8*sin(x)^2*x^3,
y(x),singsol=all)
```

$$y = \sin(x^2) c_2 + \cos(x^2) c_1 + 1 - \cos(2x) - \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{\pi} \sqrt{2} \sin(x^2 + 1)}{2} \\ + \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x-1)}{\sqrt{\pi}}\right) \sqrt{\pi} \sqrt{2} \cos(x^2 + 1)}{2} + \frac{\text{FresnelC}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{\pi} \sqrt{2} \sin(x^2 + 1)}{2} \\ - \frac{\text{FresnelS}\left(\frac{\sqrt{2}(x+1)}{\sqrt{\pi}}\right) \sqrt{\pi} \sqrt{2} \cos(x^2 + 1)}{2}$$

Mathematica DSolve solution

Solving time : 1.292 (sec)

Leaf size : 147

```
DSolve[{x*D[y[x]},{x,2]}-D[y[x],x]+4*x^3*y[x]==8*x^3*Sin[x]^2,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{2} \left(-\sqrt{2\pi} \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(x-1)\right) \sin(x^2 + 1) \right. \\ \left. + \sqrt{2\pi} \text{FresnelC}\left(\sqrt{\frac{2}{\pi}}(x+1)\right) \sin(x^2 + 1) \right. \\ \left. + \sqrt{2\pi} \text{FresnelS}\left(\sqrt{\frac{2}{\pi}}(x-1)\right) \cos(x^2 + 1) \right. \\ \left. - \sqrt{2\pi} \text{FresnelS}\left(\sqrt{\frac{2}{\pi}}(x+1)\right) \cos(x^2 + 1) + 2c_1 \cos(x^2) + 2c_2 \sin(x^2) \right. \\ \left. - 2 \cos(2x) + 2 \right)$$

2.2.24 problem 25

Solved as second order ode using change of variable on x
 method 2 679
 Solved as second order ode using change of variable on x
 method 1 682
 Solved as second order Bessel ode 684
 Solved as second order ode using Kovacic algorithm 687
 Solved as second order ode adjoint method 693
 Maple step by step solution 695
 Maple trace 695
 Maple dsolve solution 695
 Mathematica DSolve solution 695

Internal problem ID [8823]

Book : Second order enumerated odes

Section : section 2

Problem number : 25

Date solved : Thursday, December 12, 2024 at 09:52:46 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$xy'' - y' + 4x^3y = x^5$$

Solved as second order ode using change of variable on x method 2

Time used: 0.374 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$xy'' - y' + 4x^3y = 0$$

In normal form the ode

$$xy'' - y' + 4x^3y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{1}{x}$$

$$q(x) = 4x^2$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned}\tau &= \int e^{-\int p(x)dx} dx \\ &= \int e^{-\int -\frac{1}{x}dx} dx \\ &= \int e^{\ln(x)} dx \\ &= \int x dx \\ &= \frac{x^2}{2}\end{aligned}\tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{4x^2}{x^2} \\ &= 4\end{aligned}\tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + 4y(\tau) &= 0\end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\tau\lambda} + 4e^{\tau\lambda} = 0\tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 4 = 0\tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +2i \\ \lambda_2 &= -2i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 2i \\ \lambda_2 &= -2i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau}(c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0(c_1 \cos(2\tau) + c_2 \sin(2\tau))$$

Or

$$y(\tau) = c_1 \cos(2\tau) + c_2 \sin(2\tau)$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(x^2) + c_2 \sin(x^2)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x^2) + c_2 \sin(x^2)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^5$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3, x^4, x^5\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x^2), \sin(x^2)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_6 x^5 + A_5 x^4 + A_4 x^3 + A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5, A_6\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned}x(20x^3 A_6 + 12x^2 A_5 + 6x A_4 + 2A_3) - 5x^4 A_6 - 4x^3 A_5 - 3x^2 A_4 \\ - 2x A_3 - A_2 + 4x^3(A_6 x^5 + A_5 x^4 + A_4 x^3 + A_3 x^2 + A_2 x + A_1) = x^5\end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = 0, A_3 = \frac{1}{4}, A_4 = 0, A_5 = 0, A_6 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x^2) + c_2 \sin(x^2)) + \left(\frac{x^2}{4}\right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{4} + c_1 \cos(x^2) + c_2 \sin(x^2)$$

Solved as second order ode using change of variable on x method 1

Time used: 0.227 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x, B = -1, C = 4x^3, f(x) = x^5$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$xy'' - y' + 4x^3y = 0$$

In normal form the ode

$$xy'' - y' + 4x^3y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$\begin{aligned} p(x) &= -\frac{1}{x} \\ q(x) &= 4x^2 \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{2\sqrt{x^2}}{c} \\ \tau'' &= \frac{2x}{c\sqrt{x^2}}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{2x}{c\sqrt{x^2}} - \frac{1}{x}\frac{2\sqrt{x^2}}{c}}{\left(\frac{2\sqrt{x^2}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int 2\sqrt{x^2} dx}{c} \\ &= \frac{x\sqrt{x^2}}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos(x\sqrt{x^2}) + c_2 \sin(x\sqrt{x^2})$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^5$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3, x^4, x^5\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \cos(x\sqrt{x^2}), \sin(x\sqrt{x^2}) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_6x^5 + A_5x^4 + A_4x^3 + A_3x^2 + A_2x + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5, A_6\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$x(20x^3A_6 + 12x^2A_5 + 6xA_4 + 2A_3) - 5x^4A_6 - 4x^3A_5 - 3x^2A_4 - 2xA_3 - A_2 + 4x^3(A_6x^5 + A_5x^4 + A_4x^3 + A_3x^2 + A_2x + A_1) = x^5$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = 0, A_3 = \frac{1}{4}, A_4 = 0, A_5 = 0, A_6 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2}{4}$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1 \cos(x\sqrt{x^2}) + c_2 \sin(x\sqrt{x^2}) \right) + \left(\frac{x^2}{4} \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos(x\sqrt{x^2}) + c_2 \sin(x\sqrt{x^2}) + \frac{x^2}{4}$$

Solved as second order Bessel ode

Time used: 0.371 (sec)

Writing the ode as

$$x^2y'' - y'x + 4x^4y = x^6 \tag{1}$$

Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE and y_p is a particular solution to the non-homogeneous ODE. Bessel ode has the form

$$x^2y'' + y'x + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\alpha = 1 \\ \beta = 1 \\ n = \frac{1}{2} \\ \gamma = 2$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

$$y_2 = -\frac{x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} & -\frac{x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \\ \frac{d}{dx} \left(\frac{x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) & \frac{d}{dx} \left(-\frac{x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} & -\frac{x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \\ \frac{\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} & -\frac{\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \end{vmatrix}$$

Therefore

$$W = \left(\frac{x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) \left(-\frac{\sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) - \left(-\frac{x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) \left(\frac{\sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{x^2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2x^2 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right)$$

Which simplifies to

$$W = \frac{4x \left(\sin(x^2)^2 + \cos(x^2)^2 \right)}{\pi}$$

Which simplifies to

$$W = \frac{4x}{\pi}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-\frac{x^7 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}}{\frac{4x^3}{\pi}} dx$$

Which simplifies for $0 < x$ to

$$u_1 = - \int -\frac{x^3 \sqrt{2} \sqrt{\pi} \cos(x^2)}{4} dx$$

Hence

$$u_1 = \frac{\sqrt{2} \sqrt{\pi} \left(\frac{\cos(x^2)}{2} + \frac{x^2 \sin(x^2)}{2} \right)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{x^7 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}}{\frac{4x^3}{\pi}} dx$$

Which simplifies for $0 < x$ to

$$u_2 = \int \frac{x^3 \sqrt{2} \sqrt{\pi} \sin(x^2)}{4} dx$$

Hence

$$u_2 = \frac{\sqrt{2} \sqrt{\pi} \left(\frac{\sin(x^2)}{2} - \frac{x^2 \cos(x^2)}{2} \right)}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(\frac{\cos(x^2)}{2} + \frac{x^2 \sin(x^2)}{2} \right) x \sin(x^2)}{2\sqrt{x^2}} - \frac{x \cos(x^2) \left(\frac{\sin(x^2)}{2} - \frac{x^2 \cos(x^2)}{2} \right)}{2\sqrt{x^2}}$$

Which simplifies to

$$y_p(x) = \frac{x\sqrt{x^2}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right) + \left(\frac{x\sqrt{x^2}}{4} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{x\sqrt{x^2}}{4}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.347 (sec)

Writing the ode as

$$xy'' - y' + 4x^3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= -1 \\ C &= 4x^3 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16x^4 + 3}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -16x^4 + 3 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{-16x^4 + 3}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.80: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 4 \\ &= -2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is -2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Therefore

$$L = [1, 2]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -4x^2 + \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $O_r(\infty) = -2$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{2}{2} = 1$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^1 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^1$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx 2ix - \frac{3i}{16x^3} - \frac{9i}{1024x^7} - \frac{27i}{32768x^{11}} - \frac{405i}{4194304x^{15}} - \frac{1701i}{134217728x^{19}} - \frac{15309i}{8589934592x^{23}} - \frac{72171i}{274877906944x^{27}} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = 2i$$

From Eq. (9) the sum up to $v = 1$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^1 a_i x^i \\ &= 2ix \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^0 = 1$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = -4x^2$$

This shows that the coefficient of 1 in the above is 0. Now we need to find the coefficient of 1 in r . How this is done depends on if $v = 0$ or not. Since $v = 1$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of 1 in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{-16x^4 + 3}{4x^2} \\ &= Q + \frac{R}{4x^2} \\ &= (-4x^2) + \left(\frac{3}{4x^2}\right) \\ &= -4x^2 + \frac{3}{4x^2} \end{aligned}$$

We see that the coefficient of the term x in the quotient is 0. Now b can be found.

$$\begin{aligned} b &= (0) - (0) \\ &= 0 \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 2ix \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{0}{2i} - 1 \right) = -\frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{0}{2i} - 1 \right) = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{-16x^4 + 3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-2	$2ix$	$-\frac{1}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-)(2ix) \\ &= -\frac{1}{2x} - 2ix \\ &= -\frac{1}{2x} - 2ix \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-\frac{1}{2x} - 2ix \right) (0) + \left(\left(\frac{1}{2x^2} - 2i \right) + \left(-\frac{1}{2x} - 2ix \right)^2 - \left(\frac{-16x^4 + 3}{4x^2} \right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \left(-\frac{1}{2x} - 2ix \right) dx} \\ &= \frac{e^{-ix^2}}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{x} dx} \\ &= z_1 e^{\frac{\ln(x)}{2}} \\ &= z_1 (\sqrt{x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-ix^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{x} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{ie^{2ix^2}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-ix^2} \right) + c_2 \left(e^{-ix^2} \left(-\frac{ie^{2ix^2}}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$xy'' - y' + 4x^3y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-ix^2} - \frac{ic_2 e^{ix^2}}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{-ix^2} \\ y_2 &= -\frac{ie^{ix^2}}{4} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-ix^2} & -\frac{ie^{ix^2}}{4} \\ \frac{d}{dx}(e^{-ix^2}) & \frac{d}{dx}\left(-\frac{ie^{ix^2}}{4}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-ix^2} & -\frac{ie^{ix^2}}{4} \\ -2ix e^{-ix^2} & \frac{x e^{ix^2}}{2} \end{vmatrix}$$

Therefore

$$W = (e^{-ix^2}) \left(\frac{x e^{ix^2}}{2} \right) - \left(-\frac{ie^{ix^2}}{4} \right) (-2ix e^{-ix^2})$$

Which simplifies to

$$W = e^{-ix^2} x e^{ix^2}$$

Which simplifies to

$$W = x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-\frac{ie^{ix^2} x^5}{4}}{x^2} dx$$

Which simplifies to

$$u_1 = - \int -\frac{ie^{ix^2} x^3}{4} dx$$

Hence

$$u_1 = -\frac{i(ix^2 - 1) e^{ix^2}}{8}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-ix^2} x^5}{x^2} dx$$

Which simplifies to

$$u_2 = \int e^{-ix^2} x^3 dx$$

Hence

$$u_2 = \frac{(ix^2 + 1) e^{-ix^2}}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{i(ix^2 - 1) e^{ix^2} e^{-ix^2}}{8} - \frac{ie^{ix^2} (ix^2 + 1) e^{-ix^2}}{8}$$

Which simplifies to

$$y_p(x) = \frac{x^2}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-ix^2} - \frac{ic_2 e^{ix^2}}{4} \right) + \left(\frac{x^2}{4} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-ix^2} - \frac{ic_2 e^{ix^2}}{4} + \frac{x^2}{4}$$

Solved as second order ode adjoint method

Time used: 2.179 (sec)

In normal form the ode

$$xy'' - y' + 4x^3y = x^5 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{1}{x} \\ q(x) &= 4x^2 \\ r(x) &= x^4 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(-\frac{\xi(x)}{x} \right)' + (4x^2\xi(x)) &= 0 \\ \xi''(x) + \frac{\xi'(x)}{x} + \frac{(4x^4 - 1)\xi(x)}{x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi''x^2 + \xi'x + (4x^4 - 1)\xi = 0 \quad (1)$$

Bessel ode has the form

$$\xi''x^2 + \xi'x + (-n^2 + x^2)\xi = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$\xi''x^2 + (1 - 2\alpha)x\xi' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)\xi = 0 \quad (3)$$

With the standard solution

$$\xi = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= 0 \\ \beta &= 1 \\ n &= -\frac{1}{2} \\ \gamma &= 2 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$\xi = \frac{c_1 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{c_2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(-\frac{1}{x} - \frac{-\frac{c_1 \sqrt{2} \cos(x^2)x}{\sqrt{\pi} (x^2)^{3/2}} - \frac{2c_1 x \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} - \frac{c_2 \sqrt{2} \sin(x^2)x}{\sqrt{\pi} (x^2)^{3/2}} + \frac{2c_2 x \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}}}{\frac{c_1 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{c_2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}}} \right) = \frac{\sqrt{2} \operatorname{csgn}(x) \left(\frac{c_1 (\cos(x^2) + x^2 \sin(x^2))}{2} \right)}{\left(\frac{c_1 \sqrt{2} \cos(x^2)}{\sqrt{\pi} \sqrt{x^2}} + \frac{c_2 \sqrt{2} \sin(x^2)}{\sqrt{\pi} \sqrt{x^2}} \right)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{2x(-c_1 \sin(x^2) + c_2 \cos(x^2))}{c_1 \cos(x^2) + c_2 \sin(x^2)} \\ p(x) &= \frac{((-c_2 x^2 + c_1) \cos(x^2) + \sin(x^2) (c_1 x^2 + c_2)) x}{2c_1 \cos(x^2) + 2c_2 \sin(x^2)} \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{2x(-c_1 \sin(x^2) + c_2 \cos(x^2))}{c_1 \cos(x^2) + c_2 \sin(x^2)} dx} \\ &= \frac{1}{c_1 \cos(x^2) + c_2 \sin(x^2)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{((-c_2 x^2 + c_1) \cos(x^2) + \sin(x^2) (c_1 x^2 + c_2)) x}{2c_1 \cos(x^2) + 2c_2 \sin(x^2)} \right) \\ \frac{d}{dx} \left(\frac{y}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right) &= \left(\frac{1}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right) \left(\frac{((-c_2 x^2 + c_1) \cos(x^2) + \sin(x^2) (c_1 x^2 + c_2)) x}{2c_1 \cos(x^2) + 2c_2 \sin(x^2)} \right) \\ d \left(\frac{y}{c_1 \cos(x^2) + c_2 \sin(x^2)} \right) &= \left(\frac{((-c_2 x^2 + c_1) \cos(x^2) + \sin(x^2) (c_1 x^2 + c_2)) x}{(2c_1 \cos(x^2) + 2c_2 \sin(x^2)) (c_1 \cos(x^2) + c_2 \sin(x^2))} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_1 \cos(x^2) + c_2 \sin(x^2)} &= \int \frac{((-c_2 x^2 + c_1) \cos(x^2) + \sin(x^2) (c_1 x^2 + c_2)) x}{(2c_1 \cos(x^2) + 2c_2 \sin(x^2)) (c_1 \cos(x^2) + c_2 \sin(x^2))} dx \\ &= \frac{-\frac{x^2}{4} - \frac{x^2 \tan(\frac{x^2}{2})^2}{2} - \frac{x^2 \tan(\frac{x^2}{2})^4}{4}}{\left(1 + \tan\left(\frac{x^2}{2}\right)^2\right) \left(\tan\left(\frac{x^2}{2}\right)^2 c_1 - 2 \tan\left(\frac{x^2}{2}\right) c_2 - c_1\right)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_1 \cos(x^2) + c_2 \sin(x^2)}$ gives the final solution

$$y = c_1 c_3 \cos(x^2) + c_2 c_3 \sin(x^2) + \frac{x^2}{4}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_1 c_3 \cos(x^2) + c_2 c_3 \sin(x^2) + \frac{x^2}{4}$$

The constants can be merged to give

$$y = \frac{x^2}{4} + c_1 \cos(x^2) + c_2 \sin(x^2)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{x^2}{4} + c_1 \cos(x^2) + c_2 \sin(x^2)$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    <- linear_1 successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.007 (sec)

Leaf size : 22

```

dsolve(x*diff(diff(y(x),x),x)-diff(y(x),x)+4*y(x)*x^3 = x^5,
        y(x),singsol=all)

```

$$y = \sin(x^2) c_2 + \cos(x^2) c_1 + \frac{x^2}{4}$$

Mathematica DSolve solution

Solving time : 0.064 (sec)

Leaf size : 27

```

DSolve[{x*D[y[x],{x,2}]-D[y[x],x]+4*x^3*y[x]==x^5,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{x^2}{4} + c_1 \cos(x^2) + c_2 \sin(x^2)$$

2.2.25 problem 25

Solved as second order ode using change of variable on x
 method 2 696
 Solved as second order ode using change of variable on x
 method 1 700
 Maple step by step solution 704
 Maple trace 704
 Maple dsolve solution 704
 Mathematica DSolve solution 704

Internal problem ID [8824]

Book : Second order enumerated odes

Section : section 2

Problem number : 25

Date solved : Thursday, December 12, 2024 at 09:52:50 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$\cos(x)y'' + \sin(x)y' - 2y\cos(x)^3 = 2\cos(x)^5$$

Solved as second order ode using change of variable on x method 2

Time used: 0.868 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$\cos(x)y'' + \sin(x)y' - 2y\cos(x)^3 = 0$$

In normal form the ode

$$\cos(x)y'' + \sin(x)y' - 2y\cos(x)^3 = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \tan(x)$$

$$q(x) = -2\cos(x)^2$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\int p(x) dx} dx \\ &= \int e^{-\int \tan(x) dx} dx \\ &= \int e^{\ln(\cos(x))} dx \\ &= \int \cos(x) dx \\ &= \sin(x) \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-2 \cos(x)^2}{\cos(x)^2} \\ &= -2 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2} y(\tau) + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) - 2y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -2$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 2e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-2)} \\ &= \pm \sqrt{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +\sqrt{2} \\ \lambda_2 &= -\sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \sqrt{2} \\ \lambda_2 &= -\sqrt{2}\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y(\tau) &= c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau} \\ y(\tau) &= c_1 e^{(\sqrt{2})\tau} + c_2 e^{(-\sqrt{2})\tau}\end{aligned}$$

Or

$$y(\tau) = c_1 e^{\tau\sqrt{2}} + c_2 e^{-\tau\sqrt{2}}$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 e^{\sin(x)\sqrt{2}} + c_2 e^{-\sin(x)\sqrt{2}}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{\sin(x)\sqrt{2}} + c_2 e^{-\sin(x)\sqrt{2}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= e^{-\sin(x)\sqrt{2}} \\ y_2 &= e^{\sin(x)\sqrt{2}}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-\sin(x)\sqrt{2}} & e^{\sin(x)\sqrt{2}} \\ \frac{d}{dx} \left(e^{-\sin(x)\sqrt{2}} \right) & \frac{d}{dx} \left(e^{\sin(x)\sqrt{2}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-\sin(x)\sqrt{2}} & e^{\sin(x)\sqrt{2}} \\ -e^{-\sin(x)\sqrt{2}} \cos(x) \sqrt{2} & \cos(x) \sqrt{2} e^{\sin(x)\sqrt{2}} \end{vmatrix}$$

Therefore

$$W = \left(e^{-\sin(x)\sqrt{2}} \right) \left(\cos(x) \sqrt{2} e^{\sin(x)\sqrt{2}} \right) - \left(e^{\sin(x)\sqrt{2}} \right) \left(-e^{-\sin(x)\sqrt{2}} \cos(x) \sqrt{2} \right)$$

Which simplifies to

$$W = 2 \cos(x) \sqrt{2}$$

Which simplifies to

$$W = 2 \cos(x) \sqrt{2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2 e^{\sin(x)\sqrt{2}} \cos(x)^5}{2 \cos(x)^2 \sqrt{2}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{\sin(x)\sqrt{2}} \cos(x)^3 \sqrt{2}}{2} dx$$

Hence

$$u_1 = \frac{e^{\sin(x)\sqrt{2}} \sin(x)^2}{2} - \frac{\sin(x) \sqrt{2} e^{\sin(x)\sqrt{2}}}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2 e^{-\sin(x)\sqrt{2}} \cos(x)^5}{2 \cos(x)^2 \sqrt{2}} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-\sin(x)\sqrt{2}} \cos(x)^3 \sqrt{2}}{2} dx$$

Hence

$$u_2 = \frac{e^{-\sin(x)\sqrt{2}} \sin(x)^2}{2} + \frac{\sin(x) \sqrt{2} e^{-\sin(x)\sqrt{2}}}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{e^{\sin(x)\sqrt{2}} \sin(x)^2}{2} - \frac{\sin(x) \sqrt{2} e^{\sin(x)\sqrt{2}}}{2} \right) e^{-\sin(x)\sqrt{2}} \\ + e^{\sin(x)\sqrt{2}} \left(\frac{e^{-\sin(x)\sqrt{2}} \sin(x)^2}{2} + \frac{\sin(x) \sqrt{2} e^{-\sin(x)\sqrt{2}}}{2} \right)$$

Which simplifies to

$$y_p(x) = \sin(x)^2$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1 e^{\sin(x)\sqrt{2}} + c_2 e^{-\sin(x)\sqrt{2}} \right) + (\sin(x)^2)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \sin(x)^2 + c_1 e^{\sin(x)\sqrt{2}} + c_2 e^{-\sin(x)\sqrt{2}}$$

Solved as second order ode using change of variable on x method 1

Time used: 1.630 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = \cos(x)$, $B = \sin(x)$, $C = -2 \cos(x)^3$, $f(x) = 2 \cos(x)^5$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$\cos(x) y'' + \sin(x) y' - 2y \cos(x)^3 = 0$$

In normal form the ode

$$\cos(x) y'' + \sin(x) y' - 2y \cos(x)^3 = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= \tan(x) \\ q(x) &= -2 \cos(x)^2 \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c} \sqrt{q} \\ &= \frac{\sqrt{-2 \cos(x)^2}}{c} \\ \tau'' &= \frac{2 \sin(x) \cos(x)}{c \sqrt{-2 \cos(x)^2}} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{2 \sin(x) \cos(x)}{c \sqrt{-2 \cos(x)^2}} + \tan(x) \frac{\sqrt{-2 \cos(x)^2}}{c}}{\left(\frac{\sqrt{-2 \cos(x)^2}}{c} \right)^2} \\ &= 0 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{-2 \cos(x)^2} dx}{c} \\ &= -\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2} c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right) - c_2 \sin\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \cos\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right) \\ y_2 &= -\sin\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right) \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right) & -\sin\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right) \\ \frac{d}{dx}\left(\cos\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right)\right) & \frac{d}{dx}\left(-\sin\left(\frac{\sin(2x)}{\sqrt{-2 \cos(x)^2}}\right)\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) & -\sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \\ -\left(\frac{2\cos(2x)}{\sqrt{-2\cos(x)^2}} - \frac{2\sin(2x)\sin(x)\cos(x)}{(-2\cos(x)^2)^{3/2}}\right) \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) & -\left(\frac{2\cos(2x)}{\sqrt{-2\cos(x)^2}} - \frac{2\sin(2x)\sin(x)\cos(x)}{(-2\cos(x)^2)^{3/2}}\right) \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \end{vmatrix}$$

Therefore

$$W = \left(\cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right) \left(-\left(\frac{2\cos(2x)}{\sqrt{-2\cos(x)^2}} - \frac{2\sin(2x)\sin(x)\cos(x)}{(-2\cos(x)^2)^{3/2}}\right) \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right) \\ - \left(-\sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right) \left(-\left(\frac{2\cos(2x)}{\sqrt{-2\cos(x)^2}} - \frac{2\sin(2x)\sin(x)\cos(x)}{(-2\cos(x)^2)^{3/2}}\right) \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right)$$

Which simplifies to

$$W = \frac{2\cos(x)(\sin(2x)\sin(x) + 2\cos(2x)\cos(x)) \left(\cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right)^2 + \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right)^2 \right)}{(-2\cos(x)^2)^{3/2}}$$

Which simplifies to

$$W = -\frac{\cos(x)^2 \sqrt{2}}{\sqrt{-\cos(x)^2}}$$

Therefore Eq. (2) becomes

$$u_1 = -\int \frac{-2\sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \cos(x)^5}{-\frac{\cos(x)^3 \sqrt{2}}{\sqrt{-\cos(x)^2}}} dx$$

Which simplifies to

$$u_1 = -\int \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \cos(x)^2 \sqrt{2} \sqrt{-\cos(x)^2} dx$$

Hence

$$u_1 = -\int_0^x \sin\left(\frac{\sin(2\alpha)}{\sqrt{-2\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{2} \sqrt{-\cos(\alpha)^2} d\alpha$$

And Eq. (3) becomes

$$u_2 = \int \frac{2\cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \cos(x)^5}{-\frac{\cos(x)^3 \sqrt{2}}{\sqrt{-\cos(x)^2}}} dx$$

Which simplifies to

$$u_2 = \int -\cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \cos(x)^2 \sqrt{2} \sqrt{-\cos(x)^2} dx$$

Hence

$$u_2 = \int_0^x -\cos\left(\frac{\sin(2\alpha)}{\sqrt{-2\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{2} \sqrt{-\cos(\alpha)^2} d\alpha$$

Therefore the particular solution, from equation (1) is

$$\begin{aligned} y_p(x) &= -\int_0^x \sin\left(\frac{\sin(2\alpha)}{\sqrt{-2\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{2} \sqrt{-\cos(\alpha)^2} d\alpha \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \\ &\quad - \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \int_0^x -\cos\left(\frac{\sin(2\alpha)}{\sqrt{-2\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{2} \sqrt{-\cos(\alpha)^2} d\alpha \end{aligned}$$

Which simplifies to

$$\begin{aligned} y_p(x) &= -\sqrt{2} \left(\int_0^x \sin\left(\frac{\sin(2\alpha)\sqrt{2}}{2\sqrt{-\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{-\cos(\alpha)^2} d\alpha \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right. \\ &\quad \left. - \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \int_0^x \cos\left(\frac{\sin(2\alpha)\sqrt{2}}{2\sqrt{-\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{-\cos(\alpha)^2} d\alpha \right) \end{aligned}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) - c_2 \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right) \\ &\quad + \left(-\sqrt{2} \left(\int_0^x \sin\left(\frac{\sin(2\alpha)\sqrt{2}}{2\sqrt{-\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{-\cos(\alpha)^2} d\alpha \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right. \right. \\ &\quad \left. \left. - \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \int_0^x \cos\left(\frac{\sin(2\alpha)\sqrt{2}}{2\sqrt{-\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{-\cos(\alpha)^2} d\alpha \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned} y &= c_1 \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) - c_2 \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \\ &\quad - \sqrt{2} \left(\int_0^x \sin\left(\frac{\sin(2\alpha)\sqrt{2}}{2\sqrt{-\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{-\cos(\alpha)^2} d\alpha \cos\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \right. \\ &\quad \left. - \sin\left(\frac{\sin(2x)}{\sqrt{-2\cos(x)^2}}\right) \int_0^x \cos\left(\frac{\sin(2\alpha)\sqrt{2}}{2\sqrt{-\cos(\alpha)^2}}\right) \cos(\alpha)^2 \sqrt{-\cos(\alpha)^2} d\alpha \right) \end{aligned}$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
trying symmetries linear in x and y(x)
-> Try solving first the homogeneous part of the ODE
    trying a symmetry of the form [xi=0, eta=F(x)]
    <- linear_1 successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.018 (sec)

Leaf size : 30

```

dsolve(cos(x)*diff(diff(y(x),x),x)+sin(x)*diff(y(x),x)-2*y(x)*cos(x)^3 = 2*cos(x)^5,
        y(x),singsol=all)

```

$$y = \sinh\left(\sin(x)\sqrt{2}\right)c_2 + \cosh\left(\sin(x)\sqrt{2}\right)c_1 + \frac{1}{2} - \frac{\cos(2x)}{2}$$

Mathematica DSolve solution

Solving time : 23.596 (sec)

Leaf size : 167

```

DSolve[{Cos[x]*D[y[x],{x,2}]+Sin[x]*D[y[x],x]-2*y[x]*Cos[x]^3==2*Cos[x]^5,{}},
        y[x],x,IncludeSingularSolutions->True]

```

 $y(x)$

$$\begin{aligned}
&\rightarrow \cos\left(\sqrt{-\cos(2x)-1}\tan(x)\right) \int_1^x \cos^2(K[1])\sqrt{-\cos(2K[1])-1}\sin\left(\sqrt{-\cos(2K[1])-1}\tan(K[1])\right) dK[1] \\
&\quad + \sin\left(\sqrt{-\cos(2x)-1}\tan(x)\right) \int_1^x \\
&\quad - \cos^2(K[2])\sqrt{-\cos(2K[2])-1}\cos\left(\sqrt{-\cos(2K[2])-1}\tan(K[2])\right) dK[2] \\
&\quad + c_1 \cos\left(\sqrt{-\cos(2x)-1}\tan(x)\right) + c_2 \sin\left(\sqrt{-\cos(2x)-1}\tan(x)\right)
\end{aligned}$$

2.2.26 problem 26

Solved as second order ode using change of variable on x method 2	705
Solved as second order ode using change of variable on x method 1	710
Maple step by step solution	714
Maple trace	714
Maple dsolve solution	714
Mathematica DSolve solution	714

Internal problem ID [8825]

Book : Second order enumerated odes

Section : section 2

Problem number : 26

Date solved : Thursday, December 12, 2024 at 09:52:54 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' + \left(1 - \frac{1}{x}\right) y' + 4x^2 y e^{-2x} = 4(x^3 + x^2) e^{-3x}$$

Solved as second order ode using change of variable on x method 2

Time used: 1.827 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + \frac{(x-1)y'}{x} + 4x^2 y e^{-2x} = 0$$

In normal form the ode

$$y'' + \frac{(x-1)y'}{x} + 4x^2 y e^{-2x} = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{x-1}{x}$$

$$q(x) = 4x^2 e^{-2x}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\int p(x) dx} dx \\ &= \int e^{-\int \frac{x-1}{x} dx} dx \\ &= \int e^{-x+\ln(x)} dx \\ &= \int x e^{-x} dx \\ &= -(x+1) e^{-x} \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{4x^2 e^{-2x}}{x^2 e^{-2x}} \\ &= 4 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2} y(\tau) + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + 4y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\tau\lambda} + 4 e^{\tau\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +2i \\ \lambda_2 &= -2i \end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 2i \\ \lambda_2 &= -2i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau}(c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0(c_1 \cos(2\tau) + c_2 \sin(2\tau))$$

Or

$$y(\tau) = c_1 \cos(2\tau) + c_2 \sin(2\tau)$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(2(x+1)e^{-x}) - c_2 \sin(2(x+1)e^{-x})$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2(x+1)e^{-x}) - c_2 \sin(2(x+1)e^{-x})$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= -4 \cos(xe^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) - 4 \sin(xe^{-x}) \cos(xe^{-x}) \cos(e^{-x})^2 \\ &\quad + 2 \sin(xe^{-x}) \cos(xe^{-x}) + 2 \sin(e^{-x}) \cos(e^{-x}) \\ y_2 &= 4 \cos(xe^{-x})^2 \cos(e^{-x})^2 - 2 \cos(xe^{-x})^2 - 2 \cos(e^{-x})^2 \\ &\quad + 1 - 4 \sin(xe^{-x}) \cos(xe^{-x}) \sin(e^{-x}) \cos(e^{-x})\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} -4 \cos(x e^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) - 4 \sin(x e^{-x}) \cos(x e^{-x}) \cos(e^{-x})^2 + 2 \sin(x e^{-x}) \cos(x e^{-x}) + 2 \\ \frac{d}{dx} \left(-4 \cos(x e^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) - 4 \sin(x e^{-x}) \cos(x e^{-x}) \cos(e^{-x})^2 + 2 \sin(x e^{-x}) \cos(x e^{-x}) + 2 \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 8 \cos(x e^{-x}) \sin(e^{-x}) \cos(e^{-x}) (e^{-x} - x e^{-x}) \sin(x e^{-x}) + 4 \cos(x e^{-x})^2 e^{-x} \cos(e^{-x})^2 - 4 \cos(x e^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) \end{vmatrix}$$

Therefore

$$\begin{aligned} W = & \left(-4 \cos(x e^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) - 4 \sin(x e^{-x}) \cos(x e^{-x}) \cos(e^{-x})^2 \right. \\ & \left. + 2 \sin(x e^{-x}) \cos(x e^{-x}) \right) \\ & + 2 \sin(e^{-x}) \cos(e^{-x}) \left(-8 \cos(x e^{-x}) \cos(e^{-x})^2 (e^{-x} - x e^{-x}) \sin(x e^{-x}) \right. \\ & + 8 \cos(x e^{-x})^2 \cos(e^{-x}) e^{-x} \sin(e^{-x}) + 4 \cos(x e^{-x}) (e^{-x} - x e^{-x}) \sin(x e^{-x}) \\ & - 4 \cos(e^{-x}) e^{-x} \sin(e^{-x}) - 4(e^{-x} - x e^{-x}) \cos(x e^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) \\ & \left. + 4 \sin(x e^{-x})^2 (e^{-x} - x e^{-x}) \sin(e^{-x}) \cos(e^{-x}) \right) \\ & + 4 \sin(x e^{-x}) \cos(x e^{-x}) e^{-x} \cos(e^{-x})^2 - 4 \sin(x e^{-x}) \cos(x e^{-x}) \sin(e^{-x})^2 e^{-x} \\ & - \left(4 \cos(x e^{-x})^2 \cos(e^{-x})^2 - 2 \cos(x e^{-x})^2 - 2 \cos(e^{-x})^2 + 1 \right. \\ & \left. - 4 \sin(x e^{-x}) \cos(x e^{-x}) \sin(e^{-x}) \cos(e^{-x}) \right) \left(8 \cos(x e^{-x}) \sin(e^{-x}) \cos(e^{-x}) (e^{-x} \right. \\ & - x e^{-x}) \sin(x e^{-x}) + 4 \cos(x e^{-x})^2 e^{-x} \cos(e^{-x})^2 - 4 \cos(x e^{-x})^2 \sin(e^{-x})^2 e^{-x} \\ & - 4(e^{-x} - x e^{-x}) \cos(x e^{-x})^2 \cos(e^{-x})^2 + 4 \sin(x e^{-x})^2 (e^{-x} - x e^{-x}) \cos(e^{-x})^2 \\ & - 8 \sin(x e^{-x}) \cos(x e^{-x}) \cos(e^{-x}) e^{-x} \sin(e^{-x}) + 2(e^{-x} - x e^{-x}) \cos(x e^{-x})^2 \\ & \left. - 2 \sin(x e^{-x})^2 (e^{-x} - x e^{-x}) - 2 e^{-x} \cos(e^{-x})^2 + 2 \sin(e^{-x})^2 e^{-x} \right) \end{aligned}$$

Which simplifies to

$$W = \text{Expression too large to display}$$

Which simplifies to

$$W = -2x e^{-x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{4 \left(4 \cos(x e^{-x})^2 \cos(e^{-x})^2 - 2 \cos(x e^{-x})^2 - 2 \cos(e^{-x})^2 + 1 - 4 \sin(x e^{-x}) \cos(x e^{-x}) \sin(e^{-x}) \cos(e^{-x}) \right)}{-2x e^{-x}} dx$$

Which simplifies to

$$u_1 = - \int 2x(x+1) e^{-2x} (\sin(2e^{-x}) \sin(2x e^{-x}) - \cos(2e^{-x}) \cos(2x e^{-x})) dx$$

Hence

$$u_1 = \frac{i(2x e^{-x} + i + 2e^{-x}) e^{2ie^{-x}} e^{2ix e^{-x}}}{4} - \frac{i(2x e^{-x} - i + 2e^{-x}) e^{-2ie^{-x}} e^{-2ix e^{-x}}}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{4 \left(-4 \cos(x e^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) - 4 \sin(x e^{-x}) \cos(x e^{-x}) \cos(e^{-x})^2 + 2 \sin(x e^{-x}) \cos(x e^{-x}) + \dots \right)}{-2x e^{-x}} dx$$

Which simplifies to

$$u_2 = \int 2x(x+1) e^{-2x} (\sin(2e^{-x}) \cos(2x e^{-x}) + \cos(2e^{-x}) \sin(2x e^{-x})) dx$$

Hence

$$u_2 = \frac{(2x e^{-x} + i + 2e^{-x}) e^{2ie^{-x}} e^{2ix e^{-x}}}{4} + \frac{(2x e^{-x} - i + 2e^{-x}) e^{-2ie^{-x}} e^{-2ix e^{-x}}}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{i(2x e^{-x} + i + 2e^{-x}) e^{2ie^{-x}} e^{2ix e^{-x}}}{4} - \frac{i(2x e^{-x} - i + 2e^{-x}) e^{-2ie^{-x}} e^{-2ix e^{-x}}}{4} \right) \left(-4 \cos(x e^{-x})^2 \sin(e^{-x}) \cos(e^{-x}) - 4 \sin(x e^{-x}) \cos(x e^{-x}) \cos(e^{-x})^2 + 2 \sin(x e^{-x}) \cos(x e^{-x}) + 2 \sin(e^{-x}) \cos(e^{-x}) \right) + \left(4 \cos(x e^{-x})^2 \cos(e^{-x})^2 - 2 \cos(x e^{-x})^2 - 2 \cos(e^{-x})^2 + 1 - 4 \sin(x e^{-x}) \cos(x e^{-x}) \sin(e^{-x}) \cos(e^{-x}) \right) \left(\frac{(2x e^{-x} + i + 2e^{-x}) e^{2ie^{-x}} e^{2ix e^{-x}}}{4} + \frac{(2x e^{-x} - i + 2e^{-x}) e^{-2ie^{-x}} e^{-2ix e^{-x}}}{4} \right)$$

Which simplifies to

$$y_p(x) = (x+1) e^{-x}$$

Therefore the general solution is

$$y = y_h + y_p = (c_1 \cos(2(x+1)e^{-x}) - c_2 \sin(2(x+1)e^{-x})) + ((x+1)e^{-x})$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = (x+1) e^{-x} + c_1 \cos(2(x+1)e^{-x}) - c_2 \sin(2(x+1)e^{-x})$$

Solved as second order ode using change of variable on x method 1

Time used: 0.660 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = \frac{x-1}{x}, C = 4x^2e^{-2x}, f(x) = 4x^2(x+1)e^{-3x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$y'' + \frac{(x-1)y'}{x} + 4x^2ye^{-2x} = 0$$

In normal form the ode

$$y'' + \frac{(x-1)y'}{x} + 4x^2ye^{-2x} = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{x-1}{x}$$

$$q(x) = 4x^2e^{-2x}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{2\sqrt{x^2e^{-2x}}}{c} \quad (6)$$

$$\tau'' = \frac{2xe^{-2x} - 2x^2e^{-2x}}{c\sqrt{x^2e^{-2x}}}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$

$$= \frac{\frac{2xe^{-2x} - 2x^2e^{-2x}}{c\sqrt{x^2e^{-2x}}} + \frac{x-1}{x} \frac{2\sqrt{x^2e^{-2x}}}{c}}{\left(\frac{2\sqrt{x^2e^{-2x}}}{c}\right)^2}$$

$$= 0$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 2\sqrt{x^2 e^{-2x}} dx}{c} \\ &= -\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{cx} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right) - c_2 \sin\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \cos\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right) \\ y_2 &= -\sin\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right) \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right) & -\sin\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right) \\ \frac{d}{dx}\left(\cos\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right)\right) & \frac{d}{dx}\left(-\sin\left(\frac{2(x+1)\sqrt{x^2 e^{-2x}}}{x}\right)\right) \end{vmatrix}$$

Which gives

$$W = \left| \begin{array}{cc} \cos\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) & -\sin\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) \\ -\left(\frac{2\sqrt{x^2e^{-2x}}}{x} + \frac{(x+1)(2xe^{-2x} - 2x^2e^{-2x})}{\sqrt{x^2e^{-2x}}x} - \frac{2(x+1)\sqrt{x^2e^{-2x}}}{x^2}\right) \sin\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) & -\left(\frac{2\sqrt{x^2e^{-2x}}}{x} + \frac{(x+1)(2xe^{-2x} - 2x^2e^{-2x})}{\sqrt{x^2e^{-2x}}x} - \frac{2(x+1)\sqrt{x^2e^{-2x}}}{x^2}\right) \cos\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) \end{array} \right|$$

Therefore

$$W = \left(\cos\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) \right) \left(-\left(\frac{2\sqrt{x^2e^{-2x}}}{x} + \frac{(x+1)(2xe^{-2x} - 2x^2e^{-2x})}{\sqrt{x^2e^{-2x}}x} - \frac{2(x+1)\sqrt{x^2e^{-2x}}}{x^2}\right) \cos\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) \right) \\ - \left(-\sin\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) \right) \left(-\left(\frac{2\sqrt{x^2e^{-2x}}}{x} + \frac{(x+1)(2xe^{-2x} - 2x^2e^{-2x})}{\sqrt{x^2e^{-2x}}x} - \frac{2(x+1)\sqrt{x^2e^{-2x}}}{x^2}\right) \sin\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) \right)$$

Which simplifies to

$$W = \frac{2x^2e^{-2x} \left(\cos\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right)^2 + \sin\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right)^2 \right)}{\sqrt{x^2e^{-2x}}}$$

Which simplifies to

$$W = \frac{2x^2e^{-2x}}{\sqrt{x^2e^{-2x}}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-4 \sin\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) x^2(x+1)e^{-3x}}{\frac{2x^2e^{-2x}}{\sqrt{x^2e^{-2x}}}} dx$$

Which simplifies to

$$u_1 = - \int -2(x+1)e^{-x}\sqrt{x^2e^{-2x}} \sin\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) dx$$

Hence

$$u_1 = - \int_0^x -2(\alpha+1)e^{-\alpha}\sqrt{\alpha^2e^{-2\alpha}} \sin\left(\frac{2(\alpha+1)\sqrt{\alpha^2e^{-2\alpha}}}{\alpha}\right) d\alpha$$

And Eq. (3) becomes

$$u_2 = \int \frac{4 \cos\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) x^2(x+1)e^{-3x}}{\frac{2x^2e^{-2x}}{\sqrt{x^2e^{-2x}}}} dx$$

Which simplifies to

$$u_2 = \int 2(x+1)e^{-x}\sqrt{x^2e^{-2x}} \cos\left(\frac{2(x+1)\sqrt{x^2e^{-2x}}}{x}\right) dx$$

Hence

$$u_2 = \int_0^x 2(\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \cos \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha$$

Therefore the particular solution, from equation (1) is

$$\begin{aligned} y_p(x) = & - \int_0^x -2(\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \sin \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \cos \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \\ & - \sin \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \int_0^x 2(\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \cos \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \end{aligned}$$

Which simplifies to

$$\begin{aligned} y_p(x) = & 2 \int_0^x (\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \sin \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \cos \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \\ & - 2 \sin \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \int_0^x (\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \cos \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \end{aligned}$$

Therefore the general solution is

$$\begin{aligned} y = & y_h + y_p \\ = & \left(c_1 \cos \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) - c_2 \sin \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \right) \\ & + \left(2 \int_0^x (\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \sin \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \cos \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \right. \\ & \left. - 2 \sin \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \int_0^x (\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \cos \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned} y = & c_1 \cos \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) - c_2 \sin \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \\ & + 2 \int_0^x (\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \sin \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \cos \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \\ & - 2 \sin \left(\frac{2(x + 1) \sqrt{x^2 e^{-2x}}}{x} \right) \int_0^x (\alpha + 1) e^{-\alpha \sqrt{\alpha^2 e^{-2\alpha}}} \cos \left(\frac{2(\alpha + 1) \sqrt{\alpha^2 e^{-2\alpha}}}{\alpha} \right) d\alpha \end{aligned}$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
trying symmetries linear in x and y(x)
-> Try solving first the homogeneous part of the ODE
    trying a symmetry of the form [xi=0, eta=F(x)]
    <- linear_1 successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.015 (sec)

Leaf size : 39

```

dsolve(diff(diff(y(x),x),x)+(1-1/x)*diff(y(x),x)+4*x^2*y(x)*exp(-2*x) = 4*(x^3+x^2)*exp(
y(x),singsol=all)

```

$$y = \sin(2(x+1)e^{-x})c_2 + \cos(2(x+1)e^{-x})c_1 + xe^{-x} + e^{-x}$$

Mathematica DSolve solution

Solving time : 0.759 (sec)

Leaf size : 47

```

DSolve[{D[y[x],{x,2}]+(1-1/x)*D[y[x],x]+4*x^2*y[x]*Exp[-2*x]==4*(x^2+x^3)*Exp[-3*x],{}}],
y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow c_1 \cos(2e^{-x}(x+1)) + e^{-x}(x - c_2 e^x \sin(2e^{-x}(x+1)) + 1)$$

2.2.27 problem 27

Solved as second order ode using change of variable on y
 method 2 715
 Solved as second order ode using Kovacic algorithm 719
 Maple step by step solution 726
 Maple trace 726
 Maple dsolve solution 727
 Mathematica DSolve solution 727

Internal problem ID [8826]

Book : Second order enumerated odes

Section : section 2

Problem number : 27

Date solved : Thursday, December 12, 2024 at 09:52:57 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' - x^2y' + xy = x^{m+1}$$

Solved as second order ode using change of variable on y method 2

Time used: 0.905 (sec)

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -x^2, C = x, f(x) = x^{m+1}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$y'' - x^2y' + xy = 0$$

In normal form the ode

$$y'' - x^2y' + xy = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -x^2$$

$$q(x) = x$$

Applying change of variables on the dependnt variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \tag{3}$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - nx + x = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \left(\frac{2}{x} - x^2\right)v'(x) &= 0 \\ v''(x) + \frac{(-x^3 + 2)v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(-x^3 + 2)u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form a linear first order is

$$u'(x) + q(x)u(x) = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{x^3 - 2}{x} \\ p(x) &= 0 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{x^3 - 2}{x} dx} \\ &= x^2 e^{-\frac{x^3}{3}} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx} \mu u &= 0 \\ \frac{d}{dx} \left(u x^2 e^{-\frac{x^3}{3}} \right) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned} u x^2 e^{-\frac{x^3}{3}} &= \int 0 dx + c_1 \\ &= c_1 \end{aligned}$$

Dividing throughout by the integrating factor $x^2 e^{-\frac{x^3}{3}}$ gives the final solution

$$u(x) = \frac{e^{\frac{x^3}{3}} c_1}{x^2}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= \frac{3^{2/3}(-1)^{1/3} c_1 \left(-\frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3})}{(-x^3)^{2/3}} + \frac{3^{3^{1/3}}(-1)^{2/3}e^{\frac{x^3}{3}}}{x} + \frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{(-x^3)^{2/3}} \right)}{9} + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \left(\frac{3^{2/3}(-1)^{1/3} c_1 \left(-\frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3})}{(-x^3)^{2/3}} + \frac{3^{3^{1/3}}(-1)^{2/3}e^{\frac{x^3}{3}}}{x} + \frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{(-x^3)^{2/3}} \right)}{9} + c_2 \right) x \\ &= \frac{(3c_2x - 3e^{\frac{x^3}{3}}c_1)(-x^3)^{2/3} + x^3c_13^{2/3}(\Gamma(\frac{2}{3}) - \Gamma(\frac{2}{3}, -\frac{x^3}{3}))}{3(-x^3)^{2/3}} \end{aligned}$$

Now the particular solution to this ODE is found

$$y'' - x^2y' + xy = x^{m+1}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= x \\ y_2 &= \frac{3^{2/3}x^3\Gamma(\frac{2}{3})}{3(-x^3)^{2/3}} - \frac{3^{2/3}x^3\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{2/3}} - e^{\frac{x^3}{3}} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & \frac{3^{2/3}x^3\Gamma(\frac{2}{3})}{3(-x^3)^{2/3}} - \frac{3^{2/3}x^3\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{2/3}} - e^{\frac{x^3}{3}} \\ \frac{d}{dx}(x) & \frac{d}{dx} \left(\frac{3^{2/3}x^3\Gamma(\frac{2}{3})}{3(-x^3)^{2/3}} - \frac{3^{2/3}x^3\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{2/3}} - e^{\frac{x^3}{3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & \frac{3^{2/3}x^3\Gamma(\frac{2}{3})}{3(-x^3)^{2/3}} - \frac{3^{2/3}x^3\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{2/3}} - e^{\frac{x^3}{3}} \\ 1 & \frac{3^{2/3}x^2\Gamma(\frac{2}{3})}{(-x^3)^{2/3}} + \frac{2 \cdot 3^{2/3}x^5\Gamma(\frac{2}{3})}{3(-x^3)^{5/3}} - \frac{3^{2/3}x^2\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{(-x^3)^{2/3}} - \frac{2 \cdot 3^{2/3}x^5\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{5/3}} - \frac{3^{2/3}x^5e^{\frac{x^3}{3}}}{3(-x^3)^{2/3}(-\frac{x^3}{3})^{1/3}} - x^2e^{\frac{x^3}{3}} \end{vmatrix}$$

Therefore

$$W = (x) \left(\frac{3^{2/3}x^2\Gamma(\frac{2}{3})}{(-x^3)^{2/3}} + \frac{2 \cdot 3^{2/3}x^5\Gamma(\frac{2}{3})}{3(-x^3)^{5/3}} - \frac{3^{2/3}x^2\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{(-x^3)^{2/3}} - \frac{2 \cdot 3^{2/3}x^5\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{5/3}} - \frac{3^{2/3}x^5e^{\frac{x^3}{3}}}{3(-x^3)^{2/3}(-\frac{x^3}{3})^{1/3}} - x^2e^{\frac{x^3}{3}} \right) - \left(\frac{3^{2/3}x^3\Gamma(\frac{2}{3})}{3(-x^3)^{2/3}} - \frac{3^{2/3}x^3\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{2/3}} - e^{\frac{x^3}{3}} \right) \quad (1)$$

Which simplifies to

$$W = - \frac{e^{\frac{x^3}{3}} \left(-3^{2/3}x^9 + 3 \left(-\frac{x^3}{3} \right)^{1/3} (-x^3)^{5/3} x^3 - 3(-x^3)^{5/3} \left(-\frac{x^3}{3} \right)^{1/3} \right)}{3(-x^3)^{5/3} \left(-\frac{x^3}{3} \right)^{1/3}}$$

Which simplifies to

$$W = e^{\frac{x^3}{3}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\left(\frac{3^{2/3}x^3\Gamma(\frac{2}{3})}{3(-x^3)^{2/3}} - \frac{3^{2/3}x^3\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{3(-x^3)^{2/3}} - e^{\frac{x^3}{3}} \right) x^{m+1}}{e^{\frac{x^3}{3}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{m+1} \left(-3(-x^3)^{2/3} + x^3 3^{2/3} e^{-\frac{x^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right)}{3(-x^3)^{2/3}} dx$$

Hence

$$u_1 = - \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{3(-\alpha^3)^{2/3}} d\alpha$$

And Eq. (3) becomes

$$u_2 = \int \frac{x x^{m+1}}{e^{\frac{x^3}{3}}} dx$$

Which simplifies to

$$u_2 = \int x^{m+2} e^{-\frac{x^3}{3}} dx$$

Hence

$$u_2 = \frac{3^{\frac{m}{6}+1} x^m (x^3)^{-\frac{m}{6}} e^{-\frac{x^3}{3}} \text{WhittakerM} \left(\frac{m}{6}, \frac{m}{6} + \frac{1}{2}, \frac{x^3}{3} \right)}{m+3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{3(-\alpha^3)^{2/3}} d\alpha x$$

$$+ \frac{\left(\frac{3^{2/3} x^3 \Gamma\left(\frac{2}{3}\right)}{3(-x^3)^{2/3}} - \frac{3^{2/3} x^3 \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right)}{3(-x^3)^{2/3}} - e^{\frac{x^3}{3}} \right) 3^{\frac{m}{6}+1} x^m (x^3)^{-\frac{m}{6}} e^{-\frac{x^3}{6}} \text{WhittakerM}\left(\frac{m}{6}, \frac{m}{6} + \frac{1}{2}, \frac{x^3}{3}\right)}{m+3}$$

Which simplifies to

$$y_p(x) = \frac{(-x^3)^{2/3} x(m+3) \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{(-\alpha^3)^{2/3}} d\alpha + 9(x^3)^{-\frac{m}{6}} \left((-x^3)^{2/3} e^{\frac{x^3}{6}} x^m 3^{\frac{m}{6}} \right)}{(-x^3)^{2/3} (3m+9)}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(\left(\frac{3^{2/3} (-1)^{1/3} c_1 \left(-\frac{3x^2 (-1)^{2/3} \Gamma\left(\frac{2}{3}\right)}{(-x^3)^{2/3}} + \frac{3 3^{1/3} (-1)^{2/3} e^{\frac{x^3}{3}}}{x} + \frac{3x^2 (-1)^{2/3} \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right)}{(-x^3)^{2/3}} \right)}{9} \right) \right)$$

$$+ c_2 \left(x - \frac{(-x^3)^{2/3} x(m+3) \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{(-\alpha^3)^{2/3}} d\alpha + 9(x^3)^{-\frac{m}{6}} \left((-x^3)^{2/3} e^{\frac{x^3}{6}} x^m 3^{\frac{m}{6}} \right)}{(-x^3)^{2/3} (3m+9)} \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(\frac{3^{2/3} (-1)^{1/3} c_1 \left(-\frac{3x^2 (-1)^{2/3} \Gamma\left(\frac{2}{3}\right)}{(-x^3)^{2/3}} + \frac{3 3^{1/3} (-1)^{2/3} e^{\frac{x^3}{3}}}{x} + \frac{3x^2 (-1)^{2/3} \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right)}{(-x^3)^{2/3}} \right)}{9} \right)$$

$$+ c_2 \left(x - \frac{(-x^3)^{2/3} x(m+3) \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{(-\alpha^3)^{2/3}} d\alpha + 9(x^3)^{-\frac{m}{6}} \left((-x^3)^{2/3} e^{\frac{x^3}{6}} x^m 3^{\frac{m}{6}} \right)}{(-x^3)^{2/3} (3m+9)} \right)$$

Solved as second order ode using Kovacic algorithm

Time used: 0.505 (sec)

Writing the ode as

$$y'' - x^2 y' + xy = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -x^2 \tag{3}$$

$$C = x$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x(x^3 - 8)}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= x(x^3 - 8) \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{x(x^3 - 8)}{4} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.81: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 4 \\ &= -4 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is -4 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Attempting to find a solution using case $n = 1$.

Since the order of r at ∞ is $O_r(\infty) = -4$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{4}{2} = 2$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^2 a_i x^i \end{aligned} \quad (8)$$

Let a be the coefficient of $x^v = x^2$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{x^2}{2} - \frac{2}{x} - \frac{4}{x^4} - \frac{16}{x^7} - \frac{80}{x^{10}} - \frac{448}{x^{13}} - \frac{2688}{x^{16}} - \frac{16896}{x^{19}} + \dots \quad (9)$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 2$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^2 a_i x^i \\ &= \frac{x^2}{2} \end{aligned} \quad (10)$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^1 = x$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{x^4}{4}$$

This shows that the coefficient of x in the above is 0. Now we need to find the coefficient of x in r . How this is done depends on if $v = 0$ or not. Since $v = 2$ which is not zero, then starting $r = \frac{s}{t}$, we do long division and write this in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of x in r will be the coefficient this term in the quotient. Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x(x^3 - 8)}{4} \\ &= Q + \frac{R}{4} \\ &= \left(\frac{1}{4}x^4 - 2x \right) + (0) \\ &= \frac{1}{4}x^4 - 2x \end{aligned}$$

We see that the coefficient of the term $\frac{1}{x}$ in the quotient is -2 . Now b can be found.

$$\begin{aligned} b &= (-2) - (0) \\ &= -2 \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{x^2}{2} \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{-2}{\frac{1}{2}} - 2 \right) = -3 \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{-2}{\frac{1}{2}} - 2 \right) = 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{x(x^3 - 8)}{4}$$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
-4	$\frac{x^2}{2}$	-3	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$, and since there are no poles then

$$\begin{aligned} d &= \alpha_\infty^- \\ &= 1 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= (-)[\sqrt{r}]_\infty \\ &= 0 + (-) \left(\frac{x^2}{2} \right) \\ &= -\frac{x^2}{2} \\ &= -\frac{x^2}{2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 1$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2 \left(-\frac{x^2}{2} \right) (1) + \left((-x) + \left(-\frac{x^2}{2} \right)^2 - \left(\frac{x(x^3 - 8)}{4} \right) \right) = 0$$

$$xa_0 = 0$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 0\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x) e^{\int -\frac{x^2}{2} dx} \\ &= (x) e^{-\frac{x^3}{6}} \\ &= x e^{-\frac{x^3}{6}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x^2}{1} dx} \\ &= z_1 e^{\frac{x^3}{6}} \\ &= z_1 \left(e^{\frac{x^3}{6}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x^2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\frac{x^3}{3}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{3^{2/3}(-1)^{1/3} \left(-\frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3})}{(-x^3)^{2/3}} + \frac{33^{1/3}(-1)^{2/3}e^{\frac{x^3}{3}}}{x} + \frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{(-x^3)^{2/3}} \right)}{9} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(x) + c_2 \left(x \left(\frac{3^{2/3}(-1)^{1/3} \left(-\frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3})}{(-x^3)^{2/3}} + \frac{33^{1/3}(-1)^{2/3}e^{\frac{x^3}{3}}}{x} + \frac{3x^2(-1)^{2/3}\Gamma(\frac{2}{3}, -\frac{x^3}{3})}{(-x^3)^{2/3}} \right)}{9} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - x^2y' + xy = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1x + \frac{c_2 \left(-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right)}{3(-x^3)^{2/3}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = \frac{-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right)}{3(-x^3)^{2/3}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & \frac{-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right)}{3(-x^3)^{2/3}} \\ \frac{d}{dx}(x) & \frac{d}{dx} \left(\frac{-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right)}{3(-x^3)^{2/3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & \frac{-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right)}{3(-x^3)^{2/3}} \\ 1 & \frac{2 \left(-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right) x^2}{3(-x^3)^{5/3}} + \frac{\frac{6e^{\frac{x^3}{3}}}{(-x^3)^{1/3}} x^2 - 3(-x^3)^{2/3} x^2 e^{\frac{x^3}{3}} + 3x^2 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) - \frac{x^5 3^{2/3} e^{\frac{x^3}{3}}}{(-x^3)^{1/3}}}{3(-x^3)^{2/3}} \end{vmatrix}$$

Therefore

$$W = (x) \left(\frac{2 \left(-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right) x^2}{3(-x^3)^{5/3}} \right. \\ \left. + \frac{\frac{6e^{\frac{x^3}{3}} x^2}{(-x^3)^{1/3}} - 3(-x^3)^{2/3} x^2 e^{\frac{x^3}{3}} + 3x^2 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) - \frac{x^5 3^{2/3} e^{\frac{x^3}{3}}}{(-x^3)^{1/3}}}{3(-x^3)^{2/3}} \right) - \left(\frac{-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3}}{3(-x^3)^{5/3}} \right)$$

Which simplifies to

$$W = \frac{e^{\frac{x^3}{3}} \left(3^{2/3} x^9 - 3 \left(-\frac{x^3}{3} \right)^{1/3} (-x^3)^{5/3} x^3 + 3(-x^3)^{5/3} \left(-\frac{x^3}{3} \right)^{1/3} \right)}{3(-x^3)^{5/3} \left(-\frac{x^3}{3} \right)^{1/3}}$$

Which simplifies to

$$W = e^{\frac{x^3}{3}}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\left(-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right) x^{m+1}}{\frac{3(-x^3)^{2/3}}{e^{\frac{x^3}{3}}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{m+1} \left(-3(-x^3)^{2/3} + x^3 3^{2/3} e^{-\frac{x^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right)}{3(-x^3)^{2/3}} dx$$

Hence

$$u_1 = - \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{3(-\alpha^3)^{2/3}} d\alpha$$

And Eq. (3) becomes

$$u_2 = \int \frac{x x^{m+1}}{e^{\frac{x^3}{3}}} dx$$

Which simplifies to

$$u_2 = \int x^{m+2} e^{-\frac{x^3}{3}} dx$$

Hence

$$u_2 = \frac{3^{\frac{m}{6}+1} x^m (x^3)^{-\frac{m}{6}} e^{-\frac{x^3}{6}} \text{WhittakerM} \left(\frac{m}{6}, \frac{m}{6} + \frac{1}{2}, \frac{x^3}{3} \right)}{m+3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{3(-\alpha^3)^{2/3}} d\alpha x \\ + \frac{\left(-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right) 3^{\frac{m}{6}+1} x^m (x^3)^{-\frac{m}{6}} e^{-\frac{x^3}{6}} \text{WhittakerM} \left(\frac{m}{6}, \frac{m}{6} + \frac{1}{2}, \frac{x^3}{3} \right)}{3(-x^3)^{2/3} (m+3)}$$

Which simplifies to

$$y_p(x) = \frac{(-x^3)^{2/3} x(m+3) \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{(-\alpha^3)^{2/3}} d\alpha + 9(x^3)^{-\frac{m}{6}} \left((-x^3)^{2/3} e^{\frac{x^3}{6}} x^m 3^{\frac{m}{6}} + 3 \right)}{(-x^3)^{2/3} (3m+9)}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 x + \frac{c_2 \left(-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right)}{3(-x^3)^{2/3}} \right) + \left(\frac{(-x^3)^{2/3} x(m+3) \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{(-\alpha^3)^{2/3}} d\alpha + 9(x^3)^{-\frac{m}{6}} \left((-x^3)^{2/3} e^{\frac{x^3}{6}} x^m 3^{\frac{m}{6}} + 3 \right)}{(-x^3)^{2/3} (3m+9)} \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 x + \frac{c_2 \left(-3(-x^3)^{2/3} e^{\frac{x^3}{3}} + x^3 3^{2/3} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right)}{3(-x^3)^{2/3}} + \frac{(-x^3)^{2/3} x(m+3) \int_0^x \frac{\alpha^{m+1} \left(-3(-\alpha^3)^{2/3} + \alpha^3 3^{2/3} e^{-\frac{\alpha^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{\alpha^3}{3}\right) \right) \right)}{(-\alpha^3)^{2/3}} d\alpha + 9(x^3)^{-\frac{m}{6}} \left((-x^3)^{2/3} e^{\frac{x^3}{6}} x^m 3^{\frac{m}{6}} + 3 \right)}{(-x^3)^{2/3} (3m+9)}$$

Maple step by step solution

Maple trace

```

Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
    <- Kovacic's algorithm successful
<- solving first the homogeneous part of the ODE successful

```

Maple dsolve solution

Solving time : 0.042 (sec)

Leaf size : 210

```
dsolve(diff(diff(y(x),x),x)-diff(y(x),x)*x^2+x*y(x) = x^(m+1),
        y(x),singsol=all)
```

$$y = \frac{(-x^3)^{2/3} x^{m+3} \left(\int -\frac{x^{m+1} \left(-3(-x^3)^{2/3} + x^3 2^{2/3} e^{-\frac{x^3}{3}} \left(\Gamma\left(\frac{2}{3}\right) - \Gamma\left(\frac{2}{3}, -\frac{x^3}{3}\right) \right) \right)}{(-x^3)^{2/3}} dx \right)}{3} - \frac{\left(-9(-x^3)^{2/3} e^{\frac{x^3}{6}} x^m 3^{\frac{m}{6}} + x^{m+3} 3^{\frac{5}{6} + \frac{m}{6}} e^{-\dots} \right)}{3}$$

Mathematica DSolve solution

Solving time : 0.705 (sec)

Leaf size : 144

```
DSolve[{D[y[x],{x,2}]-x^2*D[y[x],x]+x*y[x]==x^(m+1),{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x \int_1^x \frac{e^{-\frac{1}{3}K[1]^3} \Gamma\left(-\frac{1}{3}, -\frac{1}{3}K[1]^3\right) K[1]^{m+1} \sqrt[3]{-K[1]^3}}{3\sqrt[3]{3}} dK[1] - \frac{\sqrt[3]{-x^3} (x^3)^{-m/3} \Gamma\left(-\frac{1}{3}, -\frac{x^3}{3}\right) \left(-3^{m/3} x^m \Gamma\left(\frac{m+3}{3}, \frac{x^3}{3}\right) + c_2 (x^3)^{m/3} \right)}{3\sqrt[3]{3}} + c_1 x$$

2.2.28 problem 28

Solved as second order ode using Kovacic algorithm 728
 Solved as second order ode adjoint method 732
 Maple step by step solution 737
 Maple trace 737
 Maple dsolve solution 737
 Mathematica DSolve solution 737

Internal problem ID [8827]

Book : Second order enumerated odes

Section : section 2

Problem number : 28

Date solved : Thursday, December 12, 2024 at 09:53:00 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$y'' - \frac{y'}{\sqrt{x}} + \frac{y(-8 + \sqrt{x} + x)}{4x^2} = 0$$

Solved as second order ode using Kovacic algorithm

Time used: 0.105 (sec)

Writing the ode as

$$y'' - \frac{y'}{\sqrt{x}} + \left(-\frac{2}{x^2} + \frac{1}{4x^{3/2}} + \frac{1}{4x}\right)y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -\frac{1}{\sqrt{x}} \\ C &= -\frac{2}{x^2} + \frac{1}{4x^{3/2}} + \frac{1}{4x} \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.82: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{x} + (-)(0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \tag{1A}$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{\sqrt{x}} dx} \\ &= z_1 e^{\sqrt{x}} \\ &= z_1 \left(e^{\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{e^{\sqrt{x}}}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{\sqrt{x}} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2\sqrt{x}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{e^{\sqrt{x}}}{x} \right) + c_2 \left(\frac{e^{\sqrt{x}}}{x} \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 e^{\sqrt{x}}}{x} + \frac{c_2 x^2 e^{\sqrt{x}}}{3}$$

Solved as second order ode adjoint method

Time used: 0.949 (sec)

In normal form the ode

$$y'' - \frac{y'}{\sqrt{x}} + \frac{y(-8 + \sqrt{x} + x)}{4x^2} = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{1}{\sqrt{x}} \\ q(x) &= -\frac{2}{x^2} + \frac{1}{4x^{3/2}} + \frac{1}{4x} \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(-\frac{\xi(x)}{\sqrt{x}}\right)' + \left(\left(-\frac{2}{x^2} + \frac{1}{4x^{3/2}} + \frac{1}{4x}\right)\xi(x)\right) &= 0 \\ \xi''(x) + \frac{\xi'(x)}{\sqrt{x}} - \frac{(\sqrt{x} - x + 8)\xi(x)}{4x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi'' + \frac{\xi'}{\sqrt{x}} + \left(-\frac{1}{4x^{3/2}} - \frac{2}{x^2} + \frac{1}{4x}\right)\xi = 0 \quad (1)$$

$$A\xi'' + B\xi' + C\xi = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= \frac{1}{\sqrt{x}} \\ C &= -\frac{1}{4x^{3/2}} - \frac{2}{x^2} + \frac{1}{4x} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = \xi e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then ξ is found using the inverse transformation

$$\xi = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.83: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\text{gcd}(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{x} + (-)(0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \tag{1A}$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

The first solution to the original ode in ξ is found from

$$\begin{aligned} \xi_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{\sqrt{x}} dx} \\ &= z_1 e^{-\sqrt{x}} \\ &= z_1 \left(e^{-\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$\xi_1 = \frac{e^{-\sqrt{x}}}{x}$$

The second solution ξ_2 to the original ode is found using reduction of order

$$\xi_2 = \xi_1 \int \frac{e^{\int -\frac{B}{A} dx}}{\xi_1^2} dx$$

Substituting gives

$$\begin{aligned} \xi_2 &= \xi_1 \int \frac{e^{\int -\frac{1}{2} \frac{1}{\sqrt{x}} dx}}{(\xi_1)^2} dx \\ &= \xi_1 \int \frac{e^{-2\sqrt{x}}}{(\xi_1)^2} dx \\ &= \xi_1 \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} \xi &= c_1 \xi_1 + c_2 \xi_2 \\ &= c_1 \left(\frac{e^{-\sqrt{x}}}{x} \right) + c_2 \left(\frac{e^{-\sqrt{x}}}{x} \left(\frac{x^3 e^{2\sqrt{x}} e^{-2\sqrt{x}}}{3} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(-\frac{1}{\sqrt{x}} - \frac{-\frac{c_1 e^{-\sqrt{x}}}{2x^{3/2}} - \frac{c_1 e^{-\sqrt{x}}}{x^2} + \frac{2c_2 x e^{-\sqrt{x}}}{3} - \frac{c_2 x^{3/2} e^{-\sqrt{x}}}{6}}{\frac{c_1 e^{-\sqrt{x}}}{x} + \frac{c_2 x^2 e^{-\sqrt{x}}}{3}} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{4c_2 x^{7/2} - 6c_1 \sqrt{x} + x(c_2 x^3 + 3c_1)}{x^{3/2} (2c_2 x^3 + 6c_1)} \\ p(x) &= 0\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{4c_2 x^{7/2} - 6c_1 \sqrt{x} + x(c_2 x^3 + 3c_1)}{x^{3/2} (2c_2 x^3 + 6c_1)} dx} \\ &= \frac{x e^{-\sqrt{x}}}{c_2 x^3 + 3c_1}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{yx e^{-\sqrt{x}}}{c_2 x^3 + 3c_1} \right) &= 0\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{yx e^{-\sqrt{x}}}{c_2 x^3 + 3c_1} &= \int 0 dx + c_3 \\ &= c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{x e^{-\sqrt{x}}}{c_2 x^3 + 3c_1}$ gives the final solution

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}} c_3}{x}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}} c_3}{x}$$

The constants can be merged to give

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}}}{x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{(c_2 x^3 + 3c_1) e^{\sqrt{x}}}{x}$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
<- Kovacic's algorithm successful`

```

Maple dsolve solution

Solving time : 0.007 (sec)

Leaf size : 19

```

dsolve(diff(diff(y(x),x),x)-1/x^(1/2)*diff(y(x),x)+1/4/x^2*(x+x^(1/2)-8)*y(x) = 0,
        y(x),singsol=all)

```

$$y = \frac{e^{\sqrt{x}}(c_2x^3 + c_1)}{x}$$

Mathematica DSolve solution

Solving time : 0.057 (sec)

Leaf size : 30

```

DSolve[{D[y[x],{x,2}]-1/x^(1/2)*D[y[x],x]+y[x]/(4*x^2)*(-8+x^(1/2)+x)==0,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{e^{\sqrt{x}}(c_2x^3 + 3c_1)}{3x}$$

2.2.29 problem 29

Solved as second order ode using change of variable on y
 method 1 738
 Solved as second order ode adjoint method 740
 Maple step by step solution 743
 Maple trace 743
 Maple dsolve solution 744
 Mathematica DSolve solution 744

Internal problem ID [8828]

Book : Second order enumerated odes

Section : section 2

Problem number : 29

Date solved : Thursday, December 12, 2024 at 09:53:01 AM

CAS classification : [_Lienard]

Solve

$$\cos(x)^2 y'' - 2 \cos(x) \sin(x) y' + y \cos(x)^2 = 0$$

Solved as second order ode using change of variable on y method 1

Time used: 0.460 (sec)

In normal form the given ode is written as

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$p(x) = -\frac{\sin(2x)}{\cos(x)^2}$$

$$q(x) = 1$$

Calculating the Liouville ode invariant Q given by

$$Q = q - \frac{p'}{2} - \frac{p^2}{4}$$

$$= 1 - \frac{\left(-\frac{\sin(2x)}{\cos(x)^2}\right)'}{2} - \frac{\left(-\frac{\sin(2x)}{\cos(x)^2}\right)^2}{4}$$

$$= 1 - \frac{\left(-\frac{2 \cos(2x)}{\cos(x)^2} - \frac{2 \sin(2x) \sin(x)}{\cos(x)^3}\right)}{2} - \frac{\left(\frac{\sin(2x)^2}{\cos(x)^4}\right)}{4}$$

$$= 1 - \left(-\frac{\cos(2x)}{\cos(x)^2} - \frac{\sin(2x) \sin(x)}{\cos(x)^3}\right) - \frac{\sin(2x)^2}{4 \cos(x)^4}$$

$$= 2$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x) z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$z(x) = e^{-\int \frac{p(x)}{2} dx}$$

$$= e^{-\int \frac{-\frac{\sin(2x)}{\cos(x)^2}}{2} dx}$$

$$= \sec(x) \tag{5}$$

Hence (3) becomes

$$y = v(x) \sec(x) \quad (4)$$

Applying this change of variable to the original ode results in

$$\cos(x) (2v(x) + v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$2v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 2$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + 2e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(2)} \\ &= \pm i\sqrt{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{2}$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x))$$

Or

$$v(x) = c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= \left(c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x) \right) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \sec(x)$$

Hence (7) becomes

$$y = \left(c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x) \right) \sec(x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x) \right) \sec(x)$$

Solved as second order ode adjoint method

Time used: 2.226 (sec)

In normal form the ode

$$\cos(x)^2 y'' - 2 \cos(x) \sin(x) y' + y \cos(x)^2 = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -2 \tan(x) \\ q(x) &= 1 \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (-2 \tan(x) \xi(x))' + (\xi(x)) &= 0 \\ 2 \tan(x) \xi'(x) + \xi(x) + \xi''(x) + 2 \sec(x)^2 \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x) \xi' + q(x) \xi = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= 2 \tan(x) \\ q(x) &= 2 \sec(x)^2 + 1 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 2 \sec(x)^2 + 1 - \frac{(2 \tan(x))'}{2} - \frac{(2 \tan(x))^2}{4} \\ &= 2 \sec(x)^2 + 1 - \frac{(2 + 2 \tan(x)^2)}{2} - \frac{(4 \tan(x)^2)}{4} \\ &= 2 \sec(x)^2 + 1 - (1 + \tan(x)^2) - \tan(x)^2 \\ &= 2 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x) z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{2 \tan(x)}{2} dx} \\ &= \cos(x) \end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = v(x) \cos(x) \quad (4)$$

Applying this change of variable to the original ode results in

$$(2v(x) + v''(x)) \cos(x) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$2v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 2$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + 2e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(2)} \\ &= \pm i\sqrt{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{2}$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0(c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x))$$

Or

$$v(x) = c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} \xi &= v(x)z(x) \\ &= (c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x))z(x) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \cos(x)$$

Hence (7) becomes

$$\xi = (c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)) \cos(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x)y' - y\xi'(x) + \xi(x)p(x)y &= \int \xi(x)r(x) dx \\ y' + y\left(p(x) - \frac{\xi'(x)}{\xi(x)}\right) &= \frac{\int \xi(x)r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y\left(-2 \tan(x) - \frac{(-c_1\sqrt{2} \sin(\sqrt{2}x) + c_2\sqrt{2} \cos(\sqrt{2}x)) \cos(x) - (c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)) \sin(x)}{(c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)) \cos(x)}\right)$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{(-c_1\sqrt{2} + \tan(x)c_2) \sin(\sqrt{2}x) + (\tan(x)c_1 + c_2\sqrt{2}) \cos(\sqrt{2}x)}{c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(-c_1\sqrt{2} + \tan(x)c_2) \sin(\sqrt{2}x) + (\tan(x)c_1 + c_2\sqrt{2}) \cos(\sqrt{2}x)}{c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)} dx} \\ &= e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{2}x}{2}\right)^2\right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx} \left(y e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) + \ln\left(1+\tan\left(\frac{\sqrt{2}x}{2}\right)^2\right)} \right) = 0$$

Integrating gives

$$y e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) + \ln\left(1+\tan\left(\frac{\sqrt{2}x}{2}\right)^2\right)} = \int 0 dx + c_3 = c_3$$

Dividing throughout by the integrating factor $e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) + \ln\left(1+\tan\left(\frac{\sqrt{2}x}{2}\right)^2\right)}$ gives the final solution

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2} c_3}{1 + \tan\left(\frac{\sqrt{2}x}{2}\right)^2}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2} c_3}{1 + \tan\left(\frac{\sqrt{2}x}{2}\right)^2}$$

The constants can be merged to give

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2}}{1 + \tan\left(\frac{\sqrt{2}x}{2}\right)^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2}}{1 + \tan\left(\frac{\sqrt{2}x}{2}\right)^2}$$

Maple step by step solution

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`
```

Maple dsolve solution

Solving time : 0.007 (sec)

Leaf size : 24

```
dsolve(cos(x)^2*diff(diff(y(x),x),x)-2*cos(x)*sin(x)*diff(y(x),x)+cos(x)^2*y(x) = 0,
y(x),singsol=all)
```

$$y = \sec(x) \left(c_1 \sin(\sqrt{2}x) + c_2 \cos(\sqrt{2}x) \right)$$

Mathematica DSolve solution

Solving time : 0.096 (sec)

Leaf size : 51

```
DSolve[{Cos[x]^2*D[y[x],{x,2}]-2*Cos[x]*Sin[x]*D[y[x],x]+y[x]*Cos[x]^2==0,{},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{4} e^{-i\sqrt{2}x} \left(4c_1 - i\sqrt{2}c_2 e^{2i\sqrt{2}x} \right) \sec(x)$$

2.2.30 problem 30

Solved as second order ode using change of variable on y	
method 1	745
Solved as second order ode using Kovacic algorithm	749
Solved as second order ode adjoint method	752
Maple step by step solution	756
Maple trace	756
Maple dsolve solution	756
Mathematica DSolve solution	757

Internal problem ID [8829]

Book : Second order enumerated odes

Section : section 2

Problem number : 30

Date solved : Thursday, December 12, 2024 at 09:53:05 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' - 4xy' + (4x^2 - 1)y = -3e^{x^2} \sin(x)$$

Solved as second order ode using change of variable on y method 1

Time used: 0.482 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4xy' + (4x^2 - 1)y = 0$$

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$\begin{aligned} p(x) &= -4x \\ q(x) &= 4x^2 - 1 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 4x^2 - 1 - \frac{(-4x)'}{2} - \frac{(-4x)^2}{4} \\ &= 4x^2 - 1 - \frac{(-4)}{2} - \frac{(16x^2)}{4} \\ &= 4x^2 - 1 - (-2) - 4x^2 \\ &= 1 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{-4x}{2}} \\ &= e^{x^2} \end{aligned} \tag{5}$$

Hence (3) becomes

$$y = v(x) e^{x^2} \tag{4}$$

Applying this change of variable to the original ode results in

$$e^{x^2}(v(x) + v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0(c_1 \cos(x) + c_2 \sin(x))$$

Or

$$v(x) = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 \cos(x) + c_2 \sin(x)) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = e^{x^2}$$

Hence (7) becomes

$$y = (c_1 \cos(x) + c_2 \sin(x)) e^{x^2}$$

Therefore the homogeneous solution y_h is

$$y_h = (c_1 \cos(x) + c_2 \sin(x)) e^{x^2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x) e^{x^2}$$

$$y_2 = e^{x^2} \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) e^{x^2} & e^{x^2} \sin(x) \\ \frac{d}{dx}(\cos(x) e^{x^2}) & \frac{d}{dx}(e^{x^2} \sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) e^{x^2} & e^{x^2} \sin(x) \\ -e^{x^2} \sin(x) + 2 \cos(x) x e^{x^2} & 2x e^{x^2} \sin(x) + \cos(x) e^{x^2} \end{vmatrix}$$

Therefore

$$W = (\cos(x) e^{x^2}) (2x e^{x^2} \sin(x) + \cos(x) e^{x^2}) - (e^{x^2} \sin(x)) (-e^{x^2} \sin(x) + 2 \cos(x) x e^{x^2})$$

Which simplifies to

$$W = e^{2x^2} \cos(x)^2 + e^{2x^2} \sin(x)^2$$

Which simplifies to

$$W = e^{2x^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-3 e^{2x^2} \sin(x)^2}{e^{2x^2}} dx$$

Which simplifies to

$$u_1 = - \int -3 \sin(x)^2 dx$$

Hence

$$u_1 = - \frac{3 \sin(x) \cos(x)}{2} + \frac{3x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{-3 \cos(x) e^{2x^2} \sin(x)}{e^{2x^2}} dx$$

Which simplifies to

$$u_2 = \int -\frac{3 \sin(2x)}{2} dx$$

Hence

$$u_2 = \frac{3 \cos(2x)}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{3 \sin(x) \cos(x)}{2} + \frac{3x}{2} \right) \cos(x) e^{x^2} + \frac{3 e^{x^2} \sin(x) \cos(2x)}{4}$$

Which simplifies to

$$y_p(x) = \frac{3 e^{x^2} (-\sin(x) + 2 \cos(x) x)}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left((c_1 \cos(x) + c_2 \sin(x)) e^{x^2} \right) + \left(\frac{3 e^{x^2} (-\sin(x) + 2 \cos(x) x)}{4} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = (c_1 \cos(x) + c_2 \sin(x)) e^{x^2} + \frac{3 e^{x^2} (-\sin(x) + 2 \cos(x) x)}{4}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.269 (sec)

Writing the ode as

$$y'' - 4xy' + (4x^2 - 1)y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -4x \\ C &= 4x^2 - 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.84: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-4x}{1} dx} \\ &= z_1 e^{x^2} \\ &= z_1 (e^{x^2}) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x) e^{x^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-4x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x^2}}{(y_1)^2} dx \\ &= y_1 (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x) e^{x^2}) + c_2 (\cos(x) e^{x^2} (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4xy' + (4x^2 - 1)y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) e^{x^2} + c_2 e^{x^2} \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \cos(x) e^{x^2} \\ y_2 &= e^{x^2} \sin(x) \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) e^{x^2} & e^{x^2} \sin(x) \\ \frac{d}{dx}(\cos(x) e^{x^2}) & \frac{d}{dx}(e^{x^2} \sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) e^{x^2} & e^{x^2} \sin(x) \\ -e^{x^2} \sin(x) + 2 \cos(x) x e^{x^2} & 2x e^{x^2} \sin(x) + \cos(x) e^{x^2} \end{vmatrix}$$

Therefore

$$W = (\cos(x) e^{x^2}) (2x e^{x^2} \sin(x) + \cos(x) e^{x^2}) - (e^{x^2} \sin(x)) (-e^{x^2} \sin(x) + 2 \cos(x) x e^{x^2})$$

Which simplifies to

$$W = e^{2x^2} \cos(x)^2 + e^{2x^2} \sin(x)^2$$

Which simplifies to

$$W = e^{2x^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-3 e^{2x^2} \sin(x)^2}{e^{2x^2}} dx$$

Which simplifies to

$$u_1 = - \int -3 \sin(x)^2 dx$$

Hence

$$u_1 = -\frac{3 \sin(x) \cos(x)}{2} + \frac{3x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{-3 \cos(x) e^{2x^2} \sin(x)}{e^{2x^2}} dx$$

Which simplifies to

$$u_2 = \int -\frac{3 \sin(2x)}{2} dx$$

Hence

$$u_2 = \frac{3 \cos(2x)}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{3 \sin(x) \cos(x)}{2} + \frac{3x}{2} \right) \cos(x) e^{x^2} + \frac{3 e^{x^2} \sin(x) \cos(2x)}{4}$$

Which simplifies to

$$y_p(x) = \frac{3 e^{x^2} (-\sin(x) + 2 \cos(x) x)}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(x) e^{x^2} + c_2 e^{x^2} \sin(x) \right) + \left(\frac{3 e^{x^2} (-\sin(x) + 2 \cos(x) x)}{4} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos(x) e^{x^2} + c_2 e^{x^2} \sin(x) + \frac{3 e^{x^2} (-\sin(x) + 2 \cos(x) x)}{4}$$

Solved as second order ode adjoint method

Time used: 4.780 (sec)

In normal form the ode

$$y'' - 4xy' + (4x^2 - 1)y = -3e^{x^2} \sin(x) \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -4x \\ q(x) &= 4x^2 - 1 \\ r(x) &= -3e^{x^2} \sin(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned}\xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (-4x\xi(x))' + ((4x^2 - 1)\xi(x)) &= 0 \\ 4\xi(x)x^2 + 4x\xi'(x) + \xi''(x) + 3\xi(x) &= 0\end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x)\xi' + q(x)\xi = 0 \quad (2)$$

Where

$$\begin{aligned}p(x) &= 4x \\ q(x) &= 4x^2 + 3\end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned}Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 4x^2 + 3 - \frac{(4x)'}{2} - \frac{(4x)^2}{4} \\ &= 4x^2 + 3 - \frac{(4)}{2} - \frac{(16x^2)}{4} \\ &= 4x^2 + 3 - (2) - 4x^2 \\ &= 1\end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned}z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{4x}{2} dx} \\ &= e^{-x^2}\end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = v(x)e^{-x^2} \quad (4)$$

Applying this change of variable to the original ode results in

$$e^{-x^2}(v''(x) + v(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$v''(x) + v(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$v(x) = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} \xi &= v(x) z(x) \\ &= (c_1 \cos(x) + c_2 \sin(x)) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = e^{-x^2}$$

Hence (7) becomes

$$\xi = (c_1 \cos(x) + c_2 \sin(x)) e^{-x^2}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(-4x - \frac{((-c_1 \sin(x) + c_2 \cos(x)) e^{-x^2} - 2(c_1 \cos(x) + c_2 \sin(x)) x e^{-x^2}) e^{x^2}}{c_1 \cos(x) + c_2 \sin(x)} \right) = \frac{e^{x^2} \left(\frac{3 \cos(x)^2 c_1 - 3 \cos(x) c_2}{2} - c_2 x \right)}{c_1 \cos(x) + c_2 \sin(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{\sin(x)(2c_2x - c_1) + 2(c_1x + \frac{c_2}{2})\cos(x)}{c_1 \cos(x) + c_2 \sin(x)}$$

$$p(x) = \frac{3(\cos(x)^2 c_1 + c_2 \sin(x) \cos(x) - c_2 x) e^{x^2}}{2c_1 \cos(x) + 2c_2 \sin(x)}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{\sin(x)(2c_2x - c_1) + 2(c_1x + \frac{c_2}{2})\cos(x)}{c_1 \cos(x) + c_2 \sin(x)} dx} \\ &= \frac{\sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{3(\cos(x)^2 c_1 + c_2 \sin(x) \cos(x) - c_2 x) e^{x^2}}{2c_1 \cos(x) + 2c_2 \sin(x)} \right) \\ \frac{d}{dx} \left(\frac{y \sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1} \right) &= \left(\frac{\sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1} \right) \left(\frac{3(\cos(x)^2 c_1 + c_2 \sin(x) \cos(x) - c_2 x) e^{x^2}}{2c_1 \cos(x) + 2c_2 \sin(x)} \right) \\ d \left(\frac{y \sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1} \right) &= \left(\frac{3(\cos(x)^2 c_1 + c_2 \sin(x) \cos(x) - c_2 x) e^{x^2} \sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{(2c_1 \cos(x) + 2c_2 \sin(x)) (c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1)} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y \sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1} &= \int \frac{3(\cos(x)^2 c_1 + c_2 \sin(x) \cos(x) - c_2 x) e^{x^2} \sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{(2c_1 \cos(x) + 2c_2 \sin(x)) (c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1)} dx \\ &= \frac{6x}{4ic_2 - 4c_1} - \frac{3ic_2x}{(ic_1 + c_2)(e^{2ix}c_2 + ic_1 e^{2ix} - c_2 + ic_1)} + c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{\sec\left(\frac{x}{2}\right)^2 e^{-x^2}}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1}$ gives the final solution

$$y = -\frac{(c_1 \cos(x) + c_2 \sin(x)) e^{x^2} (-2e^{2ix} c_1^2 c_3 + 2e^{2ix} c_2^2 c_3 + 3e^{2ix} c_1 x - 2c_1^2 c_3 - 2c_2^2 c_3 + 4ie^{2ix} c_1 c_2 c_3 + 3c_1 x)}{2(ic_1 + c_2)(e^{2ix}c_2 + ic_1 e^{2ix} - c_2 + ic_1)}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{(c_1 \cos(x) + c_2 \sin(x)) e^{x^2} (-2 e^{2ix} c_1^2 c_3 + 2 e^{2ix} c_2^2 c_3 + 3 e^{2ix} c_1 x - 2 c_1^2 c_3 - 2 c_2^2 c_3 + 4 i e^{2ix} c_1 c_2 c_3 + 3 c_1 x - 3 i e^{2ix} c_2 x - 3 i e^{2ix} c_2 c_3)}{2 (i c_1 + c_2) (e^{2ix} c_2 + i c_1 e^{2ix} - c_2 + i c_1)}$$

The constants can be merged to give

$$y = \frac{(c_1 \cos(x) + c_2 \sin(x)) e^{x^2} (-2 e^{2ix} c_1^2 + 2 e^{2ix} c_2^2 + 3 e^{2ix} c_1 x - 2 c_1^2 - 2 c_2^2 + 4 i e^{2ix} c_1 c_2 + 3 c_1 x - 3 i e^{2ix} c_2 x - 3 i e^{2ix} c_2 c_3)}{2 (i c_1 + c_2) (e^{2ix} c_2 + i c_1 e^{2ix} - c_2 + i c_1)}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{(c_1 \cos(x) + c_2 \sin(x)) e^{x^2} (-2 e^{2ix} c_1^2 + 2 e^{2ix} c_2^2 + 3 e^{2ix} c_1 x - 2 c_1^2 - 2 c_2^2 + 4 i e^{2ix} c_1 c_2 + 3 c_1 x - 3 i e^{2ix} c_2 x - 3 i e^{2ix} c_2 c_3)}{2 (i c_1 + c_2) (e^{2ix} c_2 + i c_1 e^{2ix} - c_2 + i c_1)}$$

Maple step by step solution

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacic's algorithm successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.024 (sec)

Leaf size : 29

```

dsolve(diff(diff(y(x), x), x) - 4*diff(y(x), x)*x + (4*x^2 - 1)*y(x) = -3*exp(x^2)*sin(x),
        y(x), singsol=all)

```

$$y = \frac{((3x + 2c_2) \cos(x) + \sin(x) (2c_1 - 3)) e^{x^2}}{2}$$

Mathematica DSolve solution

Solving time : 0.113 (sec)

Leaf size : 50

```
DSolve[{D[y[x],{x,2}]-4*x*D[y[x],x]+(4*x^2-1)*y[x]==-3*Exp[x^2]*Sin[x],{}}],  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{8} e^{x(x-i)} (6x + e^{2ix} (6x + 3i - 4ic_2) - 3i + 8c_1)$$

2.2.31 problem 31

Solved as second order ode using change of variable on y	
method 1	758
Solved as second order ode using Kovacic algorithm	762
Solved as second order ode adjoint method	766
Maple step by step solution	770
Maple trace	770
Maple dsolve solution	770
Mathematica DSolve solution	770

Internal problem ID [8830]

Book : Second order enumerated odes

Section : section 2

Problem number : 31

Date solved : Thursday, December 12, 2024 at 09:53:11 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' - 2bxy' + b^2x^2y = x$$

Solved as second order ode using change of variable on y method 1

Time used: 0.633 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2bxy' + b^2x^2y = 0$$

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -2xb$$

$$q(x) = x^2b^2$$

Calculating the Liouville ode invariant Q given by

$$Q = q - \frac{p'}{2} - \frac{p^2}{4}$$

$$= x^2b^2 - \frac{(-2xb)'}{2} - \frac{(-2xb)^2}{4}$$

$$= x^2b^2 - \frac{(-2b)}{2} - \frac{(4x^2b^2)}{4}$$

$$= x^2b^2 - (-b) - x^2b^2$$

$$= b$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{-2xb}{2} dx} \\ &= e^{\frac{bx^2}{2}} \end{aligned} \quad (5)$$

Hence (3) becomes

$$y = v(x) e^{\frac{bx^2}{2}} \quad (4)$$

Applying this change of variable to the original ode results in

$$e^{\frac{bx^2}{2}} (bv(x) + v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$bv(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = b$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + b e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + b = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = b$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(b)} \\ &= \pm \sqrt{-b} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +\sqrt{-b} \\ \lambda_2 &= -\sqrt{-b} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \sqrt{-b} \\ \lambda_2 &= -\sqrt{-b} \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} v(x) &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ v(x) &= c_1 e^{(\sqrt{-b})x} + c_2 e^{(-\sqrt{-b})x} \end{aligned}$$

Or

$$v(x) = c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x}$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= \left(c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x} \right) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = e^{\frac{bx^2}{2}}$$

Hence (7) becomes

$$y = \left(c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x} \right) e^{\frac{bx^2}{2}}$$

Therefore the homogeneous solution y_h is

$$y_h = \left(c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x} \right) e^{\frac{bx^2}{2}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}}$$

$$y_2 = e^{\sqrt{-b}x} e^{\frac{bx^2}{2}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}} & e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} \\ \frac{d}{dx} \left(e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}} \right) & \frac{d}{dx} \left(e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}} & e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} \\ -e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}} \sqrt{-b} + e^{-\sqrt{-b}x} x b e^{\frac{bx^2}{2}} & \sqrt{-b} e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} + e^{\sqrt{-b}x} x b e^{\frac{bx^2}{2}} \end{vmatrix}$$

Therefore

$$W = \left(e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}} \right) \left(\sqrt{-b} e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} + e^{\sqrt{-b}x} x b e^{\frac{bx^2}{2}} \right) \\ - \left(e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} \right) \left(-e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}} \sqrt{-b} + e^{-\sqrt{-b}x} x b e^{\frac{bx^2}{2}} \right)$$

Which simplifies to

$$W = 2 e^{bx^2} \sqrt{-b}$$

Which simplifies to

$$W = 2 e^{bx^2} \sqrt{-b}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} x}{2 e^{bx^2} \sqrt{-b}} dx$$

Which simplifies to

$$u_1 = - \int \frac{x e^{-\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} dx$$

Hence

$$u_1 = - \frac{-\frac{e^{-\frac{bx^2}{2} + \sqrt{-b}x}}{b} + \frac{\sqrt{-b} \sqrt{\pi} e^{-\frac{1}{2}\sqrt{2}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{b}x - \sqrt{-b}\sqrt{2}}{2\sqrt{b}}\right)}{2b^{3/2}}}{2\sqrt{-b}}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}} x}{2 e^{bx^2} \sqrt{-b}} dx$$

Which simplifies to

$$u_2 = \int \frac{x e^{-\frac{x(xb+2\sqrt{-b})}{2}}}{2\sqrt{-b}} dx$$

Hence

$$u_2 = \frac{-\frac{e^{-\frac{bx^2}{2} - \sqrt{-b}x}}{b} - \frac{\sqrt{-b} \sqrt{\pi} e^{-\frac{1}{2}\sqrt{2}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{b}x + \sqrt{-b}\sqrt{2}}{2\sqrt{b}}\right)}{2b^{3/2}}}{2\sqrt{-b}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{\left(-\frac{e^{-\frac{bx^2}{2} + \sqrt{-b}x}}{b} + \frac{\sqrt{-b} \sqrt{\pi} e^{-\frac{1}{2}\sqrt{2}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{b}x - \sqrt{-b}\sqrt{2}}{2\sqrt{b}}\right)}{2b^{3/2}} \right) e^{-\sqrt{-b}x} e^{\frac{bx^2}{2}}}{2\sqrt{-b}} \\ + \frac{e^{\sqrt{-b}x} e^{\frac{bx^2}{2}} \left(-\frac{e^{-\frac{bx^2}{2} - \sqrt{-b}x}}{b} - \frac{\sqrt{-b} \sqrt{\pi} e^{-\frac{1}{2}\sqrt{2}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{b}x + \sqrt{-b}\sqrt{2}}{2\sqrt{b}}\right)}{2b^{3/2}} \right)}{2\sqrt{-b}}$$

Which simplifies to

$$y_p(x) = - \frac{e^{\frac{x(xb-2\sqrt{-b})}{2}} \sqrt{\pi} \sqrt{2} \left(e^{-\frac{1}{2}+2\sqrt{-b}x} \operatorname{erf}\left(\frac{\sqrt{2}(xb+\sqrt{-b})}{2\sqrt{b}}\right) - e^{-\frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb+\sqrt{-b})}{2\sqrt{b}}\right) \right)}{4b^{3/2}}$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left((c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x}) e^{\frac{bx^2}{2}} \right) \\
 &\quad + \left(-\frac{e^{\frac{x(xb-2\sqrt{-b})}{2}} \sqrt{\pi} \sqrt{2} \left(e^{-\frac{1}{2}+2\sqrt{-b}x} \operatorname{erf} \left(\frac{\sqrt{2}(xb+\sqrt{-b})}{2\sqrt{b}} \right) - e^{-\frac{1}{2}} \operatorname{erf} \left(\frac{\sqrt{2}(-xb+\sqrt{-b})}{2\sqrt{b}} \right) \right)}{4b^{3/2}} \right)
 \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$\begin{aligned}
 y &= \left(c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x} \right) e^{\frac{bx^2}{2}} \\
 &\quad - \frac{e^{\frac{x(xb-2\sqrt{-b})}{2}} \sqrt{\pi} \sqrt{2} \left(e^{-\frac{1}{2}+2\sqrt{-b}x} \operatorname{erf} \left(\frac{\sqrt{2}(xb+\sqrt{-b})}{2\sqrt{b}} \right) - e^{-\frac{1}{2}} \operatorname{erf} \left(\frac{\sqrt{2}(-xb+\sqrt{-b})}{2\sqrt{b}} \right) \right)}{4b^{3/2}}
 \end{aligned}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.236 (sec)

Writing the ode as

$$y'' - 2bxy' + b^2x^2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}
 A &= 1 \\
 B &= -2xb \\
 C &= x^2b^2
 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}
 r &= \frac{s}{t} \\
 &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}
 \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-b}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned}
 s &= -b \\
 t &= 1
 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = (-b)z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.85: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -b$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\sqrt{-b}x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2xb}{1} dx} \\ &= z_1 e^{\frac{bx^2}{2}} \\ &= z_1 \left(e^{\frac{bx^2}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{\frac{x(xb+2\sqrt{-b})}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2xb}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{bx^2}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{e^{bx^2} e^{-x(xb+2\sqrt{-b})}}{2\sqrt{-b}} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{\frac{x(xb+2\sqrt{-b})}{2}} \right) + c_2 \left(e^{\frac{x(xb+2\sqrt{-b})}{2}} \left(-\frac{e^{bx^2} e^{-x(xb+2\sqrt{-b})}}{2\sqrt{-b}} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2bx y' + b^2 x^2 y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{\frac{x(xb+2\sqrt{-b})}{2}} - \frac{c_2 e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{\frac{x(xb+2\sqrt{-b})}{2}} \\ y_2 &= -\frac{e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} dx \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} dx \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{\frac{x(xb+2\sqrt{-b})}{2}} & -\frac{e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \\ \frac{d}{dx} \left(e^{\frac{x(xb+2\sqrt{-b})}{2}} \right) & \frac{d}{dx} \left(-\frac{e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{\frac{x(xb+2\sqrt{-b})}{2}} & -\frac{e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \\ (xb + \sqrt{-b}) e^{\frac{x(xb+2\sqrt{-b})}{2}} & -\frac{(xb - \sqrt{-b}) e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= \left(e^{\frac{x(xb+2\sqrt{-b})}{2}} \right) \left(-\frac{(xb - \sqrt{-b}) e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \right) \\ &\quad - \left(-\frac{e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \right) \left((xb + \sqrt{-b}) e^{\frac{x(xb+2\sqrt{-b})}{2}} \right) \end{aligned}$$

Which simplifies to

$$W = e^{\frac{x(xb+2\sqrt{-b})}{2}} e^{-\frac{x(-xb+2\sqrt{-b})}{2}}$$

Which simplifies to

$$W = e^{bx^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-\frac{e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} x}{e^{bx^2}} dx$$

Which simplifies to

$$u_1 = - \int -\frac{x e^{-\frac{x(xb+2\sqrt{-b})}{2}}}{2\sqrt{-b}} dx$$

Hence

$$u_1 = \frac{e^{-\frac{bx^2}{2} - \sqrt{-b}x}}{b} - \frac{\sqrt{-b} \sqrt{\pi} e^{-\frac{1}{2}} \sqrt{2} \operatorname{erf} \left(\frac{\sqrt{2} \sqrt{b} x + \sqrt{-b} \sqrt{2}}{2\sqrt{b}} \right)}{2b^{3/2}}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{\frac{x(xb+2\sqrt{-b})}{2}} x}{e^{bx^2}} dx$$

Which simplifies to

$$u_2 = \int x e^{-\frac{x(xb-2\sqrt{-b})}{2}} dx$$

Hence

$$u_2 = -\frac{e^{-\frac{bx^2}{2} + \sqrt{-b}x}}{b} + \frac{\sqrt{-b} \sqrt{\pi} e^{-\frac{1}{2}} \sqrt{2} \operatorname{erf} \left(\frac{\sqrt{2} \sqrt{b} x - \sqrt{-b} \sqrt{2}}{2\sqrt{b}} \right)}{2b^{3/2}}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\left(-\frac{e^{-\frac{bx^2}{2} - \sqrt{-b}x}}{b} - \frac{\sqrt{-b}\sqrt{\pi}e^{-\frac{1}{2}\sqrt{2}\operatorname{erf}\left(\frac{\sqrt{2}\sqrt{b}x + \frac{\sqrt{-b}\sqrt{2}}{2\sqrt{b}}\right)}}{2b^{3/2}} \right) e^{\frac{x(xb+2\sqrt{-b})}{2}}}{2\sqrt{-b}} - \frac{e^{\frac{x(xb-2\sqrt{-b})}{2}} \left(-\frac{e^{-\frac{bx^2}{2} + \sqrt{-b}x}}{b} + \frac{\sqrt{-b}\sqrt{\pi}e^{-\frac{1}{2}\sqrt{2}\operatorname{erf}\left(\frac{\sqrt{2}\sqrt{b}x - \frac{\sqrt{-b}\sqrt{2}}{2\sqrt{b}}\right)}}{2b^{3/2}} \right)}{2\sqrt{-b}}$$

Which simplifies to

$$y_p(x) = \frac{\sqrt{\pi}\sqrt{2}\left(e^{\frac{bx^2}{2} - \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) - e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right)\right)}{4b^{3/2}}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 e^{\frac{x(xb+2\sqrt{-b})}{2}} - \frac{c_2 e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} \right) + \left(\frac{\sqrt{\pi}\sqrt{2}\left(e^{\frac{bx^2}{2} - \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) - e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right)\right)}{4b^{3/2}} \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{\frac{x(xb+2\sqrt{-b})}{2}} - \frac{c_2 e^{\frac{x(xb-2\sqrt{-b})}{2}}}{2\sqrt{-b}} + \frac{\sqrt{\pi}\sqrt{2}\left(e^{\frac{bx^2}{2} - \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) - e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right)\right)}{4b^{3/2}}$$

Solved as second order ode adjoint method

Time used: 0.562 (sec)

In normal form the ode

$$y'' - 2bxy' + b^2x^2y = x \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -2xb \\ q(x) &= x^2b^2 \\ r(x) &= x \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (-2xb\xi(x))' + (x^2b^2\xi(x)) &= 0 \\ \xi''(x) + 2xb\xi'(x) + (x^2b^2 + 2b)\xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x)\xi' + q(x)\xi = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= 2xb \\ q(x) &= x^2b^2 + 2b \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= x^2b^2 + 2b - \frac{(2xb)'}{2} - \frac{(2xb)^2}{4} \\ &= x^2b^2 + 2b - \frac{(2b)}{2} - \frac{(4x^2b^2)}{4} \\ &= x^2b^2 + 2b - (b) - x^2b^2 \\ &= b \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{2xb}{2} dx} \\ &= e^{-\frac{bx^2}{2}} \end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = v(x)e^{-\frac{bx^2}{2}} \quad (4)$$

Applying this change of variable to the original ode results in

$$e^{-\frac{bx^2}{2}}(bv(x) + v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$bv(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = b$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + b e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + b = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = b$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(b)} \\ &= \pm \sqrt{-b}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +\sqrt{-b} \\ \lambda_2 &= -\sqrt{-b}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \sqrt{-b} \\ \lambda_2 &= -\sqrt{-b}\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}v(x) &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ v(x) &= c_1 e^{(\sqrt{-b})x} + c_2 e^{(-\sqrt{-b})x}\end{aligned}$$

Or

$$v(x) = c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x}$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned}\xi &= v(x) z(x) \\ &= \left(c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x} \right) (z(x))\end{aligned}\tag{7}$$

But from (5)

$$z(x) = e^{-\frac{bx^2}{2}}$$

Hence (7) becomes

$$\xi = \left(c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x} \right) e^{-\frac{bx^2}{2}}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(-2xb - \frac{\left(\left(c_1 \sqrt{-b} e^{\sqrt{-b}x} - c_2 \sqrt{-b} e^{-\sqrt{-b}x} \right) e^{-\frac{bx^2}{2}} - \left(c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x} \right) xb e^{-\frac{bx^2}{2}} \right) e^{\frac{bx^2}{2}}}{c_1 e^{\sqrt{-b}x} + c_2 e^{-\sqrt{-b}x}} \right) = \frac{e^{\frac{bx^2}{2}}}{\xi(x)}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{c_1(xb + \sqrt{-b})e^{2\sqrt{-b}x} + c_2(xb - \sqrt{-b})}{c_1e^{2\sqrt{-b}x} + c_2}$$

$$p(x) = -\frac{c_1\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) + c_2\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right)}{b^{3/2}(2c_1e^{2\sqrt{-b}x} + 2c_2)}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{c_1(xb + \sqrt{-b})e^{2\sqrt{-b}x} + c_2(xb - \sqrt{-b})}{c_1e^{2\sqrt{-b}x} + c_2} dx}$$

Therefore the solution is

$$y = \left(\int -\frac{(c_1\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) + c_2\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right))}{b^{3/2}(2c_1e^{2\sqrt{-b}x} + 2c_2)} dx \right)$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \left(\int -\frac{(c_1\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) + c_2\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right) + 2c_3}{b^{3/2}(2c_1e^{2\sqrt{-b}x} + 2c_2)} dx \right. \\ \left. + c_3 \right) e^{-\int -\frac{c_1(xb + \sqrt{-b})e^{2\sqrt{-b}x} + c_2(xb - \sqrt{-b})}{c_1e^{2\sqrt{-b}x} + c_2} dx}$$

The constants can be merged to give

$$y = \left(\int -\frac{(c_1\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) + c_2\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right) + 2c_3}{b^{3/2}(2c_1e^{2\sqrt{-b}x} + 2c_2)} dx \right. \\ \left. + 1 \right) e^{-\int -\frac{c_1(xb + \sqrt{-b})e^{2\sqrt{-b}x} + c_2(xb - \sqrt{-b})}{c_1e^{2\sqrt{-b}x} + c_2} dx}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(\int -\frac{(c_1\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(-xb + \sqrt{-b})}{2\sqrt{b}}\right) + c_2\sqrt{2}\sqrt{\pi}\sqrt{-b}e^{\frac{bx^2}{2} + \sqrt{-b}x - \frac{1}{2}} \operatorname{erf}\left(\frac{\sqrt{2}(xb + \sqrt{-b})}{2\sqrt{b}}\right) + 2c_3}{b^{3/2}(2c_1e^{2\sqrt{-b}x} + 2c_2)} dx \right. \\ \left. + 1 \right) e^{-\int -\frac{c_1(xb + \sqrt{-b})e^{2\sqrt{-b}x} + c_2(xb - \sqrt{-b})}{c_1e^{2\sqrt{-b}x} + c_2} dx}$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
    <- Kovacic's algorithm successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.016 (sec)

Leaf size : 137

```

dsolve(diff(diff(y(x),x),x)-2*b*x*diff(y(x),x)+b^2*x^2*y(x) = x,
        y(x),singsol=all)

```

$$y = \frac{4e^{\frac{x(bx-2\sqrt{-b})}{2}}c_1b^{3/2} - \operatorname{erf}\left(\frac{\sqrt{2}(bx+\sqrt{-b})}{2\sqrt{b}}\right)\sqrt{\pi}\sqrt{2}e^{\frac{bx^2}{2}+x\sqrt{-b}-\frac{1}{2}} + 4e^{\frac{x(bx+2\sqrt{-b})}{2}}c_2b^{3/2} + \sqrt{\pi}e^{\frac{bx^2}{2}-x\sqrt{-b}-\frac{1}{2}}\sqrt{2}\operatorname{erf}\left(\frac{\sqrt{2}(bx+\sqrt{-b})}{2\sqrt{b}}\right)}{4b^{3/2}}$$

Mathematica DSolve solution

Solving time : 0.783 (sec)

Leaf size : 139

```

DSolve[{D[y[x],{x,2}]-2*b*x*D[y[x],x]+b^2*x^2*y[x]==x,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{e^{\frac{1}{2}(\sqrt{bx}-i)^2}\left(-\sqrt{2\pi}e^{2i\sqrt{bx}}\operatorname{erf}\left(\frac{\sqrt{bx}+i}{\sqrt{2}}\right) + i\sqrt{2\pi}\operatorname{erfi}\left(\frac{1+i\sqrt{bx}}{\sqrt{2}}\right) + 2\sqrt{eb}\left(2\sqrt{b}c_1 - ic_2e^{2i\sqrt{bx}}\right)\right)}{4b^{3/2}}$$

2.2.32 problem 32

Solved as second order ode using change of variable on y	
method 1	771
Solved as second order ode using Kovacic algorithm	775
Solved as second order ode adjoint method	779
Maple step by step solution	782
Maple trace	782
Maple dsolve solution	782
Mathematica DSolve solution	782

Internal problem ID [8831]

Book : Second order enumerated odes

Section : section 2

Problem number : 32

Date solved : Thursday, December 12, 2024 at 09:53:13 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' - 4xy' + (4x^2 - 3)y = e^{x^2}$$

Solved as second order ode using change of variable on y method 1

Time used: 0.472 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4xy' + (4x^2 - 3)y = 0$$

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$\begin{aligned} p(x) &= -4x \\ q(x) &= 4x^2 - 3 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 4x^2 - 3 - \frac{(-4x)'}{2} - \frac{(-4x)^2}{4} \\ &= 4x^2 - 3 - \frac{(-4)}{2} - \frac{(16x^2)}{4} \\ &= 4x^2 - 3 - (-2) - 4x^2 \\ &= -1 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{-4x}{2}} \\ &= e^{x^2} \end{aligned} \tag{5}$$

Hence (3) becomes

$$y = v(x) e^{x^2} \tag{4}$$

Applying this change of variable to the original ode results in

$$e^{x^2} (v''(x) - v(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$v''(x) - v(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - e^{x\lambda} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$v(x) = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$v(x) = c_1 e^{(1)x} + c_2 e^{(-1)x}$$

Or

$$v(x) = c_1 e^x + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 e^x + c_2 e^{-x}) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = e^{x^2}$$

Hence (7) becomes

$$y = (c_1 e^x + c_2 e^{-x}) e^{x^2}$$

Therefore the homogeneous solution y_h is

$$y_h = (c_1 e^x + c_2 e^{-x}) e^{x^2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{-x} e^{x^2} \\ y_2 &= e^x e^{x^2} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-x} e^{x^2} & e^x e^{x^2} \\ \frac{d}{dx} (e^{-x} e^{x^2}) & \frac{d}{dx} (e^x e^{x^2}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-x} e^{x^2} & e^x e^{x^2} \\ -e^{-x} e^{x^2} + 2e^{-x} x e^{x^2} & e^x e^{x^2} + 2e^x x e^{x^2} \end{vmatrix}$$

Therefore

$$W = (e^{-x} e^{x^2}) (e^x e^{x^2} + 2e^x x e^{x^2}) - (e^x e^{x^2}) (-e^{-x} e^{x^2} + 2e^{-x} x e^{x^2})$$

Which simplifies to

$$W = 2e^{2x^2}$$

Which simplifies to

$$W = 2e^{2x^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^x e^{2x^2}}{2e^{2x^2}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^x}{2} dx$$

Hence

$$u_1 = -\frac{e^x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-x} e^{2x^2}}{2e^{2x^2}} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-x}}{2} dx$$

Hence

$$u_2 = -\frac{e^{-x}}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{e^{x^2}}{2} - \frac{e^x e^{x^2} e^{-x}}{2}$$

Which simplifies to

$$y_p(x) = -e^{x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left((c_1 e^x + c_2 e^{-x}) e^{x^2} \right) + \left(-e^{x^2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = (c_1 e^x + c_2 e^{-x}) e^{x^2} - e^{x^2}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.208 (sec)

Writing the ode as

$$y'' - 4xy' + (4x^2 - 3)y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -4x \\ C &= 4x^2 - 3 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.86: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-4x}{1} dx} \\ &= z_1 e^{x^2} \\ &= z_1 (e^{x^2}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{x(x-1)}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-4x}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x^2}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{2x^2} e^{-2x(x-1)}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{x(x-1)}) + c_2 \left(e^{x(x-1)} \left(\frac{e^{2x^2} e^{-2x(x-1)}}{2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4xy' + (4x^2 - 3)y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{x(x-1)} + \frac{c_2 e^{x(x+1)}}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{x(x-1)}$$

$$y_2 = \frac{e^{x(x+1)}}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{x(x-1)} & \frac{e^{x(x+1)}}{2} \\ \frac{d}{dx}(e^{x(x-1)}) & \frac{d}{dx}\left(\frac{e^{x(x+1)}}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{x(x-1)} & \frac{e^{x(x+1)}}{2} \\ (2x-1)e^{x(x-1)} & \frac{(2x+1)e^{x(x+1)}}{2} \end{vmatrix}$$

Therefore

$$W = (e^{x(x-1)}) \left(\frac{(2x+1)e^{x(x+1)}}{2} \right) - \left(\frac{e^{x(x+1)}}{2} \right) ((2x-1)e^{x(x-1)})$$

Which simplifies to

$$W = e^{x(x-1)} e^{x(x+1)}$$

Which simplifies to

$$W = e^{2x^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{x(x+1)} e^{x^2}}{2 e^{2x^2}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^x}{2} dx$$

Hence

$$u_1 = -\frac{e^x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{x(x-1)} e^{x^2}}{e^{2x^2}} dx$$

Which simplifies to

$$u_2 = \int e^{-x} dx$$

Hence

$$u_2 = -e^{-x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{e^x e^{x(x-1)}}{2} - \frac{e^{x(x+1)} e^{-x}}{2}$$

Which simplifies to

$$y_p(x) = -e^{x^2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{x(x-1)} + \frac{c_2 e^{x(x+1)}}{2} \right) + \left(-e^{x^2} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{x(x-1)} + \frac{c_2 e^{x(x+1)}}{2} - e^{x^2}$$

Solved as second order ode adjoint method

Time used: 0.423 (sec)

In normal form the ode

$$y'' - 4xy' + (4x^2 - 3)y = e^{x^2} \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -4x \\ q(x) &= 4x^2 - 3 \\ r(x) &= e^{x^2} \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (-4x\xi(x))' + ((4x^2 - 3)\xi(x)) &= 0 \\ 4\xi(x)x^2 + 4x\xi'(x) + \xi''(x) + \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x)\xi' + q(x)\xi = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= 4x \\ q(x) &= 4x^2 + 1 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 4x^2 + 1 - \frac{(4x)'}{2} - \frac{(4x)^2}{4} \\ &= 4x^2 + 1 - \frac{(4)}{2} - \frac{(16x^2)}{4} \\ &= 4x^2 + 1 - (2) - 4x^2 \\ &= -1 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{4x}{2} dx} \\ &= e^{-x^2} \end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = v(x)e^{-x^2} \quad (4)$$

Applying this change of variable to the original ode results in

$$e^{-x^2}(-v(x) + v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$-v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +1 \\ \lambda_2 &= -1 \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 1 \\ \lambda_2 &= -1 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} v(x) &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ v(x) &= c_1 e^{(1)x} + c_2 e^{(-1)x} \end{aligned}$$

Or

$$v(x) = c_1 e^x + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} \xi &= v(x) z(x) \\ &= (c_1 e^x + c_2 e^{-x}) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = e^{-x^2}$$

Hence (7) becomes

$$\xi = (c_1 e^x + c_2 e^{-x}) e^{-x^2}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(-4x - \frac{((c_1 e^x - c_2 e^{-x}) e^{-x^2} - 2(c_1 e^x + c_2 e^{-x}) x e^{-x^2}) e^{x^2}}{c_1 e^x + c_2 e^{-x}} \right) = \frac{e^{x^2} (c_1 e^x - c_2 e^{-x})}{c_1 e^x + c_2 e^{-x}}$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} \\ p(x) &= \frac{e^{x^2}(c_1 e^{2x} - c_2)}{c_1 e^{2x} + c_2} \end{aligned}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}$$

Therefore the solution is

$$y = \left(\int \frac{e^{x^2}(c_1 e^{2x} - c_2) e^{\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}}{c_1 e^{2x} + c_2} dx + c_3 \right) e^{-\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \left(\int \frac{e^{x^2}(c_1 e^{2x} - c_2) e^{\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}}{c_1 e^{2x} + c_2} dx + c_3 \right) e^{-\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}$$

The constants can be merged to give

$$y = \left(\int \frac{e^{x^2}(c_1 e^{2x} - c_2) e^{\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}}{c_1 e^{2x} + c_2} dx + 1 \right) e^{-\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(\int \frac{e^{x^2}(c_1 e^{2x} - c_2) e^{\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}}{c_1 e^{2x} + c_2} dx + 1 \right) e^{-\int -\frac{c_1(2x+1)e^{2x} + c_2(2x-1)}{c_1 e^{2x} + c_2} dx}$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacic's algorithm successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.009 (sec)
 Leaf size : 27

```

dsolve(diff(diff(y(x),x),x)-4*diff(y(x),x)*x+(4*x^2-3)*y(x) = exp(x^2),
        y(x),singsol=all)

```

$$y = e^{x(x+1)}c_2 + e^{x(x-1)}c_1 - e^{x^2}$$

Mathematica DSolve solution

Solving time : 0.06 (sec)
 Leaf size : 34

```

DSolve[{D[y[x],{x,2}]-4*x*D[y[x],x]+(4*x^2-3)*y[x]==Exp[x^2],{}}],
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{1}{2}e^{(x-1)x}(-2e^x + c_2e^{2x} + 2c_1)$$

2.2.33 problem 33

Solved as second order ode using change of variable on y
 method 1 783
 Solved as second order ode using Kovacic algorithm 787
 Solved as second order ode adjoint method 791
 Maple step by step solution 796
 Maple trace 796
 Maple dsolve solution 796
 Mathematica DSolve solution 796

Internal problem ID [8832]

Book : Second order enumerated odes

Section : section 2

Problem number : 33

Date solved : Thursday, December 12, 2024 at 09:53:15 AM

CAS classification : [[_2nd_order, _linear, _nonhomogeneous]]

Solve

$$y'' - 2 \tan(x) y' + 5y = e^{x^2} \sec(x)$$

Solved as second order ode using change of variable on y method 1

Time used: 0.809 (sec)

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2 \tan(x) y' + 5y = 0$$

In normal form the given ode is written as

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$p(x) = -2 \tan(x)$$

$$q(x) = 5$$

Calculating the Liouville ode invariant Q given by

$$Q = q - \frac{p'}{2} - \frac{p^2}{4}$$

$$= 5 - \frac{(-2 \tan(x))'}{2} - \frac{(-2 \tan(x))^2}{4}$$

$$= 5 - \frac{(-2 - 2 \tan(x)^2)}{2} - \frac{(4 \tan(x)^2)}{4}$$

$$= 5 - (-1 - \tan(x)^2) - \tan(x)^2$$

$$= 6$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x) z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{-2 \tan(x)}{2} dx} \\ &= \sec(x) \end{aligned} \quad (5)$$

Hence (3) becomes

$$y = v(x) \sec(x) \quad (4)$$

Applying this change of variable to the original ode results in

$$\sec(x) (v''(x) + 6v(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$v''(x) + 6v(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 6$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + 6 e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(6)} \\ &= \pm i\sqrt{6} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i\sqrt{6} \\ \lambda_2 &= -i\sqrt{6} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i\sqrt{6} \\ \lambda_2 &= -i\sqrt{6} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{6}$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 \left(c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x) \right)$$

Or

$$v(x) = c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= \left(c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x) \right) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \sec(x)$$

Hence (7) becomes

$$y = \left(c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x) \right) \sec(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \left(c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x) \right) \sec(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(\sqrt{6}x) \sec(x)$$

$$y_2 = \sin(\sqrt{6}x) \sec(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(\sqrt{6}x) \sec(x) & \sin(\sqrt{6}x) \sec(x) \\ \frac{d}{dx}(\cos(\sqrt{6}x) \sec(x)) & \frac{d}{dx}(\sin(\sqrt{6}x) \sec(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(\sqrt{6}x) \sec(x) & \sin(\sqrt{6}x) \sec(x) \\ -\sqrt{6} \sin(\sqrt{6}x) \sec(x) + \cos(\sqrt{6}x) \sec(x) \tan(x) & \sqrt{6} \cos(\sqrt{6}x) \sec(x) + \sin(\sqrt{6}x) \sec(x) \tan(x) \end{vmatrix}$$

Therefore

$$W = \left(\cos(\sqrt{6}x) \sec(x) \right) \left(\sqrt{6} \cos(\sqrt{6}x) \sec(x) + \sin(\sqrt{6}x) \sec(x) \tan(x) \right) - \left(\sin(\sqrt{6}x) \sec(x) \right) \left(-\sqrt{6} \sin(\sqrt{6}x) \sec(x) + \cos(\sqrt{6}x) \sec(x) \tan(x) \right)$$

Which simplifies to

$$W = \sec(x)^2 \sqrt{6} \cos(\sqrt{6}x)^2 + \sec(x)^2 \sqrt{6} \sin(\sqrt{6}x)^2$$

Which simplifies to

$$W = \sqrt{6} \sec(x)^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(\sqrt{6}x) \sec(x)^2 e^{x^2}}{\sqrt{6} \sec(x)^2} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin(\sqrt{6}x) e^{x^2} \sqrt{6}}{6} dx$$

Hence

$$u_1 = \frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{24} - \frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{24}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(\sqrt{6}x) \sec(x)^2 e^{x^2}}{\sqrt{6} \sec(x)^2} dx$$

Which simplifies to

$$u_2 = \int \frac{\cos(\sqrt{6}x) e^{x^2} \sqrt{6}}{6} dx$$

Hence

$$u_2 = - \frac{i\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{24} - \frac{i\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{24}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{24} - \frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{24} \right) \cos(\sqrt{6}x) \sec(x) + \sin(\sqrt{6}x) \sec(x) \left(- \frac{i\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{24} - \frac{i\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{24} \right)$$

Which simplifies to

$$y_p(x) = \frac{\left((i \sin(\sqrt{6}x) - \cos(\sqrt{6}x)) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (i \sin(\sqrt{6}x) + \cos(\sqrt{6}x)) \right) \sqrt{\pi} e^{\frac{3}{2}\sqrt{6}x}}{24}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left((c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x)) \sec(x) \right) \\ &\quad + \left(-\frac{\left((i \sin(\sqrt{6}x) - \cos(\sqrt{6}x)) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (i \sin(\sqrt{6}x) + \cos(\sqrt{6}x)) \right) \sqrt{\pi} e^{\frac{3}{2}\sqrt{6}x}}{24} \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x) \right) \sec(x) - \frac{\left((i \sin(\sqrt{6}x) - \cos(\sqrt{6}x)) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (i \sin(\sqrt{6}x) + \cos(\sqrt{6}x)) \right) \sqrt{\pi} e^{\frac{3}{2}\sqrt{6}x}}{24}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.377 (sec)

Writing the ode as

$$y'' - 2 \tan(x) y' + 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \tan(x) \\ C &= 5 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-6}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -6 \\ t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -6z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.87: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -6$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(\sqrt{6}x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2 \tan(x)}{1} dx} \\ &= z_1 e^{-\ln(\cos(x))} \\ &= z_1 (\sec(x))\end{aligned}$$

Which simplifies to

$$y_1 = \cos(\sqrt{6}x) \sec(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2 \tan(x)}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2 \ln(\cos(x))}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{6} \tan(\sqrt{6}x)}{6} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\cos(\sqrt{6}x) \sec(x) \right) + c_2 \left(\cos(\sqrt{6}x) \sec(x) \left(\frac{\sqrt{6} \tan(\sqrt{6}x)}{6} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2 \tan(x) y' + 5y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(\sqrt{6}x) \sec(x) + \frac{c_2 \sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \cos(\sqrt{6}x) \sec(x) \\ y_2 &= \frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(\sqrt{6}x) \sec(x) & \frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6} \\ \frac{d}{dx}(\cos(\sqrt{6}x) \sec(x)) & \frac{d}{dx}\left(\frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(\sqrt{6}x) \sec(x) & \frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6} \\ -\sqrt{6} \sin(\sqrt{6}x) \sec(x) + \cos(\sqrt{6}x) \sec(x) \tan(x) & \cos(\sqrt{6}x) \sec(x) + \frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x) \tan(x)}{6} \end{vmatrix}$$

Therefore

$$W = \left(\cos(\sqrt{6}x) \sec(x) \right) \left(\cos(\sqrt{6}x) \sec(x) + \frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x) \tan(x)}{6} \right) - \left(\frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6} \right) \left(-\sqrt{6} \sin(\sqrt{6}x) \sec(x) + \cos(\sqrt{6}x) \sec(x) \tan(x) \right)$$

Which simplifies to

$$W = \cos(\sqrt{6}x)^2 \sec(x)^2 + \sec(x)^2 \sin(\sqrt{6}x)^2$$

Which simplifies to

$$W = \sec(x)^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x)^2 e^{x^2}}{6}}{\sec(x)^2} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin(\sqrt{6}x) e^{x^2} \sqrt{6}}{6} dx$$

Hence

$$u_1 = \frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{24} - \frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{24}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(\sqrt{6}x) \sec(x)^2 e^{x^2}}{\sec(x)^2} dx$$

Which simplifies to

$$u_2 = \int \cos(\sqrt{6}x) e^{x^2} dx$$

Hence

$$u_2 = -\frac{i\sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{4} - \frac{i\sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{24} - \frac{\sqrt{6} \sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{24} \right) \cos(\sqrt{6}x) \sec(x) \\ + \frac{\sqrt{6} \sin(\sqrt{6}x) \sec(x) \left(-\frac{i\sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)}{4} - \frac{i\sqrt{\pi} e^{\frac{3}{2}} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)}{4} \right)}{6}$$

Which simplifies to

$$y_p(x) = \frac{\left((i \sin(\sqrt{6}x) - \cos(\sqrt{6}x)) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (i \sin(\sqrt{6}x) + \cos(\sqrt{6}x)) \right) \sqrt{\pi} e^{\frac{3}{2}} \sqrt{6} \sec(x)}{24}$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1 \cos(\sqrt{6}x) \sec(x) + \frac{c_2 \sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6} \right) \\ + \frac{\left((i \sin(\sqrt{6}x) - \cos(\sqrt{6}x)) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (i \sin(\sqrt{6}x) + \cos(\sqrt{6}x)) \right) \sqrt{\pi} e^{\frac{3}{2}} \sqrt{6} \sec(x)}{24}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos(\sqrt{6}x) \sec(x) + \frac{c_2 \sqrt{6} \sin(\sqrt{6}x) \sec(x)}{6} \\ + \frac{\left((i \sin(\sqrt{6}x) - \cos(\sqrt{6}x)) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (i \sin(\sqrt{6}x) + \cos(\sqrt{6}x)) \right) \sqrt{\pi} e^{\frac{3}{2}} \sqrt{6} \sec(x)}{24}$$

Solved as second order ode adjoint method

Time used: 13.993 (sec)

In normal form the ode

$$y'' - 2 \tan(x) y' + 5y = e^{x^2} \sec(x) \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -2 \tan(x) \\ q(x) &= 5 \\ r(x) &= e^{x^2} \sec(x) \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (-2 \tan(x) \xi(x))' + (5\xi(x)) &= 0 \\ 2 \tan(x) \xi'(x) + 5\xi(x) + \xi''(x) + 2 \sec(x)^2 \xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x) \xi' + q(x) \xi = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= 2 \tan(x) \\ q(x) &= 5 + 2 \sec(x)^2 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 5 + 2 \sec(x)^2 - \frac{(2 \tan(x))'}{2} - \frac{(2 \tan(x))^2}{4} \\ &= 5 + 2 \sec(x)^2 - \frac{(2 + 2 \tan(x)^2)}{2} - \frac{(4 \tan(x)^2)}{4} \\ &= 5 + 2 \sec(x)^2 - (1 + \tan(x)^2) - \tan(x)^2 \\ &= 6 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x) z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{2 \tan(x)}{2} dx} \\ &= \cos(x) \end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = v(x) \cos(x) \quad (4)$$

Applying this change of variable to the original ode results in

$$(6v(x) + v''(x)) \cos(x) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$6v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 6$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + 6 e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(6)} \\ &= \pm i\sqrt{6} \end{aligned}$$

Hence

$$\lambda_1 = +i\sqrt{6}$$

$$\lambda_2 = -i\sqrt{6}$$

Which simplifies to

$$\lambda_1 = i\sqrt{6}$$

$$\lambda_2 = -i\sqrt{6}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{6}$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (c_1 \cos(\sqrt{6} x) + c_2 \sin(\sqrt{6} x))$$

Or

$$v(x) = c_1 \cos(\sqrt{6} x) + c_2 \sin(\sqrt{6} x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} \xi &= v(x) z(x) \\ &= (c_1 \cos(\sqrt{6} x) + c_2 \sin(\sqrt{6} x)) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \cos(x)$$

Hence (7) becomes

$$\xi = (c_1 \cos(\sqrt{6} x) + c_2 \sin(\sqrt{6} x)) \cos(x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(-2 \tan(x) - \frac{(-c_1 \sqrt{6} \sin(\sqrt{6}x) + c_2 \sqrt{6} \cos(\sqrt{6}x)) \cos(x) - (c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x)) \sin(x)}{(c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x)) \cos(x)} \right)$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{(-\sqrt{6}c_1 + \tan(x)c_2) \sin(\sqrt{6}x) + (\tan(x)c_1 + c_2\sqrt{6}) \cos(\sqrt{6}x)}{c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x)} \\ p(x) &= -\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (ic_1 - c_2) \right) e^{\frac{3}{2} \sec(x)} \sqrt{\pi}}{4c_1 \cos(\sqrt{6}x) + 4c_2 \sin(\sqrt{6}x)}\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{(-\sqrt{6}c_1 + \tan(x)c_2) \sin(\sqrt{6}x) + (\tan(x)c_1 + c_2\sqrt{6}) \cos(\sqrt{6}x)}{c_1 \cos(\sqrt{6}x) + c_2 \sin(\sqrt{6}x)} dx} \\ &= e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2\right)}\end{aligned}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = \mu p$$

$$\frac{d}{dx}(\mu y) = (\mu) \left(-\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (ic_1 - c_2) \right) e^{\frac{3}{2} \sec(x)} \sqrt{\pi}}{4c_1 \cos(\sqrt{6}x) + 4c_2 \sin(\sqrt{6}x)} \right)$$

$$\begin{aligned}\frac{d}{dx} \left(y e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2\right)} \right) \\ = \left(e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2\right)} \right) \left(-\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (ic_1 - c_2) \right) e^{\frac{3}{2} \sec(x)} \sqrt{\pi}}{4c_1 \cos(\sqrt{6}x) + 4c_2 \sin(\sqrt{6}x)} \right)\end{aligned}$$

$$\begin{aligned}d \left(y e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2\right)} \right) \\ = \left(-\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) (ic_1 - c_2) \right) e^{\frac{3}{2} \sec(x)} \sqrt{\pi} e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2\right)}}{4c_1 \cos(\sqrt{6}x) + 4c_2 \sin(\sqrt{6}x)} \right)\end{aligned}$$

Integrating gives

$$y e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2\right)} = \int -\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)\right)}{4c_1} dx$$

$$= \int -\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right)\right)}{4c_1} dx$$

Dividing throughout by the integrating factor $e^{-\frac{\ln(1+\tan(x)^2)}{2} - \ln\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) + \ln\left(1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2\right)}$ gives the final solution

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2} \left(\int -\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)\right)(ic_1 - c_2) e^{\frac{3}{2} \sec(x)}}{4c_1} dx\right)}{1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2} \left(\int -\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)\right)(ic_1 - c_2) e^{\frac{3}{2} \sec(x)}}{4c_1} dx\right)}{1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2}$$

The constants can be merged to give

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2} \left(\int -\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)\right)(ic_1 - c_2) e^{\frac{3}{2} \sec(x)}}{4c_1} dx\right)}{1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{\left(c_1 \tan\left(\frac{\sqrt{6}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{6}x}{2}\right) - c_1\right) \sqrt{1 + \tan(x)^2} \left(\int -\frac{\left((ic_1 + c_2) \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right)\right)(ic_1 - c_2) e^{\frac{3}{2} \sec(x)}}{4c_1} dx\right)}{1 + \tan\left(\frac{\sqrt{6}x}{2}\right)^2}$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
trying symmetries linear in x and y(x)
-> Try solving first the homogeneous part of the ODE
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Trying a Liouvillian solution using Kovacic's algorithm
        A Liouvillian solution exists
        Group is reducible or imprimitive
    <- Kovacic's algorithm successful
<- solving first the homogeneous part of the ODE successful`

```

Maple dsolve solution

Solving time : 0.025 (sec)

Leaf size : 97

```

dsolve(diff(diff(y(x),x),x)-2*tan(x)*diff(y(x),x)+5*y(x) = exp(x^2)*sec(x),
        y(x),singsol=all)

```

 $y =$

$$\frac{\left(e^{\frac{3}{2}} (i \sin(\sqrt{6}x) - \cos(\sqrt{6}x)) \sqrt{6} \sqrt{\pi} \operatorname{erf}\left(ix - \frac{\sqrt{6}}{2}\right) + e^{\frac{3}{2}} (i \sin(\sqrt{6}x) + \cos(\sqrt{6}x)) \sqrt{6} \sqrt{\pi} \operatorname{erf}\left(ix + \frac{\sqrt{6}}{2}\right) \right)}{24}$$

Mathematica DSolve solution

Solving time : 0.416 (sec)

Leaf size : 118

```

DSolve[{D[y[x],{x,2}]-2*Tan[x]*D[y[x],x]+5*y[x]==Exp[x^2]*Sec[x],{}}],
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{1}{24} e^{-i\sqrt{6}x} \sec(x) \left(-e^{3/2} \sqrt{6\pi} \operatorname{erf}\left(\sqrt{\frac{3}{2}} - ix\right) - \sqrt{6\pi} e^{\frac{3}{2}+2i\sqrt{6}x} \operatorname{erf}\left(\sqrt{\frac{3}{2}} + ix\right) - 2i\sqrt{6}c_2 e^{2i\sqrt{6}x} + 24c_1 \right)$$

2.2.34 problem 34

Solved as second order ode using change of variable on y
 method 1 797
 Solved as second order Bessel ode 799
 Solved as second order ode using Kovacic algorithm 800
 Solved as second order ode adjoint method 802
 Maple step by step solution 805
 Maple trace 807
 Maple dsolve solution 807
 Mathematica DSolve solution 807

Internal problem ID [8833]

Book : Second order enumerated odes

Section : section 2

Problem number : 34

Date solved : Thursday, December 12, 2024 at 09:53:31 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$x^2y'' - 2xy' + 2(x^2 + 1)y = 0$$

Solved as second order ode using change of variable on y method 1

Time used: 0.366 (sec)

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{2}{x}$$

$$q(x) = \frac{2x^2 + 2}{x^2}$$

Calculating the Liouville ode invariant Q given by

$$Q = q - \frac{p'}{2} - \frac{p^2}{4}$$

$$= \frac{2x^2 + 2}{x^2} - \frac{(-\frac{2}{x})'}{2} - \frac{(-\frac{2}{x})^2}{4}$$

$$= \frac{2x^2 + 2}{x^2} - \frac{(\frac{2}{x^2})}{2} - \frac{(\frac{4}{x^2})}{4}$$

$$= \frac{2x^2 + 2}{x^2} - \left(\frac{1}{x^2}\right) - \frac{1}{x^2}$$

$$= 2$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$z(x) = e^{-\int \frac{p(x)}{2} dx}$$

$$= e^{-\int \frac{-2}{x} dx}$$

$$= x \tag{5}$$

Hence (3) becomes

$$y = v(x)x \quad (4)$$

Applying this change of variable to the original ode results in

$$x^3(2v(x) + v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$2v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 2$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + 2e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(2)} \\ &= \pm i\sqrt{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{2}$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x}(c_3 \cos(\beta x) + c_4 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 \left(c_3 \cos(\sqrt{2}x) + c_4 \sin(\sqrt{2}x) \right)$$

Or

$$v(x) = c_3 \cos(\sqrt{2}x) + c_4 \sin(\sqrt{2}x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= \left(c_3 \cos(\sqrt{2}x) + c_4 \sin(\sqrt{2}x) \right) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = x$$

Hence (7) becomes

$$y = \left(c_3 \cos(\sqrt{2}x) + c_4 \sin(\sqrt{2}x) \right) x$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(c_3 \cos(\sqrt{2}x) + c_4 \sin(\sqrt{2}x) \right) x$$

Solved as second order Bessel ode

Time used: 0.062 (sec)

Writing the ode as

$$x^2 y'' - 2xy' + (2x^2 + 2)y = 0 \quad (1)$$

Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2)y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha)xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2)y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= \frac{3}{2} \\ \beta &= \sqrt{2} \\ n &= -\frac{1}{2} \\ \gamma &= 1 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1 x^{3/2} \sqrt{2} \cos(\sqrt{2}x)}{\sqrt{\pi} \sqrt{\sqrt{2}x}} + \frac{c_2 x^{3/2} \sqrt{2} \sin(\sqrt{2}x)}{\sqrt{\pi} \sqrt{\sqrt{2}x}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 x^{3/2} \sqrt{2} \cos(\sqrt{2}x)}{\sqrt{\pi} \sqrt{\sqrt{2}x}} + \frac{c_2 x^{3/2} \sqrt{2} \sin(\sqrt{2}x)}{\sqrt{\pi} \sqrt{\sqrt{2}x}}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.133 (sec)

Writing the ode as

$$x^2 y'' - 2xy' + (2x^2 + 2)y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= -2x \\ C &= 2x^2 + 2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-2}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -2 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -2z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.88: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(\sqrt{2}x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2x}{x^2} dx} \\ &= z_1 e^{\ln(x)} \\ &= z_1(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(\sqrt{2}x) x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2 \ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{2} \tan(\sqrt{2}x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\cos(\sqrt{2}x) x \right) + c_2 \left(\cos(\sqrt{2}x) x \left(\frac{\sqrt{2} \tan(\sqrt{2}x)}{2} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 \cos(\sqrt{2}x) x + \frac{c_2 \sqrt{2} x \sin(\sqrt{2}x)}{2}$$

Solved as second order ode adjoint method

Time used: 2.292 (sec)

In normal form the ode

$$x^2 y'' - 2xy' + 2(x^2 + 1)y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{2}{x} \\ q(x) &= \frac{2x^2 + 2}{x^2} \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(-\frac{2\xi(x)}{x}\right)' + \left(\frac{(2x^2 + 2)\xi(x)}{x^2}\right) &= 0 \\ \xi''(x) + \frac{2\xi'(x)}{x} + 2\xi(x) &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x)\xi' + q(x)\xi = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{2}{x} \\ q(x) &= 2 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 2 - \frac{\left(\frac{2}{x}\right)'}{2} - \frac{\left(\frac{2}{x}\right)^2}{4} \\ &= 2 - \frac{\left(-\frac{2}{x^2}\right)}{2} - \frac{\left(\frac{4}{x^2}\right)}{4} \\ &= 2 - \left(-\frac{1}{x^2}\right) - \frac{1}{x^2} \\ &= 2 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{2}{x} dx} \\ &= \frac{1}{x} \end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = \frac{v(x)}{x} \quad (4)$$

Applying this change of variable to the original ode results in

$$\frac{v''(x) + 2v(x)}{x} = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$v''(x) + 2v(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 2$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + 2e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(2)} \\ &= \pm i\sqrt{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i\sqrt{2} \\ \lambda_2 &= -i\sqrt{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{2}$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x))$$

Or

$$v(x) = c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} \xi &= v(x)z(x) \\ &= (c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x))(z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \frac{1}{x}$$

Hence (7) becomes

$$\xi = \frac{c_1 \cos(\sqrt{2}x) + c_2 \sin(\sqrt{2}x)}{x}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x)y' - y\xi'(x) + \xi(x)p(x)y &= \int \xi(x)r(x)dx \\ y' + y\left(p(x) - \frac{\xi'(x)}{\xi(x)}\right) &= \frac{\int \xi(x)r(x)dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(-\frac{2}{x} - \frac{\left(\frac{-c_1\sqrt{2}\sin(\sqrt{2}x) + c_2\sqrt{2}\cos(\sqrt{2}x)}{x} - \frac{c_1\cos(\sqrt{2}x) + c_2\sin(\sqrt{2}x)}{x^2} \right) x}{c_1\cos(\sqrt{2}x) + c_2\sin(\sqrt{2}x)} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{(\sqrt{2}xc_2 + c_1)\cos(\sqrt{2}x) - \sin(\sqrt{2}x)(xc_1\sqrt{2} - c_2)}{x(c_1\cos(\sqrt{2}x) + c_2\sin(\sqrt{2}x))} \\ p(x) &= 0 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{(\sqrt{2}xc_2 + c_1)\cos(\sqrt{2}x) - \sin(\sqrt{2}x)(xc_1\sqrt{2} - c_2)}{x(c_1\cos(\sqrt{2}x) + c_2\sin(\sqrt{2}x))} dx} \\ &= \frac{\sec\left(\frac{\sqrt{2}x}{2}\right)^2}{x\left(c_1\tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2\tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right)} \end{aligned}$$

The ode becomes

$$\frac{d}{dx} \left(\frac{y \sec\left(\frac{\sqrt{2}x}{2}\right)^2}{x\left(c_1\tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2\tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right)} \right) = 0$$

Integrating gives

$$\frac{y \sec\left(\frac{\sqrt{2}x}{2}\right)^2}{x\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right)} = \int 0 dx + c_3$$

$$= c_3$$

Dividing throughout by the integrating factor $\frac{\sec\left(\frac{\sqrt{2}x}{2}\right)^2}{x\left(c_1 \tan\left(\frac{\sqrt{2}x}{2}\right)^2 - 2c_2 \tan\left(\frac{\sqrt{2}x}{2}\right) - c_1\right)}$ gives the final solution

$$y = c_3 x \left(-c_1 \cos(\sqrt{2}x) - c_2 \sin(\sqrt{2}x)\right)$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = c_3 x \left(-c_1 \cos(\sqrt{2}x) - c_2 \sin(\sqrt{2}x)\right)$$

The constants can be merged to give

$$y = x \left(-c_1 \cos(\sqrt{2}x) - c_2 \sin(\sqrt{2}x)\right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x \left(-c_1 \cos(\sqrt{2}x) - c_2 \sin(\sqrt{2}x)\right)$$

Maple step by step solution

Let's solve

$$x^2 \left(\frac{d^2}{dx^2} y(x)\right) - 2x \left(\frac{d}{dx} y(x)\right) + 2(x^2 + 1)y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = -\frac{2(x^2+1)y(x)}{x^2} + \frac{2\left(\frac{d}{dx} y(x)\right)}{x}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is lin

$$\frac{d^2}{dx^2} y(x) - \frac{2\left(\frac{d}{dx} y(x)\right)}{x} + \frac{2(x^2+1)y(x)}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{2}{x}, P_3(x) = \frac{2(x^2+1)}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) - 2x \left(\frac{d}{dx} y(x) \right) + (2x^2 + 2) y(x) = 0$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y(x)$ to series expansion for $m = 0..2$

$$x^m \cdot y(x) = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y(x) = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot \left(\frac{d}{dx} y(x) \right)$ to series expansion

$$x \cdot \left(\frac{d}{dx} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right)$ to series expansion

$$x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-1+r)(-2+r)x^r + a_1r(-1+r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r-1)(k+r-2) + 2a_{k-2}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-1+r)(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{1, 2\}$$

- Each term must be 0

$$a_1r(-1+r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r-1)(k+r-2) + 2a_{k-2} = 0$$

- Shift index using $k- > k + 2$

$$a_{k+2}(k+1+r)(k+r) + 2a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{2a_k}{(k+1+r)(k+r)}$$

- Recursion relation for $r = 1$

$$a_{k+2} = -\frac{2a_k}{(k+2)(k+1)}$$

- Solution for $r = 1$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+2} = -\frac{2a_k}{(k+2)(k+1)}, a_1 = 0 \right]$$

- Recursion relation for $r = 2$

$$a_{k+2} = -\frac{2a_k}{(k+3)(k+2)}$$

- Solution for $r = 2$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+2} = -\frac{2a_k}{(k+3)(k+2)}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y(x) = \left(\sum_{k=0}^{\infty} a_k x^{k+1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+2} \right), a_{k+2} = -\frac{2a_k}{(k+2)(k+1)}, a_1 = 0, b_{k+2} = -\frac{2b_k}{(k+3)(k+2)}, b_1 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacic's algorithm successful`

```

Maple dsolve solution

Solving time : 0.013 (sec)

Leaf size : 23

```

dsolve(x^2*diff(diff(y(x),x),x)-2*diff(y(x),x)*x+2*(x^2+1)*y(x) = 0,
        y(x),singsol=all)

```

$$y = x \left(c_1 \sin(\sqrt{2}x) + c_2 \cos(\sqrt{2}x) \right)$$

Mathematica DSolve solution

Solving time : 0.069 (sec)

Leaf size : 48

```

DSolve[{x^2*D[y[x],{x,2}]-2*x*D[y[x],x]+2*(1+x^2)*y[x]==0,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow c_1 e^{-i\sqrt{2}x} x - \frac{ic_2 e^{i\sqrt{2}x} x}{2\sqrt{2}}$$

2.2.35 problem 35

Solved as second order ode using Kovacic algorithm	808
Solved as second order ode adjoint method	812
Maple step by step solution	817
Maple trace	818
Maple dsolve solution	819
Mathematica DSolve solution	819

Internal problem ID [8834]

Book : Second order enumerated odes

Section : section 2

Problem number : 35

Date solved : Thursday, December 12, 2024 at 09:53:35 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$4x^2y'' + 4x^5y' + (x^8 + 6x^4 + 4)y = 0$$

Solved as second order ode using Kovacic algorithm

Time used: 0.261 (sec)

Writing the ode as

$$4x^2y'' + 4x^5y' + (x^8 + 6x^4 + 4)y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 4x^2$$

$$B = 4x^5 \quad (3)$$

$$C = x^8 + 6x^4 + 4$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = x^2$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{x^2}\right)z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.90: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -1$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -1$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + \frac{i\sqrt{3}}{2}$	$\frac{1}{2} - \frac{i\sqrt{3}}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + \frac{i\sqrt{3}}{2}$	$\frac{1}{2} - \frac{i\sqrt{3}}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{2} - \frac{i\sqrt{3}}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{2} - \frac{i\sqrt{3}}{2} - \left(\frac{1}{2} - \frac{i\sqrt{3}}{2} \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= \frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} + (-) (0) \\ &= \frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} \\ &= \frac{1 - i\sqrt{3}}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2 \left(\frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} \right) (0) + \left(\left(-\frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x^2} \right) + \left(\frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} \right)^2 - \left(-\frac{1}{x^2} \right) \right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} dx} \\ &= x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4x^5}{4x^2} dx} \\ &= z_1 e^{-\frac{x^4}{8}} \\ &= z_1 \left(e^{-\frac{x^4}{8}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{4x^5}{4x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\frac{x^4}{4}}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{ix\sqrt{3} e^{-\frac{x^4}{4}} e^{\frac{x^4}{4}} x^{i\sqrt{3}-1}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \right) + c_2 \left(e^{-\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(-\frac{ix\sqrt{3} e^{-\frac{x^4}{4}} e^{\frac{x^4}{4}} x^{i\sqrt{3}-1}}{3} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 e^{-\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} - \frac{ic_2 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \sqrt{3} e^{-\frac{x^4}{8}}}{3}$$

Solved as second order ode adjoint method

Time used: 1.031 (sec)

In normal form the ode

$$4x^2y'' + 4x^5y' + (x^8 + 6x^4 + 4)y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= x^3 \\ q(x) &= \frac{x^8 + 6x^4 + 4}{4x^2} \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - (x^3\xi(x))' + \left(\frac{(x^8 + 6x^4 + 4)\xi(x)}{4x^2} \right) &= 0 \\ \xi''(x) - x^3\xi'(x) + \frac{(x^8 - 6x^4 + 4)\xi(x)}{4x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi'' - x^3\xi' + \left(\frac{x^6}{4} - \frac{3x^2}{2} + \frac{1}{x^2} \right) \xi = 0 \quad (1)$$

$$A\xi'' + B\xi' + C\xi = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -x^3 \\ C &= \frac{x^6}{4} - \frac{3x^2}{2} + \frac{1}{x^2} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = \xi e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then ξ is found using the inverse transformation

$$\xi = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.91: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -1$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -1$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + \frac{i\sqrt{3}}{2}$	$\frac{1}{2} - \frac{i\sqrt{3}}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + \frac{i\sqrt{3}}{2}$	$\frac{1}{2} - \frac{i\sqrt{3}}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{2} - \frac{i\sqrt{3}}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{2} - \frac{i\sqrt{3}}{2} - \left(\frac{1}{2} - \frac{i\sqrt{3}}{2} \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= \frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} + (-) (0) \\ &= \frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} \\ &= \frac{1 - i\sqrt{3}}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2 \left(\frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} \right) (0) + \left(\left(-\frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x^2} \right) + \left(\frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} \right)^2 - \left(-\frac{1}{x^2} \right) \right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \frac{\frac{1}{2} - \frac{i\sqrt{3}}{2}}{x} dx} \\ &= x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \end{aligned}$$

The first solution to the original ode in ξ is found from

$$\begin{aligned} \xi_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x^3}{1} dx} \\ &= z_1 e^{\frac{x^4}{8}} \\ &= z_1 \left(e^{\frac{x^4}{8}} \right) \end{aligned}$$

Which simplifies to

$$\xi_1 = e^{\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}}$$

The second solution ξ_2 to the original ode is found using reduction of order

$$\xi_2 = \xi_1 \int \frac{e^{\int -\frac{B}{A} dx}}{\xi_1^2} dx$$

Substituting gives

$$\begin{aligned} \xi_2 &= \xi_1 \int \frac{e^{\int -\frac{-x^3}{1} dx}}{(\xi_1)^2} dx \\ &= \xi_1 \int \frac{e^{\frac{x^4}{4}}}{(\xi_1)^2} dx \\ &= \xi_1 \left(-\frac{ix\sqrt{3} e^{-\frac{x^4}{4}} e^{\frac{x^4}{4}} x^{i\sqrt{3}-1}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} \xi &= c_1 \xi_1 + c_2 \xi_2 \\ &= c_1 \left(e^{\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \right) + c_2 \left(e^{\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(-\frac{ix\sqrt{3} e^{-\frac{x^4}{4}} e^{\frac{x^4}{4}} x^{i\sqrt{3}-1}}{3} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(x^3 - \frac{c_1 x^3 e^{\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}}}{2} + \frac{c_1 e^{\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(\frac{1}{2} - \frac{i\sqrt{3}}{2} \right)}{x} - \frac{ic_2 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(\frac{1}{2} + \frac{i\sqrt{3}}{2} \right) \sqrt{3} e^{\frac{x^4}{8}}}{3x} - \frac{ic_2 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \sqrt{3} x^3 e^{\frac{x^4}{8}}}{6} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = - \frac{c_1 x^{-\frac{i\sqrt{3}}{2}} (-ix^4 + i + \sqrt{3}) + x^{\frac{i\sqrt{3}}{2}} c_2 \left(i + \frac{(-x^4+1)\sqrt{3}}{3} \right)}{2x \left(ix^{-\frac{i\sqrt{3}}{2}} c_1 + x^{\frac{i\sqrt{3}}{2}} \frac{\sqrt{3} c_2}{3} \right)}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{-\int \frac{c_1 x^{-\frac{i\sqrt{3}}{2}} (-ix^4 + i + \sqrt{3}) + x^{\frac{i\sqrt{3}}{2}} c_2 \left(i + \frac{(-x^4+1)\sqrt{3}}{3} \right)}{2x \left(ix^{-\frac{i\sqrt{3}}{2}} c_1 + x^{\frac{i\sqrt{3}}{2}} \frac{\sqrt{3} c_2}{3} \right)} dx} \\ &= e^{\frac{x^{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} e^{\frac{x^4}{8}}}{i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{y x^{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} e^{\frac{x^4}{8}}}{i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1} \right) &= 0\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y x^{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} e^{\frac{x^4}{8}}}{i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1} &= \int 0 dx + c_3 \\ &= c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{x^{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} e^{\frac{x^4}{8}}}{i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1}$ gives the final solution

$$y = x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} e^{-\frac{x^4}{8}} \left(i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1 \right) c_3$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} e^{-\frac{x^4}{8}} \left(i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1 \right) c_3$$

The constants can be merged to give

$$y = x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} e^{-\frac{x^4}{8}} \left(i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1 \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} e^{-\frac{x^4}{8}} \left(i\sqrt{3} c_2 - 3x^{-i\sqrt{3}} c_1 \right)$$

Maple step by step solution

Let's solve

$$4x^2 \left(\frac{d^2}{dx^2} y(x) \right) + 4x^5 \left(\frac{d}{dx} y(x) \right) + (x^8 + 6x^4 + 4) y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = -\frac{(x^8+6x^4+4)y(x)}{4x^2} - \left(\frac{d}{dx} y(x) \right) x^3$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is lin

$$\frac{d^2}{dx^2} y(x) + \left(\frac{d}{dx} y(x) \right) x^3 + \frac{(x^8+6x^4+4)y(x)}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = x^3, P_3(x) = \frac{x^8+6x^4+4}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2 \left(\frac{d^2}{dx^2} y(x) \right) + 4x^5 \left(\frac{d}{dx} y(x) \right) + (x^8 + 6x^4 + 4) y(x) = 0$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y(x)$ to series expansion for $m = 0..8$

$$x^m \cdot y(x) = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y(x) = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^5 \cdot \left(\frac{d}{dx} y(x) \right)$ to series expansion

$$x^5 \cdot \left(\frac{d}{dx} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r+4}$$

- Shift index using $k \rightarrow k - 4$

$$x^5 \cdot \left(\frac{d}{dx} y(x) \right) = \sum_{k=4}^{\infty} a_{k-4} (k-4+r) x^{k+r}$$

- Convert $x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right)$ to series expansion

$$x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$4a_0(r^2 - r + 1) x^r + 4a_1(r^2 + r + 1) x^{1+r} + 4a_2(r^2 + 3r + 3) x^{2+r} + 4a_3(r^2 + 5r + 7) x^{3+r} + (4a_4$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$4r^2 - 4r + 4 = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ \frac{1}{2} - \frac{i\sqrt{3}}{2}, \frac{1}{2} + \frac{i\sqrt{3}}{2} \right\}$$

- The coefficients of each power of x must be 0

$$[4a_1(r^2 + r + 1) = 0, 4a_2(r^2 + 3r + 3) = 0, 4a_3(r^2 + 5r + 7) = 0, 4a_4(r^2 + 7r + 13) + 2a_0(3 + 2r) = 0]$$

- Solve for the dependent coefficient(s)

$$\left\{ a_1 = 0, a_2 = 0, a_3 = 0, a_4 = -\frac{a_0(3+2r)}{2(r^2+7r+13)}, a_5 = 0, a_6 = 0, a_7 = 0 \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$4(1 + k^2 + (2r - 1)k + r^2 - r)a_k + 2a_{k-4}(2k - 5 + 2r) + a_{k-8} = 0$$

- Shift index using $k \rightarrow k + 8$

$$4(1 + (k + 8)^2 + (2r - 1)(k + 8) + r^2 - r)a_{k+8} + 2a_{k+4}(2k + 11 + 2r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+8} = -\frac{4ka_{k+4} + 4ra_{k+4} + a_k + 22a_{k+4}}{4(k^2 + 2kr + r^2 + 15k + 15r + 57)}$$

- Recursion relation for $r = \frac{1}{2} - \frac{i\sqrt{3}}{2}$

$$a_{k+8} = -\frac{4ka_{k+4} + 4\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)a_{k+4} + a_k + 22a_{k+4}}{4\left(k^2 + 2k\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right) + \left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2 + 15k + \frac{129}{2} - \frac{15i\sqrt{3}}{2}\right)}$$

- Solution for $r = \frac{1}{2} - \frac{i\sqrt{3}}{2}$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k + \frac{1}{2} - \frac{i\sqrt{3}}{2}}, a_{k+8} = -\frac{4ka_{k+4} + 4\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)a_{k+4} + a_k + 22a_{k+4}}{4\left(k^2 + 2k\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right) + \left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2 + 15k + \frac{129}{2} - \frac{15i\sqrt{3}}{2}\right)}, a_1 = 0, a_2 = 0, a_3 = 0, a_4 = 0 \right]$$

- Recursion relation for $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$

$$a_{k+8} = -\frac{4ka_{k+4} + 4\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)a_{k+4} + a_k + 22a_{k+4}}{4\left(k^2 + 2k\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right) + \left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2 + 15k + \frac{129}{2} + \frac{15i\sqrt{3}}{2}\right)}$$

- Solution for $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k + \frac{1}{2} + \frac{i\sqrt{3}}{2}}, a_{k+8} = -\frac{4ka_{k+4} + 4\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)a_{k+4} + a_k + 22a_{k+4}}{4\left(k^2 + 2k\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right) + \left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2 + 15k + \frac{129}{2} + \frac{15i\sqrt{3}}{2}\right)}, a_1 = 0, a_2 = 0, a_3 = 0, a_4 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y(x) = \left(\sum_{k=0}^{\infty} a_k x^{k + \frac{1}{2} - \frac{i\sqrt{3}}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k + \frac{1}{2} + \frac{i\sqrt{3}}{2}} \right), a_{k+8} = -\frac{4ka_{k+4} + 4\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)a_{k+4} + a_k + 22a_{k+4}}{4\left(k^2 + 2k\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right) + \left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2 + 15k + \frac{129}{2} - \frac{15i\sqrt{3}}{2}\right)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
A Liouvillian solution exists
Group is reducible or imprimitive
<- Kovacic's algorithm successful`

```

Maple dsolve solution

Solving time : 0.015 (sec)

Leaf size : 33

```
dsolve(4*x^2*diff(diff(y(x),x),x)+4*x^5*diff(y(x),x)+(x^8+6*x^4+4)*y(x) = 0,
y(x),singsol=all)
```

$$y = \sqrt{x} e^{-\frac{x^4}{8}} \left(c_1 x^{\frac{i\sqrt{3}}{2}} + c_2 x^{-\frac{i\sqrt{3}}{2}} \right)$$

Mathematica DSolve solution

Solving time : 0.112 (sec)

Leaf size : 62

```
DSolve[{4*x^2*D[y[x],{x,2}]+4*x^5*D[y[x],x]+(x^8+6*x^4+4)*y[x]==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{3} e^{-\frac{x^4}{8}} x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(3c_1 - i\sqrt{3}c_2 x^{i\sqrt{3}} \right)$$

2.2.36 problem 36

Maple step by step solution	820
Maple trace	820
Maple dsolve solution	820
Mathematica DSolve solution	821

Internal problem ID [8835]

Book : Second order enumerated odes

Section : section 2

Problem number : 36

Date solved : Wednesday, December 18, 2024 at 02:16:41 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$x^2 y'' + (xy' - y)^2 = 0$$

Maple step by step solution**Maple trace**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrati
    --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for d
    -> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integr
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
<- linear symmetries successful`

```

Maple dsolve solution

Solving time : 0.057 (sec)

Leaf size : 22

```

dsolve(x^2*diff(diff(y(x),x),x)+(diff(y(x),x)*x-y(x))^2 = 0,
y(x),singsol=all)

```

$$y = \left(-e^{c_1} \operatorname{Ei}_1 \left(-\ln \left(\frac{1}{x} \right) + c_1 \right) + c_2 \right) x$$

Mathematica DSolve solution

Solving time : 43.474 (sec)

Leaf size : 33

```
DSolve[{x^2*D[y[x],{x,2}]+(x*D[y[x],x]-y[x])^2==0,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x(e^{c_1} \text{ExpIntegralEi}(-c_1 - \log(x)) + c_2)$$

$$y(x) \rightarrow c_2 x$$

2.2.37 problem 37

Solved as second order ode using change of variable on y	
method 1	822
Solved as second order Bessel ode	824
Solved as second order ode using Kovacic algorithm	824
Solved as second order ode adjoint method	827
Maple step by step solution	830
Maple trace	831
Maple dsolve solution	832
Mathematica DSolve solution	832

Internal problem ID [8836]

Book : Second order enumerated odes

Section : section 2

Problem number : 37

Date solved : Thursday, December 12, 2024 at 09:53:38 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$xy'' + 2y' - xy = 0$$

Solved as second order ode using change of variable on y method 1

Time used: 0.306 (sec)

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{2}{x} \\ q(x) &= -1 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= -1 - \frac{\left(\frac{2}{x}\right)'}{2} - \frac{\left(\frac{2}{x}\right)^2}{4} \\ &= -1 - \frac{\left(-\frac{2}{x^2}\right)}{2} - \frac{\left(\frac{4}{x^2}\right)}{4} \\ &= -1 - \left(-\frac{1}{x^2}\right) - \frac{1}{x^2} \\ &= -1 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{2}{2x} dx} \\ &= \frac{1}{x} \end{aligned} \quad (5)$$

Hence (3) becomes

$$y = \frac{v(x)}{x} \quad (4)$$

Applying this change of variable to the original ode results in

$$v''(x) - v(x) = 0$$

Which is now solved for $v(x)$.

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$v(x) = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$v(x) = c_1 e^{(1)x} + c_2 e^{(-1)x}$$

Or

$$v(x) = c_1 e^x + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 e^x + c_2 e^{-x}) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \frac{1}{x}$$

Hence (7) becomes

$$y = \frac{c_1 e^x + c_2 e^{-x}}{x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 e^x + c_2 e^{-x}}{x}$$

Solved as second order Bessel ode

Time used: 0.067 (sec)

Writing the ode as

$$x^2 y'' + 2xy' - x^2 y = 0 \quad (1)$$

Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= -\frac{1}{2} \\ \beta &= i \\ n &= \frac{1}{2} \\ \gamma &= 1 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = \frac{ic_1 \sqrt{2} \sinh(x)}{\sqrt{x} \sqrt{\pi} \sqrt{ix}} - \frac{c_2 \sqrt{2} \cosh(x)}{\sqrt{x} \sqrt{\pi} \sqrt{ix}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{ic_1 \sqrt{2} \sinh(x)}{\sqrt{x} \sqrt{\pi} \sqrt{ix}} - \frac{c_2 \sqrt{2} \cosh(x)}{\sqrt{x} \sqrt{\pi} \sqrt{ix}}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.051 (sec)

Writing the ode as

$$xy'' + 2y' - xy = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= 2 \\ C &= -x \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.93: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{x} dx} \\ &= z_1 e^{-\ln(x)} \\ &= z_1 \left(\frac{1}{x} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{e^{-x}}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{x} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{2x}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{e^{-x}}{x} \right) + c_2 \left(\frac{e^{-x}}{x} \left(\frac{e^{2x}}{2} \right) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 e^{-x}}{x} + \frac{c_2 e^x}{2x}$$

Solved as second order ode adjoint method

Time used: 0.637 (sec)

In normal form the ode

$$xy'' + 2y' - xy = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{2}{x} \\ q(x) &= -1 \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(\frac{2\xi(x)}{x}\right)' + (-\xi(x)) &= 0 \\ \frac{\xi''(x)x^2 - \xi(x)x^2 - 2\xi'(x)x + 2\xi(x)}{x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x)\xi' + q(x)\xi = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{2}{x} \\ q(x) &= \frac{2}{x^2} - 1 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= \frac{2}{x^2} - 1 - \frac{\left(-\frac{2}{x}\right)'}{2} - \frac{\left(-\frac{2}{x}\right)^2}{4} \\ &= \frac{2}{x^2} - 1 - \frac{\left(\frac{2}{x^2}\right)}{2} - \frac{\left(\frac{4}{x^2}\right)}{4} \\ &= \frac{2}{x^2} - 1 - \left(\frac{1}{x^2}\right) - \frac{1}{x^2} \\ &= -1 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{-2}{x} dx} \\ &= x \end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = v(x) x \quad (4)$$

Applying this change of variable to the original ode results in

$$-x(v(x) - v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$-v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} - e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$v(x) = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$v(x) = c_1 e^{(1)x} + c_2 e^{(-1)x}$$

Or

$$v(x) = c_1 e^x + c_2 e^{-x}$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} \xi &= v(x) z(x) \\ &= (c_1 e^x + c_2 e^{-x}) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = x$$

Hence (7) becomes

$$\xi = (c_1 e^x + c_2 e^{-x}) x$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(\frac{2}{x} - \frac{(c_1 e^x - c_2 e^{-x}) x + c_1 e^x + c_2 e^{-x}}{(c_1 e^x + c_2 e^{-x}) x} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= -\frac{c_1(x-1)e^{2x} - c_2(x+1)}{(c_1 e^{2x} + c_2) x} \\ p(x) &= 0 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{c_1(x-1)e^{2x} - c_2(x+1)}{(c_1 e^{2x} + c_2) x} dx} \\ &= \frac{x e^x}{c_1 e^{2x} + c_2} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{yx e^x}{c_1 e^{2x} + c_2} \right) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{yx e^x}{c_1 e^{2x} + c_2} &= \int 0 dx + c_3 \\ &= c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{x e^x}{c_1 e^{2x} + c_2}$ gives the final solution

$$y = \frac{c_3(c_1 e^x + c_2 e^{-x})}{x}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{c_3(c_1 e^x + c_2 e^{-x})}{x}$$

The constants can be merged to give

$$y = \frac{c_1 e^x + c_2 e^{-x}}{x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 e^x + c_2 e^{-x}}{x}$$

Maple step by step solution

Let's solve

$$\left(\frac{d^2}{dx^2}y(x)\right)x + 2\frac{d}{dx}y(x) - xy(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2}y(x) = y(x) - \frac{2\left(\frac{d}{dx}y(x)\right)}{x}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dx^2}y(x) + \frac{2\left(\frac{d}{dx}y(x)\right)}{x} - y(x) = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{2}{x}, P_3(x) = -1]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$\left(\frac{d^2}{dx^2}y(x)\right)x + 2\frac{d}{dx}y(x) - xy(x) = 0$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y(x)$ to series expansion

$$x \cdot y(x) = \sum_{k=0}^{\infty} a_k x^{k+r+1}$$

- Shift index using $k- > k - 1$

$$x \cdot y(x) = \sum_{k=1}^{\infty} a_{k-1} x^{k+r}$$

- Convert $\frac{d}{dx}y(x)$ to series expansion

$$\frac{d}{dx}y(x) = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k + 1$

$$\frac{d}{dx}y(x) = \sum_{k=-1}^{\infty} a_{k+1}(k+r+1)x^{k+r}$$

- Convert $x \cdot \left(\frac{d^2}{dx^2}y(x)\right)$ to series expansion

$$x \cdot \left(\frac{d^2}{dx^2}y(x)\right) = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r-1}$$

- Shift index using $k- > k+1$

$$x \cdot \left(\frac{d^2}{dx^2}y(x)\right) = \sum_{k=-1}^{\infty} a_{k+1}(k+r+1)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0r(1+r)x^{-1+r} + a_1(1+r)(2+r)x^r + \left(\sum_{k=1}^{\infty} (a_{k+1}(k+r+1)(k+2+r) - a_{k-1})x^{k+r}\right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $r(1+r) = 0$
- Values of r that satisfy the indicial equation
 $r \in \{-1, 0\}$
- Each term must be 0
 $a_1(1+r)(2+r) = 0$
- Each term in the series must be 0, giving the recursion relation
 $a_{k+1}(k+r+1)(k+2+r) - a_{k-1} = 0$
- Shift index using $k- > k+1$
 $a_{k+2}(k+2+r)(k+3+r) - a_k = 0$
- Recursion relation that defines series solution to ODE
 $a_{k+2} = \frac{a_k}{(k+2+r)(k+3+r)}$
- Recursion relation for $r = -1$
 $a_{k+2} = \frac{a_k}{(k+1)(k+2)}$
- Solution for $r = -1$
 $\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = \frac{a_k}{(k+1)(k+2)}, 0 = 0 \right]$
- Recursion relation for $r = 0$
 $a_{k+2} = \frac{a_k}{(k+2)(k+3)}$
- Solution for $r = 0$
 $\left[y(x) = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = \frac{a_k}{(k+2)(k+3)}, 2a_1 = 0 \right]$
- Combine solutions and rename parameters
 $\left[y(x) = \left(\sum_{k=0}^{\infty} a_k x^{k-1}\right) + \left(\sum_{k=0}^{\infty} b_k x^k\right), a_{k+2} = \frac{a_k}{(k+2)(k+1)}, 0 = 0, b_{k+2} = \frac{b_k}{(k+3)(k+2)}, 2b_1 = 0 \right]$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
  A Liouvillian solution exists
  Reducible group (found an exponential solution)
  Reducible group (found another exponential solution)
<- Kovacic's algorithm successful`

```

Maple dsolve solution

Solving time : 0.008 (sec)

Leaf size : 17

```
dsolve(x*diff(diff(y(x),x),x)+2*diff(y(x),x)-x*y(x) = 0,  
y(x),singsol=all)
```

$$y = \frac{c_1 \sinh(x) + c_2 \cosh(x)}{x}$$

Mathematica DSolve solution

Solving time : 0.041 (sec)

Leaf size : 28

```
DSolve[{x*D[y[x]},{x,2]}+2*D[y[x],x]-x*y[x]==0,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{2c_1 e^{-x} + c_2 e^x}{2x}$$

2.2.38 problem 38

Solved as second order ode using change of variable on y	
method 1	833
Solved as second order Bessel ode	835
Solved as second order ode using Kovacic algorithm	836
Solved as second order ode adjoint method	838
Maple step by step solution	841
Maple trace	843
Maple dsolve solution	843
Mathematica DSolve solution	843

Internal problem ID [8837]

Book : Second order enumerated odes

Section : section 2

Problem number : 38

Date solved : Thursday, December 12, 2024 at 09:53:39 AM

CAS classification : [_Lienard]

Solve

$$xy'' + 2y' + xy = 0$$

Solved as second order ode using change of variable on y method 1

Time used: 0.313 (sec)

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{2}{x} \\ q(x) &= 1 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= 1 - \frac{\left(\frac{2}{x}\right)'}{2} - \frac{\left(\frac{2}{x}\right)^2}{4} \\ &= 1 - \frac{\left(-\frac{2}{x^2}\right)}{2} - \frac{\left(\frac{4}{x^2}\right)}{4} \\ &= 1 - \left(-\frac{1}{x^2}\right) - \frac{1}{x^2} \\ &= 1 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{2}{x} dx} \\ &= \frac{1}{x} \end{aligned} \quad (5)$$

Hence (3) becomes

$$y = \frac{v(x)}{x} \quad (4)$$

Applying this change of variable to the original ode results in

$$v''(x) + v(x) = 0$$

Which is now solved for $v(x)$.

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$v(x) = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 \cos(x) + c_2 \sin(x)) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \frac{1}{x}$$

Hence (7) becomes

$$y = \frac{c_1 \cos(x) + c_2 \sin(x)}{x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 \cos(x) + c_2 \sin(x)}{x}$$

Solved as second order Bessel ode

Time used: 0.063 (sec)

Writing the ode as

$$x^2 y'' + 2xy' + x^2 y = 0 \quad (1)$$

Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= -\frac{1}{2} \\ \beta &= 1 \\ n &= \frac{1}{2} \\ \gamma &= 1 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1 \sqrt{2} \sin(x)}{x\sqrt{\pi}} - \frac{c_2 \sqrt{2} \cos(x)}{x\sqrt{\pi}}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 \sqrt{2} \sin(x)}{x\sqrt{\pi}} - \frac{c_2 \sqrt{2} \cos(x)}{x\sqrt{\pi}}$$

Solved as second order ode using Kovacic algorithm

Time used: 0.108 (sec)

Writing the ode as

$$xy'' + 2y' + xy = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= 2 \\ C &= x \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.95: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{x} dx} \\ &= z_1 e^{-\ln(x)} \\ &= z_1 \left(\frac{1}{x} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{\cos(x)}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{x} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2\ln(x)}}{(y_1)^2} dx \\ &= y_1 (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{\cos(x)}{x} \right) + c_2 \left(\frac{\cos(x)}{x} (\tan(x)) \right) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 \cos(x)}{x} + \frac{c_2 \sin(x)}{x}$$

Solved as second order ode adjoint method

Time used: 0.927 (sec)

In normal form the ode

$$xy'' + 2y' + xy = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{2}{x} \\ q(x) &= 1 \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(\frac{2\xi(x)}{x}\right)' + (\xi(x)) &= 0 \\ \frac{\xi''(x)x^2 + \xi(x)x^2 - 2\xi'(x)x + 2\xi(x)}{x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. In normal form the given ode is written as

$$\xi'' + p(x)\xi' + q(x)\xi = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{2}{x} \\ q(x) &= \frac{2}{x^2} + 1 \end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= \frac{2}{x^2} + 1 - \frac{\left(-\frac{2}{x}\right)'}{2} - \frac{\left(-\frac{2}{x}\right)^2}{4} \\ &= \frac{2}{x^2} + 1 - \frac{\left(\frac{2}{x^2}\right)}{2} - \frac{\left(\frac{4}{x^2}\right)}{4} \\ &= \frac{2}{x^2} + 1 - \left(\frac{1}{x^2}\right) - \frac{1}{x^2} \\ &= 1 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$\xi = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{-2}{2} dx} \\ &= x \end{aligned} \quad (5)$$

Hence (3) becomes

$$\xi = v(x) x \quad (4)$$

Applying this change of variable to the original ode results in

$$x(v(x) + v''(x)) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$v(x) + v''(x) = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{x\lambda} + e^{x\lambda} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$v(x) = c_1 \cos(x) + c_2 \sin(x)$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned}\xi &= v(x) z(x) \\ &= (c_1 \cos(x) + c_2 \sin(x)) (z(x))\end{aligned}\tag{7}$$

But from (5)

$$z(x) = x$$

Hence (7) becomes

$$\xi = (c_1 \cos(x) + c_2 \sin(x)) x$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned}\xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)}\end{aligned}$$

Or

$$y' + y \left(\frac{2}{x} - \frac{(-c_1 \sin(x) + c_2 \cos(x)) x + c_1 \cos(x) + c_2 \sin(x)}{(c_1 \cos(x) + c_2 \sin(x)) x} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned}q(x) &= -\frac{(c_2 x - c_1) \cos(x) - \sin(x) (c_1 x + c_2)}{(c_1 \cos(x) + c_2 \sin(x)) x} \\ p(x) &= 0\end{aligned}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int q dx} \\ &= e^{\int -\frac{(c_2 x - c_1) \cos(x) - \sin(x) (c_1 x + c_2)}{(c_1 \cos(x) + c_2 \sin(x)) x} dx} \\ &= \frac{x \sec\left(\frac{x}{2}\right)^2}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{yx \sec\left(\frac{x}{2}\right)^2}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1} \right) &= 0\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{yx \sec\left(\frac{x}{2}\right)^2}{c_1 \tan\left(\frac{x}{2}\right)^2 - 2c_2 \tan\left(\frac{x}{2}\right) - c_1} &= \int 0 dx + c_3 \\ &= c_3\end{aligned}$$

Dividing throughout by the integrating factor $\frac{x \sec(\frac{x}{2})^2}{c_1 \tan(\frac{x}{2})^2 - 2c_2 \tan(\frac{x}{2}) - c_1}$ gives the final solution

$$y = \frac{c_3(-c_2 \sin(x) - c_1 \cos(x))}{x}$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \frac{c_3(-c_2 \sin(x) - c_1 \cos(x))}{x}$$

The constants can be merged to give

$$y = \frac{-c_2 \sin(x) - c_1 \cos(x)}{x}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{-c_2 \sin(x) - c_1 \cos(x)}{x}$$

Maple step by step solution

Let's solve

$$\left(\frac{d^2}{dx^2}y(x)\right)x + 2\frac{d}{dx}y(x) + xy(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2}y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2}y(x) = -y(x) - \frac{2\left(\frac{d}{dx}y(x)\right)}{x}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is lin

$$\frac{d^2}{dx^2}y(x) + \frac{2\left(\frac{d}{dx}y(x)\right)}{x} + y(x) = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{2}{x}, P_3(x) = 1\right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$\left(x \cdot P_2(x)\right)\Big|_{x=0} = 2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$\left(x^2 \cdot P_3(x)\right)\Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$\left(\frac{d^2}{dx^2}y(x)\right)x + 2\frac{d}{dx}y(x) + xy(x) = 0$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y(x)$ to series expansion

$$x \cdot y(x) = \sum_{k=0}^{\infty} a_k x^{k+r+1}$$

- Shift index using $k- > k-1$

$$x \cdot y(x) = \sum_{k=1}^{\infty} a_{k-1} x^{k+r}$$

- Convert $\frac{d}{dx}y(x)$ to series expansion

$$\frac{d}{dx}y(x) = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k+1$

$$\frac{d}{dx}y(x) = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1) x^{k+r}$$

- Convert $x \cdot \left(\frac{d^2}{dx^2}y(x)\right)$ to series expansion

$$x \cdot \left(\frac{d^2}{dx^2}y(x)\right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k- > k+1$

$$x \cdot \left(\frac{d^2}{dx^2}y(x)\right) = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(1+r) x^{-1+r} + a_1 (1+r)(2+r) x^r + \left(\sum_{k=1}^{\infty} (a_{k+1} (k+r+1)(k+2+r) + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 0\}$$

- Each term must be 0

$$a_1 (1+r)(2+r) = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1} (k+r+1)(k+2+r) + a_{k-1} = 0$$

- Shift index using $k- > k+1$

$$a_{k+2} (k+2+r)(k+3+r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+2+r)(k+3+r)}$$

- Recursion relation for $r = -1$

$$a_{k+2} = -\frac{a_k}{(k+1)(k+2)}$$

- Solution for $r = -1$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = -\frac{a_k}{(k+1)(k+2)}, 0 = 0 \right]$$

- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{a_k}{(k+2)(k+3)}$$

- Solution for $r = 0$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{(k+2)(k+3)}, 2a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y(x) = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^k \right), a_{k+2} = -\frac{a_k}{(k+2)(k+1)}, 0 = 0, b_{k+2} = -\frac{b_k}{(k+3)(k+2)}, 2b_1 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacic's algorithm successful`

```

Maple dsolve solution

Solving time : 0.008 (sec)

Leaf size : 17

```

dsolve(x*diff(diff(y(x),x),x)+2*diff(y(x),x)+x*y(x) = 0,
        y(x),singsol=all)

```

$$y = \frac{\sin(x)c_1 + \cos(x)c_2}{x}$$

Mathematica DSolve solution

Solving time : 0.043 (sec)

Leaf size : 37

```

DSolve[{x*D[y[x],{x,2}]+2*D[y[x],x]+x*y[x]==0,{}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{2c_1e^{-ix} - ic_2e^{ix}}{2x}$$

2.2.39 problem 39

Solved as first order linear ode	844
Solved as first order Exact ode	845
Maple step by step solution	848
Maple trace	849
Maple dsolve solution	849
Mathematica DSolve solution	849

Internal problem ID [8838]

Book : Second order enumerated odes

Section : section 2

Problem number : 39

Date solved : Tuesday, December 17, 2024 at 01:04:28 PM

CAS classification : [_linear]

Solve

$$y' + y \cot(x) = 2 \cos(x)$$

Solved as first order linear ode

Time used: 0.119 (sec)

In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= \cot(x) \\ p(x) &= 2 \cos(x) \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int \cot(x) dx} \\ &= \sin(x) \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= \mu p \\ \frac{d}{dx}(\mu y) &= (\mu)(2 \cos(x)) \\ \frac{d}{dx}(y \sin(x)) &= (\sin(x))(2 \cos(x)) \\ d(y \sin(x)) &= (2 \cos(x) \sin(x)) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y \sin(x) &= \int 2 \cos(x) \sin(x) dx \\ &= \sin(x)^2 + c_1 \end{aligned}$$

Dividing throughout by the integrating factor $\sin(x)$ gives the final solution

$$y = \sin(x) + c_1 \csc(x)$$




Figure 2.160: Slope field plot
 $y' + y \cot(x) = 2 \cos(x)$

Summary of solutions found

$$y = \sin(x) + c_1 \csc(x)$$

Solved as first order Exact ode

Time used: 0.121 (sec)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= (-y \cot(x) + 2 \cos(x)) dx \\ (y \cot(x) - 2 \cos(x)) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= y \cot(x) - 2 \cos(x) \\N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y \cot(x) - 2 \cos(x)) \\&= \cot(x)\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\&= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\&= 1((\cot(x)) - (0)) \\&= \cot(x)\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\&= e^{\int \cot(x) dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\ln(\sin(x))} \\&= \sin(x)\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\&= \sin(x)(y \cot(x) - 2 \cos(x)) \\&= \cos(x)(-2 \sin(x) + y)\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\&= \sin(x)(1) \\&= \sin(x)\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\(\cos(x)(-2 \sin(x) + y)) + (\sin(x)) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (2) w.r.t. y gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial y} dy &= \int \bar{N} dy \\ \int \frac{\partial \phi}{\partial y} dy &= \int \sin(x) dy \\ \phi &= y \sin(x) + f(x) \end{aligned} \quad (3)$$

Where $f(x)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t x gives

$$\frac{\partial \phi}{\partial x} = \cos(x) y + f'(x) \quad (4)$$

But equation (1) says that $\frac{\partial \phi}{\partial x} = \cos(x)(-2 \sin(x) + y)$. Therefore equation (4) becomes

$$\cos(x)(-2 \sin(x) + y) = \cos(x) y + f'(x) \quad (5)$$

Solving equation (5) for $f'(x)$ gives

$$f'(x) = -2 \cos(x) \sin(x)$$

Integrating the above w.r.t x gives

$$\begin{aligned} \int f'(x) dx &= \int (-\sin(2x)) dx \\ f(x) &= \frac{\cos(2x)}{2} + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(x)$ into equation (3) gives ϕ

$$\phi = y \sin(x) + \frac{\cos(2x)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into the constant c_1 gives the solution as

$$c_1 = y \sin(x) + \frac{\cos(2x)}{2}$$

Solving for y gives

$$y = -\frac{\cos(2x) - 2c_1}{2 \sin(x)}$$




Figure 2.161: Slope field plot
 $y' + y \cot(x) = 2 \cos(x)$

Summary of solutions found

$$y = -\frac{\cos(2x) - 2c_1}{2 \sin(x)}$$

Maple step by step solution

Let's solve

$$\frac{d}{dx}y(x) + y(x) \cot(x) = 2 \cos(x)$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx}y(x)$$

- Solve for the highest derivative

$$\frac{d}{dx}y(x) = -y(x) \cot(x) + 2 \cos(x)$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE

$$\frac{d}{dx}y(x) + y(x) \cot(x) = 2 \cos(x)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(\frac{d}{dx}y(x) + y(x) \cot(x) \right) = 2\mu(x) \cos(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(y(x) \mu(x))$

$$\mu(x) \left(\frac{d}{dx}y(x) + y(x) \cot(x) \right) = \left(\frac{d}{dx}y(x) \right) \mu(x) + y(x) \left(\frac{d}{dx}\mu(x) \right)$$

- Isolate $\frac{d}{dx}\mu(x)$

$$\frac{d}{dx}\mu(x) = \mu(x) \cot(x)$$

- Solve to find the integrating factor

$$\mu(x) = \sin(x)$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(y(x) \mu(x)) \right) dx = \int 2\mu(x) \cos(x) dx + C1$$

- Evaluate the integral on the lhs

$$y(x) \mu(x) = \int 2\mu(x) \cos(x) dx + C1$$

- Solve for $y(x)$

$$y(x) = \frac{\int 2\mu(x) \cos(x) dx + C1}{\mu(x)}$$

- Substitute $\mu(x) = \sin(x)$

$$y(x) = \frac{\int 2 \sin(x) \cos(x) dx + C1}{\sin(x)}$$

- Evaluate the integrals on the rhs

$$y(x) = \frac{\sin(x)^2 + C1}{\sin(x)}$$

- Simplify

$$y(x) = \sin(x) + C1 \csc(x)$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

Maple dsolve solution

Solving time : 0.002 (sec)

Leaf size : 16

```
dsolve(diff(y(x),x)+y(x)*cot(x) = 2*cos(x),
        y(x),singsol=all)
```

$$y = \csc(x) \left(-\cos(x)^2 + c_1 + \frac{1}{2} \right)$$

Mathematica DSolve solution

Solving time : 0.061 (sec)

Leaf size : 20

```
DSolve[{D[y[x],x]+y[x]*Cot[x]==2*Cos[x],{}}],
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow -\frac{1}{2} \csc(x)(\cos(2x) - 2c_1)$$

2.2.40 problem 40

Solved as first order Exact ode	850
Maple step by step solution	853
Maple trace	853
Maple dsolve solution	854
Mathematica DSolve solution	854

Internal problem ID [8839]

Book : Second order enumerated odes

Section : section 2

Problem number : 40

Date solved : Tuesday, December 17, 2024 at 01:04:30 PM

CAS classification : [_rational]

Solve

$$2xy^2 - y + (y^2 + x + y) y' = 0$$

Solved as first order Exact ode

Time used: 0.255 (sec)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (y^2 + x + y) dy &= (-2xy^2 + y) dx \\ (2xy^2 - y) dx + (y^2 + x + y) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 2x y^2 - y \\N(x, y) &= y^2 + x + y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2x y^2 - y) \\&= 4yx - 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y^2 + x + y) \\&= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\&= \frac{1}{y^2 + x + y} ((4yx - 1) - (1)) \\&= \frac{4yx - 2}{y^2 + x + y}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\&= \frac{1}{2x y^2 - y} ((1) - (4yx - 1)) \\&= -\frac{2}{y}\end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned}\mu &= e^{\int B \, dy} \\&= e^{\int -\frac{2}{y} \, dy}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2 \ln(y)} \\&= \frac{1}{y^2}\end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\&= \frac{1}{y^2} (2x y^2 - y) \\&= \frac{2yx - 1}{y}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{y^2}(y^2 + x + y) \\ &= \frac{y^2 + x + y}{y^2}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{2yx - 1}{y}\right) + \left(\frac{y^2 + x + y}{y^2}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{2yx - 1}{y} dx \\ \phi &= \frac{x(yx - 1)}{y} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= \frac{x^2}{y} - \frac{x(yx - 1)}{y^2} + f'(y) \\ &= \frac{x}{y^2} + f'(y)\end{aligned} \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{y^2 + x + y}{y^2}$. Therefore equation (4) becomes

$$\frac{y^2 + x + y}{y^2} = \frac{x}{y^2} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{y + 1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{y + 1}{y}\right) dy \\ f(y) &= y + \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{x(yx - 1)}{y} + y + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into the constant c_1 gives the solution as

$$c_1 = \frac{x(yx - 1)}{y} + y + \ln(y)$$




Figure 2.162: Slope field plot
 $2xy^2 - y + (y^2 + x + y)y' = 0$

Summary of solutions found

$$\frac{x(yx - 1)}{y} + y + \ln(y) = c_1$$

Maple step by step solution

Let's solve

$$2xy(x)^2 - y(x) + (y(x)^2 + x + y(x)) \left(\frac{d}{dx}y(x)\right) = 0$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx}y(x)$$

- Solve for the highest derivative

$$\frac{d}{dx}y(x) = \frac{-2xy(x)^2 + y(x)}{y(x)^2 + x + y(x)}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
<- exact successful`
```

Maple dsolve solution

Solving time : 0.062 (sec)

Leaf size : 28

```
dsolve(2*x*y(x)^2-y(x)+(y(x)^2+x+y(x))*diff(y(x),x) = 0,
       y(x),singsol=all)
```

$$y = e^{\text{RootOf}(x^2e^{-z}+e^{2-z}+c_1e^{-z}+e^{-z}-z-x)}$$

Mathematica DSolve solution

Solving time : 0.282 (sec)

Leaf size : 22

```
DSolve[{(2*x*y[x]^2-y[x])+(y[x]^2+x+y[x])*D[y[x],x]==0,{}},
       y[x],x,IncludeSingularSolutions->True]
```

$$\text{Solve}\left[x^2 - \frac{x}{y(x)} + y(x) + \log(y(x)) = c_1, y(x)\right]$$

2.2.41 problem 41

Solved as first order ode of type reduced Riccati 855
 Maple step by step solution 856
 Maple trace 856
 Maple dsolve solution 857
 Mathematica DSolve solution 857

Internal problem ID [8840]

Book : Second order enumerated odes

Section : section 2

Problem number : 41

Date solved : Tuesday, December 17, 2024 at 01:04:32 PM

CAS classification : [[_Riccati, _special]]

Solve

$$y' = x - y^2$$

Solved as first order ode of type reduced Riccati

Time used: 0.119 (sec)

This is reduced Riccati ode of the form

$$y' = a x^n + b y^2$$

Comparing the given ode to the above shows that

$$\begin{aligned} a &= 1 \\ b &= -1 \\ n &= 1 \end{aligned}$$

Since $n \neq -2$ then the solution of the reduced Riccati ode is given by

$$\begin{aligned} w &= \sqrt{x} \begin{cases} c_1 \text{BesselJ} \left(\frac{1}{2k}, \frac{1}{k} \sqrt{abx^k} \right) + c_2 \text{BesselY} \left(\frac{1}{2k}, \frac{1}{k} \sqrt{abx^k} \right) & ab > 0 \\ c_1 \text{BesselI} \left(\frac{1}{2k}, \frac{1}{k} \sqrt{-abx^k} \right) + c_2 \text{BesselK} \left(\frac{1}{2k}, \frac{1}{k} \sqrt{-abx^k} \right) & ab < 0 \end{cases} \quad (1) \\ y &= -\frac{1}{b} \frac{w'}{w} \\ k &= 1 + \frac{n}{2} \end{aligned}$$

Since $ab < 0$ then EQ(1) gives

$$\begin{aligned} k &= \frac{3}{2} \\ w &= \sqrt{x} \left(c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + c_2 \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) \right) \end{aligned}$$

Therefore the solution becomes

$$y = -\frac{1}{b} \frac{w'}{w}$$

Substituting the value of b, w found above and simplifying gives

$$y = \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_2 \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + c_2 \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}$$

Letting $c_2 = 1$ the above becomes

$$y = \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}$$




Figure 2.163: Slope field plot
 $y' = x - y^2$

Summary of solutions found

$$y = \frac{\sqrt{x} \left(\text{BesselI} \left(-\frac{2}{3}, \frac{2x^{3/2}}{3} \right) c_1 - \text{BesselK} \left(\frac{2}{3}, \frac{2x^{3/2}}{3} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right) + \text{BesselK} \left(\frac{1}{3}, \frac{2x^{3/2}}{3} \right)}$$

Maple step by step solution

Let's solve

$$\frac{d}{dx}y(x) = x - y(x)^2$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx}y(x)$$

- Solve for the highest derivative

$$\frac{d}{dx}y(x) = x - y(x)^2$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying Riccati
trying Riccati Special
<- Riccati Special successful`
```


Maple dsolve solution

Solving time : 0.003 (sec)

Leaf size : 23

```
dsolve(diff(y(x),x) = x-y(x)^2,
        y(x),singsol=all)
```

$$y = \frac{c_1 \text{AiryAi}(1, x) + \text{AiryBi}(1, x)}{c_1 \text{AiryAi}(x) + \text{AiryBi}(x)}$$

Mathematica DSolve solution

Solving time : 0.18 (sec)

Leaf size : 223

```
DSolve[{D[y[x],x]==x-y[x]^2,{}},
        y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{-ix^{3/2} \left(2 \text{BesselJ} \left(-\frac{2}{3}, \frac{2}{3} ix^{3/2} \right) + c_1 \left(\text{BesselJ} \left(-\frac{4}{3}, \frac{2}{3} ix^{3/2} \right) - \text{BesselJ} \left(\frac{2}{3}, \frac{2}{3} ix^{3/2} \right) \right) \right) - c_1 \text{BesselJ} \left(-\frac{1}{3}, \frac{2}{3} ix^{3/2} \right)}{2x \left(\text{BesselJ} \left(\frac{1}{3}, \frac{2}{3} ix^{3/2} \right) + c_1 \text{BesselJ} \left(-\frac{1}{3}, \frac{2}{3} ix^{3/2} \right) \right)}$$

$$y(x) \rightarrow \frac{ix^{3/2} \text{BesselJ} \left(-\frac{4}{3}, \frac{2}{3} ix^{3/2} \right) - ix^{3/2} \text{BesselJ} \left(\frac{2}{3}, \frac{2}{3} ix^{3/2} \right) + \text{BesselJ} \left(-\frac{1}{3}, \frac{2}{3} ix^{3/2} \right)}{2x \text{BesselJ} \left(-\frac{1}{3}, \frac{2}{3} ix^{3/2} \right)}$$

2.2.42 problem 42

Solved as higher order constant coeff ode	858
Maple step by step solution	860
Maple trace	860
Maple dsolve solution	860
Mathematica DSolve solution	860

Internal problem ID [8841]

Book : Second order enumerated odes

Section : section 2

Problem number : 42

Date solved : Thursday, December 12, 2024 at 09:53:46 AM

CAS classification : [[_high_order, _linear, _nonhomogeneous]]

Solve

$$y'''' - y''' - 3y'' + 5y' - 2y = x e^x + 3 e^{-2x}$$

Solved as higher order constant coeff ode

Time used: 0.162 (sec)

The characteristic equation is

$$\lambda^4 - \lambda^3 - 3\lambda^2 + 5\lambda - 2 = 0$$

The roots of the above equation are

$$\lambda_1 = -2$$

$$\lambda_2 = 1$$

$$\lambda_3 = 1$$

$$\lambda_4 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = e^{-2x} c_1 + e^x c_2 + x e^x c_3 + x^2 e^x c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-2x}$$

$$y_2 = e^x$$

$$y_3 = x e^x$$

$$y_4 = x^2 e^x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y'''' - y''' - 3y'' + 5y' - 2y = 0$$

Now the particular solution to the given ODE is found

$$y'''' - y''' - 3y'' + 5y' - 2y = (x e^{3x} + 3) e^{-2x}$$

The particular solution is now found using the method of undetermined coefficients.

Looking at the RHS of the ode, which is

$$(x e^{3x} + 3) e^{-2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-2x}\}, \{x e^x, e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^x, x^2 e^x, e^x, e^{-2x}\}$$

Since e^{-2x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-2x}\}, \{x e^x, e^x\}]$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-2x}\}, \{x e^x, x^2 e^x\}]$$

Since $x e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-2x}\}, \{x^2 e^x, x^3 e^x\}]$$

Since $x^2 e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-2x}\}, \{x^3 e^x, x^4 e^x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^{-2x} + A_2 x^3 e^x + A_3 x^4 e^x$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$72A_3 x e^x + 24A_3 e^x - 27A_1 e^{-2x} + 18A_2 e^x = x e^x + 3 e^{-2x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{9}, A_2 = -\frac{1}{54}, A_3 = \frac{1}{72} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x e^{-2x}}{9} - \frac{x^3 e^x}{54} + \frac{x^4 e^x}{72}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{-2x} c_1 + e^x c_2 + x e^x c_3 + x^2 e^x c_4) + \left(-\frac{x e^{-2x}}{9} - \frac{x^3 e^x}{54} + \frac{x^4 e^x}{72} \right) \end{aligned}$$

Maple step by step solution**Maple trace**

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 4; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

Maple dsolve solution

Solving time : 0.010 (sec)

Leaf size : 52

```

dsolve(diff(diff(diff(diff(y(x),x),x),x),x)-diff(diff(diff(y(x),x),x),x)-3*diff(diff(y(x),x),x),x),singsol=all)

```

$$y = \frac{e^{-2x} \left(\left(x^4 - \frac{4x^3}{3} + (72c_4 + \frac{4}{3})x^2 + (72c_3 - \frac{8}{9})x + 72c_1 + \frac{8}{27} \right) e^{3x} - 8x + 72c_2 - 8 \right)}{72}$$

Mathematica DSolve solution

Solving time : 0.408 (sec)

Leaf size : 64

```

DSolve[{D[y[x],{x,4}]-D[y[x],{x,3}]-3*D[y[x],{x,2}]+5*D[y[x],x]-2*y[x]==x*Exp[x]+3*Exp[-2*x],{y[x],x,IncludeSingularSolutions->True]}

```

$$y(x) \rightarrow e^x \left(\frac{x^4}{72} - \frac{x^3}{54} + \left(\frac{1}{54} + c_4 \right) x^2 + \left(-\frac{1}{81} + c_3 \right) x + \frac{1}{243} + c_2 \right) - \frac{1}{9} e^{-2x} (x + 1 - 9c_1)$$

2.2.43 problem 43

Maple step by step solution 869
 Maple trace 870
 Maple dsolve solution 870
 Mathematica DSolve solution 871

Internal problem ID [8842]

Book : Second order enumerated odes

Section : section 2

Problem number : 43

Date solved : Thursday, December 12, 2024 at 09:53:47 AM

CAS classification : [[_2nd_order, _with_linear_symmetries]]

Solve

$$x^2y'' - x(x + 6)y' + 10y = 0$$

Using series expansion around $x = 0$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (-x^2 - 6x)y' + 10y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{x + 6}{x}$$

$$q(x) = \frac{10}{x^2}$$

Table 2.100: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{x+6}{x}$		$q(x) = \frac{10}{x^2}$	
singularity	type	singularity	type
$x = 0$	“regular”	$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : [0]

Irregular singular points : [∞]

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2y'' + (-x^2 - 6x)y' + 10y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + (-x^2 - 6x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 10 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) + \sum_{n=0}^{\infty} (-6x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 10a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) = \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r}) + \sum_{n=0}^{\infty} (-6x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 10a_n x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r)(n+r-1) - 6x^{n+r} a_n (n+r) + 10a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) - 6x^r a_0 r + 10a_0 x^r = 0$$

Or

$$(x^r r(-1+r) - 6x^r r + 10x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r-2)(r-5)x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$(r-2)(r-5) = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 5 \\ r_2 &= 2 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r - 2)(r - 5)x^r = 0$$

Solving for r gives the roots of the indicial equation as $[5, 2]$.

Since $r_1 - r_2 = 3$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x^5 \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^2 \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+5} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+2} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) - 6a_n(n+r) + 10a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}(n+r-1)}{n^2 + 2nr + r^2 - 7n - 7r + 10} \quad (4)$$

Which for the root $r = 5$ becomes

$$a_n = \frac{a_{n-1}(n+4)}{n(n+3)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 5$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{r}{r^2 - 5r + 4}$$

Which for the root $r = 5$ becomes

$$a_1 = \frac{5}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2-5r+4}$	$\frac{5}{4}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1+r}{r^3-8r^2+19r-12}$$

Which for the root $r = 5$ becomes

$$a_2 = \frac{3}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2-5r+4}$	$\frac{5}{4}$
a_2	$\frac{1+r}{r^3-8r^2+19r-12}$	$\frac{3}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{2+r}{r^4-10r^3+35r^2-50r+24}$$

Which for the root $r = 5$ becomes

$$a_3 = \frac{7}{24}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2-5r+4}$	$\frac{5}{4}$
a_2	$\frac{1+r}{r^3-8r^2+19r-12}$	$\frac{3}{4}$
a_3	$\frac{2+r}{r^4-10r^3+35r^2-50r+24}$	$\frac{7}{24}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{3+r}{(-1+r)^2(r-2)(r-4)(r-3)}$$

Which for the root $r = 5$ becomes

$$a_4 = \frac{1}{12}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2-5r+4}$	$\frac{5}{4}$
a_2	$\frac{1+r}{r^3-8r^2+19r-12}$	$\frac{3}{4}$
a_3	$\frac{2+r}{r^4-10r^3+35r^2-50r+24}$	$\frac{7}{24}$
a_4	$\frac{3+r}{(-1+r)^2(r-2)(r-4)(r-3)}$	$\frac{1}{12}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{4+r}{r(-1+r)^2(r-2)(r-4)(r-3)}$$

Which for the root $r = 5$ becomes

$$a_5 = \frac{3}{160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2-5r+4}$	$\frac{5}{4}$
a_2	$\frac{1+r}{r^3-8r^2+19r-12}$	$\frac{3}{4}$
a_3	$\frac{2+r}{r^4-10r^3+35r^2-50r+24}$	$\frac{7}{24}$
a_4	$\frac{3+r}{(-1+r)^2(r-2)(r-4)(r-3)}$	$\frac{1}{12}$
a_5	$\frac{4+r}{r(-1+r)^2(r-2)(r-4)(r-3)}$	$\frac{3}{160}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^5(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^5 \left(1 + \frac{5x}{4} + \frac{3x^2}{4} + \frac{7x^3}{24} + \frac{x^4}{12} + \frac{3x^5}{160} + O(x^6) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 3$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_3(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_3 \\ &= \frac{2+r}{r^4 - 10r^3 + 35r^2 - 50r + 24} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{2+r}{r^4 - 10r^3 + 35r^2 - 50r + 24} &= \lim_{r \rightarrow 2} \frac{2+r}{r^4 - 10r^3 + 35r^2 - 50r + 24} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x^2y'' + (-x^2 - 6x)y' + 10y = 0$ gives

$$\begin{aligned} & x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ & \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & + (-x^2 - 6x) \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \right) \\ & + 10Cy_1(x) \ln(x) + 10 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} & \left((x^2y_1''(x) + (-x^2 - 6x)y_1'(x) + 10y_1(x)) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) \right. \\ & \left. + \frac{(-x^2 - 6x)y_1(x)}{x} \right) C + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \quad (7) \\ & + (-x^2 - 6x) \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + 10 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned}$$

But since $y_1(x)$ is a solution to the ode, then

$$x^2y_1''(x) + (-x^2 - 6x)y_1'(x) + 10y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + \frac{(-x^2 - 6x)y_1(x)}{x} \right) C \\ & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \quad (8) \\ & + (-x^2 - 6x) \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + 10 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned}$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x - (x+7) \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 \quad (9) \\ & + (-x^2 - 6x) \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) + 10 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned}$$

Since $r_1 = 5$ and $r_2 = 2$ then the above becomes

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{n+4} a_n (n+5) \right) x - (x+7) \left(\sum_{n=0}^{\infty} a_n x^{n+5} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^n b_n (n+2) (1+n) \right) x^2 \quad (10) \\ & + (-x^2 - 6x) \left(\sum_{n=0}^{\infty} x^{1+n} b_n (n+2) \right) + 10 \left(\sum_{n=0}^{\infty} b_n x^{n+2} \right) = 0 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
 & \left(\sum_{n=0}^{\infty} 2C x^{n+5} a_n (n+5) \right) + \sum_{n=0}^{\infty} (-C x^{n+6} a_n) + \sum_{n=0}^{\infty} (-7C x^{n+5} a_n) \\
 & + \left(\sum_{n=0}^{\infty} x^{n+2} b_n (n^2 + 3n + 2) \right) + \sum_{n=0}^{\infty} (-x^{n+3} b_n (n+2)) \\
 & + \sum_{n=0}^{\infty} (-6x^{n+2} b_n (n+2)) + \left(\sum_{n=0}^{\infty} 10b_n x^{n+2} \right) = 0
 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+2$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+2} and adjusting the power and the corresponding index gives

$$\begin{aligned}
 \sum_{n=0}^{\infty} 2C x^{n+5} a_n (n+5) &= \sum_{n=3}^{\infty} 2C a_{n-3} (n+2) x^{n+2} \\
 \sum_{n=0}^{\infty} (-C x^{n+6} a_n) &= \sum_{n=4}^{\infty} (-C a_{n-4} x^{n+2}) \\
 \sum_{n=0}^{\infty} (-7C x^{n+5} a_n) &= \sum_{n=3}^{\infty} (-7C a_{n-3} x^{n+2}) \\
 \sum_{n=0}^{\infty} (-x^{n+3} b_n (n+2)) &= \sum_{n=1}^{\infty} (-b_{n-1} (1+n) x^{n+2})
 \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+2$.

$$\begin{aligned}
 & \left(\sum_{n=3}^{\infty} 2C a_{n-3} (n+2) x^{n+2} \right) + \sum_{n=4}^{\infty} (-C a_{n-4} x^{n+2}) + \sum_{n=3}^{\infty} (-7C a_{n-3} x^{n+2}) \\
 & + \left(\sum_{n=0}^{\infty} x^{n+2} b_n (n^2 + 3n + 2) \right) + \sum_{n=1}^{\infty} (-b_{n-1} (1+n) x^{n+2}) \\
 & + \sum_{n=0}^{\infty} (-6x^{n+2} b_n (n+2)) + \left(\sum_{n=0}^{\infty} 10b_n x^{n+2} \right) = 0
 \end{aligned} \tag{2B}$$

For $n=0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n=1$, Eq (2B) gives

$$-2b_1 - 2b_0 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-2b_1 - 2 = 0$$

Solving the above for b_1 gives

$$b_1 = -1$$

For $n=2$, Eq (2B) gives

$$-2b_2 - 3b_1 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-2b_2 + 3 = 0$$

Solving the above for b_2 gives

$$b_2 = \frac{3}{2}$$

For $n = N$, where $N = 3$ which is the difference between the two roots, we are free to choose $b_3 = 0$. Hence for $n = 3$, Eq (2B) gives

$$3C - 6 = 0$$

Which is solved for C . Solving for C gives

$$C = 2$$

For $n = 4$, Eq (2B) gives

$$(-a_0 + 5a_1)C - 5b_3 + 4b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{21}{2} + 4b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{21}{8}$$

For $n = 5$, Eq (2B) gives

$$(-a_1 + 7a_2)C - 6b_4 + 10b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{95}{4} + 10b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = -\frac{19}{8}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = 2$ and all b_n , then the second solution becomes

$$y_2(x) = 2 \left(x^5 \left(1 + \frac{5x}{4} + \frac{3x^2}{4} + \frac{7x^3}{24} + \frac{x^4}{12} + \frac{3x^5}{160} + O(x^6) \right) \right) \ln(x) \\ + x^2 \left(1 - x + \frac{3x^2}{2} - \frac{21x^4}{8} - \frac{19x^5}{8} + O(x^6) \right)$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x) \\ = c_1 x^5 \left(1 + \frac{5x}{4} + \frac{3x^2}{4} + \frac{7x^3}{24} + \frac{x^4}{12} + \frac{3x^5}{160} + O(x^6) \right) \\ + c_2 \left(2 \left(x^5 \left(1 + \frac{5x}{4} + \frac{3x^2}{4} + \frac{7x^3}{24} + \frac{x^4}{12} + \frac{3x^5}{160} + O(x^6) \right) \right) \ln(x) \right. \\ \left. + x^2 \left(1 - x + \frac{3x^2}{2} - \frac{21x^4}{8} - \frac{19x^5}{8} + O(x^6) \right) \right)$$

Hence the final solution is

$$y = y_h \\ = c_1 x^5 \left(1 + \frac{5x}{4} + \frac{3x^2}{4} + \frac{7x^3}{24} + \frac{x^4}{12} + \frac{3x^5}{160} + O(x^6) \right) \\ + c_2 \left(2x^5 \left(1 + \frac{5x}{4} + \frac{3x^2}{4} + \frac{7x^3}{24} + \frac{x^4}{12} + \frac{3x^5}{160} + O(x^6) \right) \ln(x) \right. \\ \left. + x^2 \left(1 - x + \frac{3x^2}{2} - \frac{21x^4}{8} - \frac{19x^5}{8} + O(x^6) \right) \right)$$

Maple step by step solution

Let's solve

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) - x(6+x) \left(\frac{d}{dx} y(x) \right) + 10y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = -\frac{10y(x)}{x^2} + \frac{(6+x) \left(\frac{d}{dx} y(x) \right)}{x}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is lin

$$\frac{d^2}{dx^2} y(x) - \frac{(6+x) \left(\frac{d}{dx} y(x) \right)}{x} + \frac{10y(x)}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{6+x}{x}, P_3(x) = \frac{10}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$\left(x \cdot P_2(x) \right) \Big|_{x=0} = -6$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$\left(x^2 \cdot P_3(x) \right) \Big|_{x=0} = 10$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) - x(6+x) \left(\frac{d}{dx} y(x) \right) + 10y(x) = 0$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot \left(\frac{d}{dx} y(x) \right)$ to series expansion for $m = 1..2$

$$x^m \cdot \left(\frac{d}{dx} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k- > k+1-m$

$$x^m \cdot \left(\frac{d}{dx} y(x) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right)$ to series expansion

$$x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-2+r)(-5+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r-2)(k+r-5) - a_{k-1}(k+r-1)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-2+r)(-5+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{2, 5\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r-2)(k+r-5) - a_{k-1}(k+r-1) = 0$$

- Shift index using $k- > k+1$

$$a_{k+1}(k+r-1)(k-4+r) - a_k(k+r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(k+r)}{(k+r-1)(k-4+r)}$$

- Recursion relation for $r = 2$

$$a_{k+1} = \frac{a_k(k+2)}{(k+1)(k-2)}$$

- Series not valid for $r = 2$, division by 0 in the recursion relation at $k = 2$

$$a_{k+1} = \frac{a_k(k+2)}{(k+1)(k-2)}$$

- Recursion relation for $r = 5$

$$a_{k+1} = \frac{a_k(k+5)}{(k+4)(k+1)}$$

- Solution for $r = 5$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k+5}, a_{k+1} = \frac{a_k(k+5)}{(k+4)(k+1)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacic's algorithm successful`

```

Maple dsolve solution

Solving time : 0.027 (sec)

Leaf size : 62

```

dsolve(x^2*diff(diff(y(x),x),x)-x*(6+x)*diff(y(x),x)+10*y(x) = 0,y(x),
series,x=0)

```

$$y = \left(c_1 x^3 \left(1 + \frac{5}{4}x + \frac{3}{4}x^2 + \frac{7}{24}x^3 + \frac{1}{12}x^4 + \frac{3}{160}x^5 + O(x^6) \right) + c_2 (\ln(x) (24x^3 + 30x^4 + 18x^5 + O(x^6)) + (12 - 12x + 18x^2 + 26x^3 + x^4 - 9x^5 + O(x^6))) \right) x^2$$

Mathematica DSolve solution

Solving time : 0.04 (sec)

Leaf size : 84

```
AsymptoticDSolveValue[{x^2*D[y[x],{x,2}]-x*(x+6)*D[y[x],x]+10*y[x]==0,{}},  
y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{1}{2} x^5 (5x + 4) \log(x) - \frac{1}{4} x^2 (3x^4 - 6x^3 - 6x^2 + 4x - 4) \right) \\ + c_2 \left(\frac{x^9}{12} + \frac{7x^8}{24} + \frac{3x^7}{4} + \frac{5x^6}{4} + x^5 \right)$$

2.2.44 problem 44

Maple step by step solution 878
 Maple trace 879
 Maple dsolve solution 880
 Mathematica DSolve solution 880

Internal problem ID [8843]

Book : Second order enumerated odes

Section : section 2

Problem number : 44

Date solved : Thursday, December 12, 2024 at 09:53:48 AM

CAS classification : [_Bessel]

Solve

$$x^2y'' + xy' + (x^2 - 5)y = 0$$

Using series expansion around $x = 0$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + xy' + (x^2 - 5)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = \frac{x^2 - 5}{x^2}$$

Table 2.102: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$		$q(x) = \frac{x^2-5}{x^2}$	
singularity	type	singularity	type
$x = 0$	“regular”	$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : [0]

Irregular singular points : [∞]

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2y'' + xy' + (x^2 - 5)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x^2 - 5) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} (-5a_n x^{n+r}) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} (-5a_n x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r)(n+r-1) + x^{n+r} a_n (n+r) - 5a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - 5a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + x^r r - 5x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 5) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 5 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= \sqrt{5} \\ r_2 &= -\sqrt{5} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 5) x^r = 0$$

Solving for r gives the roots of the indicial equation as $[\sqrt{5}, -\sqrt{5}]$.

Since $r_1 - r_2 = 2\sqrt{5}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+\sqrt{5}} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-\sqrt{5}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} - 5a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 - 5} \quad (4)$$

Which for the root $r = \sqrt{5}$ becomes

$$a_n = -\frac{a_{n-2}}{n(2\sqrt{5} + n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \sqrt{5}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 + 4r - 1}$$

Which for the root $r = \sqrt{5}$ becomes

$$a_2 = -\frac{1}{4 + 4\sqrt{5}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r-1}$	$-\frac{1}{4+4\sqrt{5}}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r-1}$	$-\frac{1}{4+4\sqrt{5}}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r^2 + 4r - 1)(r^2 + 8r + 11)}$$

Which for the root $r = \sqrt{5}$ becomes

$$a_4 = \frac{1}{32(\sqrt{5} + 1)(2 + \sqrt{5})}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r-1}$	$-\frac{1}{4+4\sqrt{5}}$
a_3	0	0
a_4	$\frac{1}{(r^2+4r-1)(r^2+8r+11)}$	$\frac{1}{32(\sqrt{5}+1)(2+\sqrt{5})}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r-1}$	$-\frac{1}{4+4\sqrt{5}}$
a_3	0	0
a_4	$\frac{1}{(r^2+4r-1)(r^2+8r+11)}$	$\frac{1}{32(\sqrt{5}+1)(2+\sqrt{5})}$
a_5	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^{\sqrt{5}}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 \dots) \\ &= x^{\sqrt{5}}\left(1 - \frac{x^2}{4 + 4\sqrt{5}} + \frac{x^4}{32(\sqrt{5} + 1)(2 + \sqrt{5})} + O(x^6)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + b_n(n+r) + b_{n-2} - 5b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{b_{n-2}}{n^2 + 2nr + r^2 - 5} \quad (4)$$

Which for the root $r = -\sqrt{5}$ becomes

$$b_n = -\frac{b_{n-2}}{n(-2\sqrt{5} + n)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\sqrt{5}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{1}{r^2 + 4r - 1}$$

Which for the root $r = -\sqrt{5}$ becomes

$$b_2 = \frac{1}{-4 + 4\sqrt{5}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+4r-1}$	$\frac{1}{-4+4\sqrt{5}}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+4r-1}$	$\frac{1}{-4+4\sqrt{5}}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(r^2 + 4r - 1)(r^2 + 8r + 11)}$$

Which for the root $r = -\sqrt{5}$ becomes

$$b_4 = \frac{1}{32(\sqrt{5} - 1)(-2 + \sqrt{5})}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+4r-1}$	$\frac{1}{-4+4\sqrt{5}}$
b_3	0	0
b_4	$\frac{1}{(r^2+4r-1)(r^2+8r+11)}$	$\frac{1}{32(\sqrt{5}-1)(-2+\sqrt{5})}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+4r-1}$	$\frac{1}{-4+4\sqrt{5}}$
b_3	0	0
b_4	$\frac{1}{(r^2+4r-1)(r^2+8r+11)}$	$\frac{1}{32(\sqrt{5}-1)(-2+\sqrt{5})}$
b_5	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^{\sqrt{5}}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 \dots) \\ &= x^{-\sqrt{5}} \left(1 + \frac{x^2}{-4 + 4\sqrt{5}} + \frac{x^4}{32(\sqrt{5} - 1)(-2 + \sqrt{5})} + O(x^6) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^{\sqrt{5}} \left(1 - \frac{x^2}{4 + 4\sqrt{5}} + \frac{x^4}{32(\sqrt{5} + 1)(2 + \sqrt{5})} + O(x^6) \right) \\ &\quad + c_2 x^{-\sqrt{5}} \left(1 + \frac{x^2}{-4 + 4\sqrt{5}} + \frac{x^4}{32(\sqrt{5} - 1)(-2 + \sqrt{5})} + O(x^6) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 x^{\sqrt{5}} \left(1 - \frac{x^2}{4 + 4\sqrt{5}} + \frac{x^4}{32(\sqrt{5} + 1)(2 + \sqrt{5})} + O(x^6) \right) \\ &\quad + c_2 x^{-\sqrt{5}} \left(1 + \frac{x^2}{-4 + 4\sqrt{5}} + \frac{x^4}{32(\sqrt{5} - 1)(-2 + \sqrt{5})} + O(x^6) \right) \end{aligned}$$

Maple step by step solution

Let's solve

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) + x \left(\frac{d}{dx} y(x) \right) + (x^2 - 5) y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = -\frac{(x^2-5)y(x)}{x^2} - \frac{\frac{d}{dx} y(x)}{x}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dx^2} y(x) + \frac{\frac{d}{dx} y(x)}{x} + \frac{(x^2-5)y(x)}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{x^2-5}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -5$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) + x \left(\frac{d}{dx} y(x) \right) + (x^2 - 5) y(x) = 0$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y(x)$ to series expansion for $m = 0..2$

$$x^m \cdot y(x) = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k - > k - m$

$$x^m \cdot y(x) = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot \left(\frac{d}{dx} y(x) \right)$ to series expansion

$$x \cdot \left(\frac{d}{dx} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right)$ to series expansion

$$x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(r^2 - 5) x^r + a_1(r^2 + 2r - 4) x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k^2 + 2kr + r^2 - 5) + a_{k-2}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 - 5 = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{\sqrt{5}, -\sqrt{5}\}$$

- Each term must be 0
 $a_1(r^2 + 2r - 4) = 0$
- Solve for the dependent coefficient(s)
 $a_1 = 0$
- Each term in the series must be 0, giving the recursion relation
 $a_k(k^2 + 2kr + r^2 - 5) + a_{k-2} = 0$
- Shift index using $k \rightarrow k + 2$
 $a_{k+2}((k+2)^2 + 2(k+2)r + r^2 - 5) + a_k = 0$
- Recursion relation that defines series solution to ODE
 $a_{k+2} = -\frac{a_k}{k^2 + 2kr + r^2 + 4k + 4r - 1}$
- Recursion relation for $r = \sqrt{5}$
 $a_{k+2} = -\frac{a_k}{k^2 + 2k\sqrt{5} + 4 + 4k + 4\sqrt{5}}$
- Solution for $r = \sqrt{5}$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k+\sqrt{5}}, a_{k+2} = -\frac{a_k}{k^2 + 2k\sqrt{5} + 4 + 4k + 4\sqrt{5}}, a_1 = 0 \right]$$
- Recursion relation for $r = -\sqrt{5}$
 $a_{k+2} = -\frac{a_k}{k^2 - 2k\sqrt{5} + 4 + 4k - 4\sqrt{5}}$
- Solution for $r = -\sqrt{5}$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k-\sqrt{5}}, a_{k+2} = -\frac{a_k}{k^2 - 2k\sqrt{5} + 4 + 4k - 4\sqrt{5}}, a_1 = 0 \right]$$
- Combine solutions and rename parameters

$$\left[y(x) = \left(\sum_{k=0}^{\infty} a_k x^{k+\sqrt{5}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-\sqrt{5}} \right), a_{k+2} = -\frac{a_k}{k^2 + 2k\sqrt{5} + 4 + 4k + 4\sqrt{5}}, a_1 = 0, b_{k+2} = -\frac{b_k}{k^2 - 2k\sqrt{5} + 4 + 4k - 4\sqrt{5}} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

Maple dsolve solution

Solving time : 0.020 (sec)

Leaf size : 89

```
dsolve(x^2*diff(diff(y(x),x),x)+diff(y(x),x)*x+(x^2-5)*y(x) = 0,y(x),
series,x=0)
```

$$y = c_1 x^{-\sqrt{5}} \left(1 + \frac{1}{-4 + 4\sqrt{5}} x^2 + \frac{1}{32} \frac{1}{(-2 + \sqrt{5})(\sqrt{5} - 1)} x^4 + O(x^6) \right) \\ + c_2 x^{\sqrt{5}} \left(1 - \frac{1}{4 + 4\sqrt{5}} x^2 + \frac{1}{32} \frac{1}{(\sqrt{5} + 2)(\sqrt{5} + 1)} x^4 + O(x^6) \right)$$

Mathematica DSolve solution

Solving time : 0.007 (sec)

Leaf size : 210

```
AsymptoticDSolveValue[{x^2*D[y[x],{x,2}]+x*D[y[x],x]+(x^2-5)*y[x]==0,{}},
y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^4}{(-3 - \sqrt{5} + (1 - \sqrt{5})(2 - \sqrt{5}))(-1 - \sqrt{5} + (3 - \sqrt{5})(4 - \sqrt{5}))} \right. \\ \left. - \frac{x^2}{-3 - \sqrt{5} + (1 - \sqrt{5})(2 - \sqrt{5})} + 1 \right) x^{-\sqrt{5}} \\ + c_1 \left(\frac{x^4}{(-3 + \sqrt{5} + (1 + \sqrt{5})(2 + \sqrt{5}))(-1 + \sqrt{5} + (3 + \sqrt{5})(4 + \sqrt{5}))} \right. \\ \left. - \frac{x^2}{-3 + \sqrt{5} + (1 + \sqrt{5})(2 + \sqrt{5})} + 1 \right) x^{\sqrt{5}}$$

2.2.45 problem 45

Solved as second order Bessel ode	881
Solved as second order ode adjoint method	882
Maple step by step solution	884
Maple trace	885
Maple dsolve solution	885
Mathematica DSolve solution	886

Internal problem ID [8844]

Book : Second order enumerated odes

Section : section 2

Problem number : 45

Date solved : Thursday, December 12, 2024 at 09:53:50 AM

CAS classification : [_Bessel]

Solve

$$x^2 y'' + xy' + (x^2 - 5)y = 0$$

Solved as second order Bessel ode

Time used: 0.051 (sec)

Writing the ode as

$$x^2 y'' + xy' + (x^2 - 5)y = 0 \quad (1)$$

Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2)y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha)xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2)y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_3 \text{BesselJ}(n, \beta x^\gamma) + c_4 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= 0 \\ \beta &= 1 \\ n &= -\sqrt{5} \\ \gamma &= 1 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)$$

Solved as second order ode adjoint method

Time used: 0.700 (sec)

In normal form the ode

$$x^2 y'' + x y' + (x^2 - 5) y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = r(x) \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{1}{x} \\ q(x) &= \frac{x^2 - 5}{x^2} \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(\frac{\xi(x)}{x}\right)' + \left(\frac{(x^2 - 5)\xi(x)}{x^2}\right) &= 0 \\ \frac{\xi''(x)x^2 + \xi(x)x^2 - \xi'(x)x - 4\xi(x)}{x^2} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. Writing the ode as

$$\xi'' x^2 - \xi' x + (x^2 - 4) \xi = 0 \quad (1)$$

Bessel ode has the form

$$\xi'' x^2 + \xi' x + (-n^2 + x^2) \xi = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$\xi'' x^2 + (1 - 2\alpha) x \xi' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) \xi = 0 \quad (3)$$

With the standard solution

$$\xi = x^\alpha (c_3 \text{BesselJ}(n, \beta x^\gamma) + c_4 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned} \alpha &= 1 \\ \beta &= 1 \\ n &= -\sqrt{5} \\ \gamma &= 1 \end{aligned}$$

Substituting all the above into (4) gives the solution as

$$\xi = c_3 x \text{BesselJ}(-\sqrt{5}, x) + c_4 x \text{BesselY}(-\sqrt{5}, x)$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(\frac{1}{x} - \frac{c_3 \text{BesselJ}(-\sqrt{5}, x) + c_3 x \left(-\text{BesselJ}(-\sqrt{5} + 1, x) - \frac{\sqrt{5} \text{BesselJ}(-\sqrt{5}, x)}{x} \right) + c_4 \text{BesselY}(-\sqrt{5}, x)}{c_3 x \text{BesselJ}(-\sqrt{5}, x) + c_4 x \text{BesselY}(-\sqrt{5}, x)} \right)$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{-\text{BesselJ}(-\sqrt{5} + 1, x) c_3 x - \text{BesselY}(-\sqrt{5} + 1, x) c_4 x - \sqrt{5} (c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x))}{x (c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x))}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{-\text{BesselJ}(-\sqrt{5}+1,x)c_3x - \text{BesselY}(-\sqrt{5}+1,x)c_4x - \sqrt{5}(c_3\text{BesselJ}(-\sqrt{5},x) + c_4\text{BesselY}(-\sqrt{5},x))}{x(c_3\text{BesselJ}(-\sqrt{5},x) + c_4\text{BesselY}(-\sqrt{5},x))} dx} \\ &= \frac{1}{c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{y}{c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)} \right) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)} &= \int 0 dx + c_5 \\ &= c_5 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)}$ gives the final solution

$$y = \left(c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x) \right) c_5$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = \left(c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x) \right) c_5$$

The constants can be merged to give

$$y = c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_3 \text{BesselJ}(-\sqrt{5}, x) + c_4 \text{BesselY}(-\sqrt{5}, x)$$

Maple step by step solution

Let's solve

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) + x \left(\frac{d}{dx} y(x) \right) + (x^2 - 5) y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = -\frac{(x^2-5)y(x)}{x^2} - \frac{\frac{d}{dx} y(x)}{x}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dx^2} y(x) + \frac{\frac{d}{dx} y(x)}{x} + \frac{(x^2-5)y(x)}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{x^2-5}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -5$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) + x \left(\frac{d}{dx} y(x) \right) + (x^2 - 5) y(x) = 0$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y(x)$ to series expansion for $m = 0..2$

$$x^m \cdot y(x) = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k - > k - m$

$$x^m \cdot y(x) = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot \left(\frac{d}{dx} y(x) \right)$ to series expansion

$$x \cdot \left(\frac{d}{dx} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right)$ to series expansion

$$x^2 \cdot \left(\frac{d^2}{dx^2} y(x) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(r^2 - 5) x^r + a_1(r^2 + 2r - 4) x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k^2 + 2kr + r^2 - 5) + a_{k-2}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 - 5 = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{\sqrt{5}, -\sqrt{5}\}$$

- Each term must be 0
 $a_1(r^2 + 2r - 4) = 0$
- Solve for the dependent coefficient(s)
 $a_1 = 0$
- Each term in the series must be 0, giving the recursion relation
 $a_k(k^2 + 2kr + r^2 - 5) + a_{k-2} = 0$
- Shift index using $k \rightarrow k + 2$
 $a_{k+2}((k+2)^2 + 2(k+2)r + r^2 - 5) + a_k = 0$
- Recursion relation that defines series solution to ODE
 $a_{k+2} = -\frac{a_k}{k^2 + 2kr + r^2 + 4k + 4r - 1}$
- Recursion relation for $r = \sqrt{5}$
 $a_{k+2} = -\frac{a_k}{k^2 + 2k\sqrt{5} + 4 + 4k + 4\sqrt{5}}$
- Solution for $r = \sqrt{5}$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k+\sqrt{5}}, a_{k+2} = -\frac{a_k}{k^2 + 2k\sqrt{5} + 4 + 4k + 4\sqrt{5}}, a_1 = 0 \right]$$
- Recursion relation for $r = -\sqrt{5}$
 $a_{k+2} = -\frac{a_k}{k^2 - 2k\sqrt{5} + 4 + 4k - 4\sqrt{5}}$
- Solution for $r = -\sqrt{5}$

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^{k-\sqrt{5}}, a_{k+2} = -\frac{a_k}{k^2 - 2k\sqrt{5} + 4 + 4k - 4\sqrt{5}}, a_1 = 0 \right]$$
- Combine solutions and rename parameters

$$\left[y(x) = \left(\sum_{k=0}^{\infty} a_k x^{k+\sqrt{5}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-\sqrt{5}} \right), a_{k+2} = -\frac{a_k}{k^2 + 2k\sqrt{5} + 4 + 4k + 4\sqrt{5}}, a_1 = 0, b_{k+2} = -\frac{b_k}{k^2 - 2k\sqrt{5} + 4 + 4k - 4\sqrt{5}} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

Maple dsolve solution

Solving time : 0.005 (sec)

Leaf size : 19

```

dsolve(x^2*diff(diff(y(x),x),x)+diff(y(x),x)*x+(x^2-5)*y(x) = 0,
        y(x),singsol=all)

```

$$y = c_1 \text{BesselJ}(\sqrt{5}, x) + c_2 \text{BesselY}(\sqrt{5}, x)$$

Mathematica DSolve solution

Solving time : 0.133 (sec)

Leaf size : 26

```
DSolve[{x^2*D[y[x],{x,2}]+x*D[y[x],x]+(x^2-5)*y[x]==0,{}},  
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow c_1 \text{BesselJ}(\sqrt{5}, x) + c_2 \text{BesselY}(\sqrt{5}, x)$$

2.2.46 problem 46

Solved as second order Euler type ode	887
Solved as second order solved by an integrating factor	888
Solved as second order ode using change of variable on x method 2	889
Solved as second order ode using change of variable on x method 1	894
Solved as second order ode using change of variable on y method 1	895
Solved as second order ode using change of variable on y method 2	896
Solved as second order ode using Kovacic algorithm	898
Solved as second order ode adjoint method	900
Maple step by step solution	902
Maple trace	903
Maple dsolve solution	903
Mathematica DSolve solution	903

Internal problem ID [8845]

Book : Second order enumerated odes

Section : section 2

Problem number : 46

Date solved : Thursday, December 12, 2024 at 09:53:52 AM

CAS classification :

[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F(x)]']]

Solve

$$x^2y'' - 4xy' + 6y = 0$$

Solved as second order Euler type ode

Time used: 0.050 (sec)

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 4rx^{r-1} + 6x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 4rx^r + 6x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - 4r + 6 = 0$$

Or

$$r^2 - 5r + 6 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 2$$

$$r_2 = 3$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_2 x^3 + c_1 x^2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2 x^3 + c_1 x^2$$

Solved as second order solved by an integrating factor

Time used: 0.032 (sec)

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -\frac{4}{x}$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p \, dx} \\ &= e^{\int -\frac{4}{x} \, dx} \\ &= \frac{1}{x^2} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)y)'' &= 0 \\ \left(\frac{y}{x^2}\right)'' &= 0 \end{aligned}$$

Integrating once gives

$$\left(\frac{y}{x^2}\right)' = c_1$$

Integrating again gives

$$\left(\frac{y}{x^2}\right) = c_1 x + c_2$$

Hence the solution is

$$y = \frac{c_1 x + c_2}{\frac{1}{x^2}}$$

Or

$$y = c_1 x^3 + c_2 x^2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_1 x^3 + c_2 x^2$$

Solved as second order ode using change of variable on x method 2

Time used: 0.426 (sec)

In normal form the ode

$$x^2 y'' - 4xy' + 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{4}{x}$$

$$q(x) = \frac{6}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-\int p(x)dx} dx \\ &= \int e^{-\int -\frac{4}{x}dx} dx \\ &= \int e^{4\ln(x)} dx \\ &= \int x^4 dx \\ &= \frac{x^5}{5} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{6}{x^2}}{x^8} \\ &= \frac{6}{x^{10}} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{6y(\tau)}{x^{10}} &= 0 \end{aligned}$$

But in terms of τ

$$\frac{6}{x^{10}} = \frac{6}{25\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{6y(\tau)}{25\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. Writing the ode as

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{6y(\tau)}{25\tau^2} = 0 \quad (1)$$

$$A\frac{d^2}{d\tau^2}y(\tau) + B\frac{d}{d\tau}y(\tau) + Cy(\tau) = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= \frac{6}{25\tau^2} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(\tau) = y(\tau) e^{\int \frac{B}{2A} d\tau}$$

Then (2) becomes

$$z''(\tau) = rz(\tau) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-6}{25\tau^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -6 \\ t &= 25\tau^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(\tau) = \left(-\frac{6}{25\tau^2}\right) z(\tau) \quad (7)$$

Equation (7) is now solved. After finding $z(\tau)$ then $y(\tau)$ is found using the inverse transformation

$$y(\tau) = z(\tau) e^{-\int \frac{B}{2A} d\tau}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.105: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 25\tau^2$. There is a pole at $\tau = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{6}{25\tau^2}$$

For the pole at $\tau = 0$ let b be the coefficient of $\frac{1}{\tau^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{6}{25}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{5} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{2}{5} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{\tau^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{6}{25\tau^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{6}{25}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{5} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{2}{5} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{6}{25\tau^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{5}$	$\frac{2}{5}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{5}$	$\frac{2}{5}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{2}{5}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{2}{5} - \left(\frac{2}{5}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{\tau - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{\tau - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= \frac{2}{5\tau} + (-) (0) \\ &= \frac{2}{5\tau} \\ &= \frac{2}{5\tau} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(\tau)$ of degree $d = 0$ to solve the ode. The polynomial $p(\tau)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(\tau) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(\frac{2}{5\tau} \right) (0) + \left(\left(-\frac{2}{5\tau^2} \right) + \left(\frac{2}{5\tau} \right)^2 - \left(-\frac{6}{25\tau^2} \right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(\tau) &= p e^{\int \omega d\tau} \\ &= e^{\int \frac{2}{5\tau} d\tau} \\ &= \tau^{2/5} \end{aligned}$$

The first solution to the original ode in $y(\tau)$ is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} d\tau}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \tau^{2/5} \end{aligned}$$

Which simplifies to

$$y_1 = \tau^{2/5}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} d\tau}}{y_1^2} d\tau$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} d\tau \\ &= \tau^{2/5} \int \frac{1}{\tau^{4/5}} d\tau \\ &= \tau^{2/5} (5\tau^{1/5}) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y(\tau) &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\tau^{2/5}) + c_2 (\tau^{2/5} (5\tau^{1/5})) \end{aligned}$$

Will add steps showing solving for IC soon.

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 5^{3/5} (x^5)^{2/5}}{5} + c_2 5^{2/5} (x^5)^{3/5}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \frac{c_1 5^{3/5} (x^5)^{2/5}}{5} + c_2 5^{2/5} (x^5)^{3/5}$$

Solved as second order ode using change of variable on x method 1

Time used: 0.128 (sec)

In normal form the ode

$$x^2 y'' - 4xy' + 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{4}{x}$$

$$q(x) = \frac{6}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{\sqrt{6}\sqrt{\frac{1}{x^2}}}{c} \quad (6)$$

$$\tau'' = -\frac{\sqrt{6}}{c\sqrt{\frac{1}{x^2}}x^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$

$$= \frac{-\frac{\sqrt{6}}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{4}{x}\frac{\sqrt{6}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{6}\sqrt{\frac{1}{x^2}}}{c}\right)^2}$$

$$= -\frac{5c\sqrt{6}}{6}$$

Therefore ode (3) now becomes

$$y(\tau)'' + p_1y(\tau)' + q_1y(\tau) = 0$$

$$\frac{d^2}{d\tau^2}y(\tau) - \frac{5c\sqrt{6}}{6}\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) = 0 \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{\frac{5\sqrt{6}c\tau}{12}} \left(c_1 \cosh\left(\frac{\sqrt{6}c\tau}{12}\right) + ic_2 \sinh\left(\frac{\sqrt{6}c\tau}{12}\right) \right)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{6} \sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{6} \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = x^{5/2} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^{5/2} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right)$$

Solved as second order ode using change of variable on y method 1

Time used: 0.128 (sec)

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$\begin{aligned}p(x) &= -\frac{4}{x} \\ q(x) &= \frac{6}{x^2}\end{aligned}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned}Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= \frac{6}{x^2} - \frac{\left(-\frac{4}{x}\right)'}{2} - \frac{\left(-\frac{4}{x}\right)^2}{4} \\ &= \frac{6}{x^2} - \frac{\left(\frac{4}{x^2}\right)}{2} - \frac{\left(\frac{16}{x^2}\right)}{4} \\ &= \frac{6}{x^2} - \left(\frac{2}{x^2}\right) - \frac{4}{x^2} \\ &= 0\end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned}z(x) &= e^{-\int \frac{p(x)}{2} dx} \\ &= e^{-\int \frac{-4}{x} dx} \\ &= x^2\end{aligned} \quad (5)$$

Hence (3) becomes

$$y = v(x) x^2 \quad (4)$$

Applying this change of variable to the original ode results in

$$x^4 v''(x) = 0$$

Which is now solved for $v(x)$.

The above ode can be simplified to

$$v''(x) = 0$$

Integrating twice gives the solution

$$v(x) = c_1 x + c_2$$

Will add steps showing solving for IC soon.

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 x + c_2) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = x^2$$

Hence (7) becomes

$$y = (c_1 x + c_2) x^2$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = (c_1 x + c_2) x^2$$

Solved as second order ode using change of variable on y method 2

Time used: 0.087 (sec)

In normal form the ode

$$x^2 y'' - 4xy' + 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= -\frac{4}{x} \\ q(x) &= \frac{6}{x^2} \end{aligned}$$

Applying change of variables on the dependent variable $y = v(x) x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p \right) v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q \right) v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{4n}{x^2} + \frac{6}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 3 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{2v'(x)}{x} &= 0 \\ v''(x) + \frac{2v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{2u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form a linear first order is

$$u'(x) + q(x)u(x) = p(x)$$

Comparing the above to the given ode shows that

$$\begin{aligned} q(x) &= \frac{2}{x} \\ p(x) &= 0 \end{aligned}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q \, dx} \\ &= e^{\int \frac{2}{x} \, dx} \\ &= x^2 \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx} \mu u &= 0 \\ \frac{d}{dx} (u x^2) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned} u x^2 &= \int 0 \, dx + c_1 \\ &= c_1 \end{aligned}$$

Dividing throughout by the integrating factor x^2 gives the final solution

$$u(x) = \frac{c_1}{x^2}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= -\frac{c_1}{x} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{c_1}{x} + c_2\right) x^3 \\&= (c_2 x - c_1) x^2\end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = \left(-\frac{c_1}{x} + c_2\right) x^3$$

Solved as second order ode using Kovacic algorithm

Time used: 0.104 (sec)

Writing the ode as

$$x^2 y'' - 4xy' + 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\B &= -4x \\C &= 6\end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 0 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.106: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-4x}{x^2} dx} \\ &= z_1 e^{2 \ln(x)} \\ &= z_1 (x^2) \end{aligned}$$

Which simplifies to

$$y_1 = x^2$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{4x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{4\ln(x)}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(x^2) + c_2(x^2(x)) \end{aligned}$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = c_2 x^3 + c_1 x^2$$

Solved as second order ode adjoint method

Time used: 0.132 (sec)

In normal form the ode

$$x^2 y'' - 4xy' + 6y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = r(x) \tag{2}$$

Where

$$\begin{aligned} p(x) &= -\frac{4}{x} \\ q(x) &= \frac{6}{x^2} \\ r(x) &= 0 \end{aligned}$$

The Lagrange adjoint ode is given by

$$\begin{aligned} \xi'' - (\xi p)' + \xi q &= 0 \\ \xi'' - \left(-\frac{4\xi(x)}{x}\right)' + \left(\frac{6\xi(x)}{x^2}\right) &= 0 \\ \xi''(x) + \frac{2\xi(x)}{x^2} + \frac{4\xi'(x)}{x} &= 0 \end{aligned}$$

Which is solved for $\xi(x)$. This is Euler second order ODE. Let the solution be $\xi = x^r$, then $\xi' = rx^{r-1}$ and $\xi'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + 4rx^{r-1} + 2x^r = 0$$

Simplifying gives

$$r(r-1)x^r + 4rx^r + 2x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + 4r + 2 = 0$$

Or

$$r^2 + 3r + 2 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -2$$

$$r_2 = -1$$

Since the roots are real and distinct, then the general solution is

$$\xi = c_1 \xi_1 + c_2 \xi_2$$

Where $\xi_1 = x^{r_1}$ and $\xi_2 = x^{r_2}$. Hence

$$\xi = \frac{c_1}{x^2} + \frac{c_2}{x}$$

Will add steps showing solving for IC soon.

The original ode now reduces to first order ode

$$\begin{aligned} \xi(x) y' - y \xi'(x) + \xi(x) p(x) y &= \int \xi(x) r(x) dx \\ y' + y \left(p(x) - \frac{\xi'(x)}{\xi(x)} \right) &= \frac{\int \xi(x) r(x) dx}{\xi(x)} \end{aligned}$$

Or

$$y' + y \left(-\frac{4}{x} - \frac{-\frac{2c_1}{x^3} - \frac{c_2}{x^2}}{\frac{c_1}{x^2} + \frac{c_2}{x}} \right) = 0$$

Which is now a first order ode. This is now solved for y . In canonical form a linear first order is

$$y' + q(x)y = p(x)$$

Comparing the above to the given ode shows that

$$q(x) = -\frac{3c_2x + 2c_1}{x(c_2x + c_1)}$$

$$p(x) = 0$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int q dx} \\ &= e^{\int -\frac{3c_2x + 2c_1}{x(c_2x + c_1)} dx} \\ &= \frac{1}{(c_2x + c_1)x^2} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} \left(\frac{y}{(c_2x + c_1)x^2} \right) &= 0 \end{aligned}$$

Integrating gives

$$\begin{aligned} \frac{y}{(c_2x + c_1)x^2} &= \int 0 dx + c_3 \\ &= c_3 \end{aligned}$$

Dividing throughout by the integrating factor $\frac{1}{(c_2x+c_1)x^2}$ gives the final solution

$$y = (c_2x + c_1)x^2c_3$$

Hence, the solution found using Lagrange adjoint equation method is

$$y = (c_2x + c_1)x^2c_3$$

The constants can be merged to give

$$y = x^2(c_2x + c_1)$$

Will add steps showing solving for IC soon.

Summary of solutions found

$$y = x^2(c_2x + c_1)$$

Maple step by step solution

Let's solve

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) - 4x \left(\frac{d}{dx} y(x) \right) + 6y(x) = 0$$

- Highest derivative means the order of the ODE is 2

$$\frac{d^2}{dx^2} y(x)$$

- Isolate 2nd derivative

$$\frac{d^2}{dx^2} y(x) = -\frac{6y(x)}{x^2} + \frac{4 \left(\frac{d}{dx} y(x) \right)}{x}$$

- Group terms with $y(x)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dx^2} y(x) - \frac{4 \left(\frac{d}{dx} y(x) \right)}{x} + \frac{6y(x)}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 \left(\frac{d^2}{dx^2} y(x) \right) - 4x \left(\frac{d}{dx} y(x) \right) + 6y(x) = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x, using the chain rule

$$\frac{d}{dx} y(x) = \left(\frac{d}{dt} y(t) \right) \left(\frac{d}{dx} t(x) \right)$$

- Compute derivative

$$\frac{d}{dx} y(x) = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x, using the chain rule

$$\frac{d^2}{dx^2} y(x) = \left(\frac{d^2}{dt^2} y(t) \right) \left(\frac{d}{dx} t(x) \right)^2 + \left(\frac{d^2}{dx^2} t(x) \right) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$\frac{d^2}{dx^2} y(x) = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - 4 \frac{d}{dt} y(t) + 6y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - 5 \frac{d}{dt} y(t) + 6y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 5r + 6 = 0$$

- Factor the characteristic polynomial
 $(r - 2)(r - 3) = 0$
- Roots of the characteristic polynomial
 $r = (2, 3)$
- 1st solution of the ODE
 $y_1(t) = e^{2t}$
- 2nd solution of the ODE
 $y_2(t) = e^{3t}$
- General solution of the ODE
 $y(t) = C_1 y_1(t) + C_2 y_2(t)$
- Substitute in solutions
 $y(t) = C_1 e^{2t} + C_2 e^{3t}$
- Change variables back using $t = \ln(x)$
 $y(x) = C_2 x^3 + C_1 x^2$
- Simplify
 $y(x) = x^2(C_2 x + C_1)$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`
```

Maple dsolve solution

Solving time : 0.006 (sec)
Leaf size : 13

```
dsolve(x^2*diff(diff(y(x),x),x)-4*diff(y(x),x)*x+6*y(x) = 0,
y(x),singsol=all)
```

$$y = x^2(c_1 x + c_2)$$

Mathematica DSolve solution

Solving time : 0.016 (sec)
Leaf size : 16

```
DSolve[{x^2*D[y[x],{x,2}]-4*x*D[y[x],x]+6*y[x]==0,{}},
y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow x^2(c_2 x + c_1)$$

2.2.47 problem 47

Maple step by step solution	904
Maple trace	905
Maple dsolve solution	905
Mathematica DSolve solution	905

Internal problem ID [8846]

Book : Second order enumerated odes

Section : section 2

Problem number : 47

Date solved : Wednesday, December 18, 2024 at 02:16:41 AM

CAS classification : [[_3rd_order, _with_linear_symmetries]]

Solve

$$y''' - xy = 0$$

Maple step by step solution

Let's solve

$$\frac{d^3}{dx^3}y(x) - xy(x) = 0$$

- Highest derivative means the order of the ODE is 3

$$\frac{d^3}{dx^3}y(x)$$

- Assume series solution for $y(x)$

$$y(x) = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y(x)$ to series expansion

$$x \cdot y(x) = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y(x) = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert $\frac{d^3}{dx^3}y(x)$ to series expansion

$$\frac{d^3}{dx^3}y(x) = \sum_{k=3}^{\infty} a_k k(k-1)(k-2) x^{k-3}$$

- Shift index using $k \rightarrow k + 3$

$$\frac{d^3}{dx^3}y(x) = \sum_{k=0}^{\infty} a_{k+3} (k+3)(k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$6a_3 + \left(\sum_{k=1}^{\infty} (a_{k+3}(k+3)(k+2)(k+1) - a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$6a_3 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^3 + 6k^2 + 11k + 6) a_{k+3} - a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^3 + 6(k+1)^2 + 11k + 17) a_{k+4} - a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y(x) = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = \frac{a_k}{k^3 + 9k^2 + 26k + 24}, 6a_3 = 0 \right]$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying high order exact linear fully integrable
trying to convert to a linear ODE with constant coefficients
trying differential order: 3; missing the dependent variable
trying Louvillian solutions for 3rd order ODEs, imprimitive case
-> pFq: Equivalence to the 3F2 or one of its 3 confluent cases under a power @ Moebius
<- pFq successful: received ODE is equivalent to the 0F2 ODE, case c = 0`

```

Maple dsolve solution

Solving time : 0.013 (sec)

Leaf size : 45

```

dsolve(diff(diff(diff(y(x),x),x),x)-x*y(x) = 0,
        y(x),singsol=all)

```

$$\begin{aligned}
 y = & c_1 \operatorname{hypergeom} \left(\left[\right], \left[\frac{1}{2}, \frac{3}{4} \right], \frac{x^4}{64} \right) + c_2 x \operatorname{hypergeom} \left(\left[\right], \left[\frac{3}{4}, \frac{5}{4} \right], \frac{x^4}{64} \right) \\
 & + c_3 x^2 \operatorname{hypergeom} \left(\left[\right], \left[\frac{5}{4}, \frac{3}{2} \right], \frac{x^4}{64} \right)
 \end{aligned}$$

Mathematica DSolve solution

Solving time : 0.012 (sec)

Leaf size : 76

```

DSolve[{D[y[x],{x,3}]-x*y[x]==0,{x}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow c_1 {}_0F_2 \left(; \frac{1}{2}, \frac{3}{4}; \frac{x^4}{64} \right) + \frac{1}{8} x \left((2 + 2i) c_2 {}_0F_2 \left(; \frac{3}{4}, \frac{5}{4}; \frac{x^4}{64} \right) + i c_3 x {}_0F_2 \left(; \frac{5}{4}, \frac{3}{2}; \frac{x^4}{64} \right) \right)$$

2.2.48 problem 48

Existence and uniqueness analysis	906
Solved as first order autonomous ode	907
Solved as first order Bernoulli ode	908
Solved as first order Exact ode	910
Solved using Lie symmetry for first order ode	913
Maple step by step solution	916
Maple trace	917
Maple dsolve solution	917
Mathematica DSolve solution	917

Internal problem ID [8847]

Book : Second order enumerated odes

Section : section 2

Problem number : 48

Date solved : Tuesday, December 17, 2024 at 01:04:33 PM

CAS classification : [_quadrature]

Solve

$$y' = y^{1/3}$$

With initial conditions

$$y(0) = 0$$

Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= y^{1/3} \end{aligned}$$

The y domain of $f(x, y)$ when $x = 0$ is

$$\{0 \leq y\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned} \frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(y^{1/3}) \\ &= \frac{1}{3y^{2/3}} \end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{0 < y\}$$

But the point $y_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. Solution exists but no guarantee that unique solution exists.

Solved as first order autonomous ode

Time used: 0.087 (sec)

Since the ode has the form $y' = f(y)$ and initial conditions (x_0, y_0) are given such that they satisfy the ode itself, then we can write

$$0 = f(y)|_{y=y_0}$$

$$0 = 0$$

And the solution is immediately written as

$$y = y_0$$

$$y = 0$$

Singular solutions are found by solving

$$y^{1/3} = 0$$

for y . This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$y = 0$$

The following diagram is the phase line diagram. It classifies each of the above equilibrium points as stable or not stable or semi-stable.






Figure 2.164: Phase line diagram

(a) Solution plot
 $y = 0$ (b) Slope field plot
 $y' = y^{1/3}$ Summary of solutions found

$$y = 0$$

Solved as first order Bernoulli ode

Time used: 0.312 (sec)

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= y^{1/3} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = (1) y^{1/3} \tag{1}$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \tag{2}$$

Comparing this to (1) shows that

$$\begin{aligned} f_0 &= 0 \\ f_1 &= 1 \end{aligned}$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_1(x) \tag{3}$$

The next step is use the substitution $v = y^{1-n}$ in equation (3) which generates a new ODE in $v(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= 0 \\ f_1(x) &= 1 \\ n &= \frac{1}{3} \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^{1/3}$ gives

$$y' \frac{1}{y^{1/3}} = 0 + 1 \tag{4}$$

Let

$$\begin{aligned} v &= y^{1-n} \\ &= y^{2/3} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$v' = \frac{2}{3y^{1/3}}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} \frac{3v'(x)}{2} &= 1 \\ v' &= \frac{2}{3} \end{aligned} \quad (7)$$

The above now is a linear ODE in $v(x)$ which is now solved.

Since the ode has the form $v'(x) = f(x)$, then we only need to integrate $f(x)$.

$$\begin{aligned} \int dv &= \int \frac{2}{3} dx \\ v(x) &= \frac{2x}{3} + c_1 \end{aligned}$$

The substitution $v = y^{1-n}$ is now used to convert the above solution back to y which results in


$$y^{2/3} = \frac{2x}{3} + c_1$$

Solving for the constant of integration from initial conditions, the solution becomes


$$y^{2/3} = \frac{2x}{3}$$

Solving for y gives

$$y = \frac{2\sqrt{2}\sqrt{3}x^{3/2}}{9}$$



(a) Solution plot
 $y = \frac{2\sqrt{2}\sqrt{3}x^{3/2}}{9}$



(b) Slope field plot
 $y' = y^{1/3}$

Summary of solutions found

$$y = \frac{2\sqrt{2}\sqrt{3}x^{3/2}}{9}$$

Solved as first order Exact ode

Time used: 0.152 (sec)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= (y^{1/3}) dx \\ (-y^{1/3}) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -y^{1/3} \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (-y^{1/3}) \\ &= -\frac{1}{3y^{2/3}} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1 \left(\left(-\frac{1}{3y^{2/3}} \right) - (0) \right) \\ &= -\frac{1}{3y^{2/3}} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= -\frac{1}{y^{1/3}} \left((0) - \left(-\frac{1}{3y^{2/3}} \right) \right) \\ &= -\frac{1}{3y} \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{1}{3y} \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-\frac{\ln(y)}{3}} \\ &= \frac{1}{y^{1/3}} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{y^{1/3}} (-y^{1/3}) \\ &= -1 \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{y^{1/3}} (1) \\ &= \frac{1}{y^{1/3}} \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (-1) + \left(\frac{1}{y^{1/3}} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -1 dx \\ \phi &= -x + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y^{1/3}}$. Therefore equation (4) becomes

$$\frac{1}{y^{1/3}} = 0 + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y^{1/3}}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y^{1/3}}\right) dy \\ f(y) &= \frac{3y^{2/3}}{2} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -x + \frac{3y^{2/3}}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into the constant c_1 gives the solution as


$$c_1 = -x + \frac{3y^{2/3}}{2}$$

Solving for the constant of integration from initial conditions, the solution becomes

$$-x + \frac{3y^{2/3}}{2} = 0$$

Solving for y gives

$$y = \frac{2\sqrt{2}\sqrt{3}x^{3/2}}{9}$$



(a) Solution plot
 $y = \frac{2\sqrt{2}\sqrt{3}x^{3/2}}{9}$

(b) Slope field plot
 $y' = y^{1/3}$

Summary of solutions found

$$y = \frac{2\sqrt{2}\sqrt{3}x^{3/2}}{9}$$

Solved using Lie symmetry for first order ode

Time used: 0.495 (sec)

Writing the ode as

$$y' = y^{1/3}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as ansatz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + y^{1/3}(b_3 - a_2) - y^{2/3}a_3 - \frac{xb_2 + yb_3 + b_1}{3y^{2/3}} = 0 \quad (\text{5E})$$

Putting the above in normal form gives

$$-\frac{3y^{4/3}a_3 + 3ya_2 - 2yb_3 - 3b_2y^{2/3} + xb_2 + b_1}{3y^{2/3}} = 0$$

Setting the numerator to zero gives

$$-3y^{4/3}a_3 + 3b_2y^{2/3} - xb_2 - 3ya_2 + 2yb_3 - b_1 = 0 \quad (\text{6E})$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, y^{2/3}, y^{4/3}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, y^{2/3} = v_3, y^{4/3} = v_4\}$$

The above PDE (6E) now becomes

$$-3v_2a_2 - 3v_4a_3 - v_1b_2 + 3b_2v_3 + 2v_2b_3 - b_1 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$-v_1b_2 + (-3a_2 + 2b_3)v_2 + 3b_2v_3 - 3v_4a_3 - b_1 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -3a_3 &= 0 \\ -b_1 &= 0 \\ -b_2 &= 0 \\ 3b_2 &= 0 \\ -3a_2 + 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= a_1 \\ a_2 &= a_2 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= \frac{3a_2}{2} \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 1 \\ \eta &= 0 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 0 - (y^{1/3}) (1) \\ &= -y^{1/3} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-y^{1/3}} dy \end{aligned}$$

Which results in

$$S = -\frac{3y^{2/3}}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y^{1/3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= 0 \\ S_y &= -\frac{1}{y^{1/3}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -1 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S .

Since the ode has the form $\frac{d}{dR}S(R) = f(R)$, then we only need to integrate $f(R)$.

$$\begin{aligned} \int dS &= \int -1 dR \\ S(R) &= -R + c_2 \end{aligned}$$



To complete the solution, we just need to transform the above back to x, y coordinates. This results in

$$-\frac{3y^{2/3}}{2} = -x + c_2$$

Which gives


$$y = \frac{(6x - 6c_2)^{3/2}}{27}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.


Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y^{1/3}$ 	$R = x$ $S = -\frac{3y^{2/3}}{2}$	$\frac{dS}{dR} = -1$ 

Solving for the constant of integration from initial conditions, the solution becomes

$$y = \frac{2x^{3/2}\sqrt{6}}{9}$$



(a) Solution plot
 $y = \frac{2x^{3/2}\sqrt{6}}{9}$



(b) Slope field plot
 $y' = y^{1/3}$

Summary of solutions found

$$y = \frac{2x^{3/2}\sqrt{6}}{9}$$

Maple step by step solution

Let's solve

$$\left[\frac{d}{dx}y(x) = y(x)^{1/3}, y(0) = 0 \right]$$

- Highest derivative means the order of the ODE is 1
- $\frac{d}{dx}y(x)$
- Solve for the highest derivative
- $\frac{d}{dx}y(x) = y(x)^{1/3}$
- Separate variables

$$\frac{\frac{d}{dx}y(x)}{y(x)^{1/3}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{\frac{d}{dx}y(x)}{y(x)^{1/3}} dx = \int 1 dx + C1$$

- Evaluate integral

$$\frac{3y(x)^{2/3}}{2} = x + C1$$

- Solve for $y(x)$

$$y(x) = \frac{(6x+6C1)^{3/2}}{27}$$

- Use initial condition $y(0) = 0$

$$0 = \frac{2\sqrt{6} C1^{3/2}}{9}$$

- Solve for $C1$

$$C1 = 0$$

- Substitute $C1 = 0$ into general solution and simplify

$$y(x) = \frac{2\sqrt{6}x^{3/2}}{9}$$

- Solution to the IVP

$$y(x) = \frac{2\sqrt{6}x^{3/2}}{9}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

Maple dsolve solution

Solving time : 0.005 (sec)

Leaf size : 5

```

dsolve([diff(y(x),x) = y(x)^(1/3),
        op([y(0) = 0])),y(x),singsol=all)

```

$$y = 0$$

Mathematica DSolve solution

Solving time : 0.006 (sec)

Leaf size : 21

```

DSolve[{D[y[x],x]==y[x]^(1/3),{y[0]==0}},
        y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow \frac{2}{3} \sqrt{\frac{2}{3}} x^{3/2}$$

2.2.49 problem 49

Solution using Matrix exponential method	918
Solution using explicit Eigenvalue and Eigenvector method	919
Maple step by step solution	922
Maple dsolve solution	923
Mathematica DSolve solution	923

Internal problem ID [8848]

Book : Second order enumerated odes

Section : section 2

Problem number : 49

Date solved : Thursday, December 12, 2024 at 09:53:56 AM

CAS classification : system_of_ODEs

$$\begin{aligned}x' &= 3x + y \\y' &= -x + y\end{aligned}$$

Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{2t}(1+t) & t e^{2t} \\ -t e^{2t} & e^{2t}(1-t) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} e^{2t}(1+t) & t e^{2t} \\ -t e^{2t} & e^{2t}(1-t) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} e^{2t}(1+t)c_1 + t e^{2t}c_2 \\ -t e^{2t}c_1 + e^{2t}(1-t)c_2 \end{bmatrix} \\ &= \begin{bmatrix} e^{2t}(tc_1 + c_2t + c_1) \\ -((-1+t)c_2 + tc_1)e^{2t} \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 3 - \lambda & 1 \\ -1 & 1 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 4\lambda + 4 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 2$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
2	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\begin{aligned} \left(\begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} \end{aligned}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 1 & 0 \\ -1 & -1 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} v_1 \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} v_1 \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$


Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} v_1 \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
2	2	1	Yes	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 2 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram



This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 2. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{2t} \\ &= \begin{bmatrix} -e^{2t} \\ e^{2t} \end{bmatrix} \end{aligned}$$

And

$$\begin{aligned} \vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} -1 \\ 1 \end{bmatrix} t + \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right) e^{2t} \\ &= \begin{bmatrix} -e^{2t}(t+2) \\ e^{2t}(1+t) \end{bmatrix} \end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x \\ y \end{bmatrix} = c_1 \begin{bmatrix} -e^{2t} \\ e^{2t} \end{bmatrix} + c_2 \begin{bmatrix} e^{2t}(-t-2) \\ e^{2t}(1+t) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -((t+2)c_2 + c_1) e^{2t} \\ e^{2t}(c_2 t + c_1 + c_2) \end{bmatrix}$$




Figure 2.170: Phase plot

Maple step by step solution

Let's solve

$$\left[\frac{d}{dt}x(t) = 3x(t) + y(t), \frac{d}{dt}y(t) = -x(t) + y(t) \right]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\frac{d}{dt}\vec{x}(t) = \begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\frac{d}{dt}\vec{x}(t) = \begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt}\vec{x}(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[2, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[2, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 2

$$\vec{x}_1(t) = e^{2t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 2$ is the eigenvalue, and \vec{v} is

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 2

$$\left(\begin{bmatrix} 3 & 1 \\ -1 & 1 \end{bmatrix} - 2 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 2

$$\vec{x}_2(t) = e^{2t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = C1 \vec{x}_1(t) + C2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = C1 e^{2t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + C2 e^{2t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{2t}(-C2t - C1 - C2) \\ e^{2t}(C2t + C1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = e^{2t}(-C2t - C1 - C2), y(t) = e^{2t}(C2t + C1)\}$$

Maple dsolve solution

Solving time : 0.038 (sec)

Leaf size : 31

```
dsolve([diff(x(t),t) = 3*x(t)+y(t), diff(y(t),t) = -x(t)+y(t)]
, {op([x(t), y(t)])})
```

$$\begin{aligned} x(t) &= e^{2t}(c_2t + c_1) \\ y(t) &= -e^{2t}(c_2t + c_1 - c_2) \end{aligned}$$

Mathematica DSolve solution

Solving time : 0.004 (sec)

Leaf size : 42

```
DSolve[{D[x[t],t]==3*x[t]+y[t],D[y[t],t]==-x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions->True]
```

$$\begin{aligned} x(t) &\rightarrow e^{2t}(c_1(t+1) + c_2t) \\ y(t) &\rightarrow e^{2t}(c_2 - (c_1 + c_2)t) \end{aligned}$$